

SCIENTIFIC AMERICAN

SEPTEMBER 1992

\$4.95

**SPECIAL
ISSUE**

MIND AND BRAIN

SEX

VISION

MEMORY


LEARNING

LANGUAGE

DISORDERS

DEVELOPMENT

CONSCIOUSNESS




The Cradle Of

First came very crude hand tools. Then cave drawings. Villages. An alphabet. And finally, two million years after the first *Homo sapiens* stood up straight, his descendants can now sit down.



Guess where.


The elaborately crafted driver's seat in the Lexus ES300 sports sedan offers 4-way power adjust-



ments. And, beneath the optional fine-grain leather upholstery, both the firmness of the springs and the densi-

ty of the padding are varied for additional comfort and greater lateral support for those spirited maneuvers

Civilization.



The ES 300

materials are extensively used to provide the hallmark Lexus quiet. And a 180-watt maximum power, 8-speaker premium sound system is furnished to subsequently shatter it.

Indeed, all that is currently missing inside the cabin is you. We suggest a visit to your Lexus dealer, where the two of you can further discuss the rise and recline of mankind.




The Relentless Pursuit Of Perfection.

through tight turns (we are, after all, talking about a sports sedan).

the cabin surrounding it. In this particular case, it is a Lexus cabin:

But a seat is only as good as noise- and vibration-dampening

Can we arrest t robs people o



he disease that f their minds?

A photograph of a person wearing a yellow short-sleeved shirt, sitting at a dark wooden table. They are looking out a window with multiple panes, which shows some greenery outside. The lighting is warm, suggesting indoor light or late afternoon sun.

Over 4 million people suffer from it, and nearly half of all nursing home patients are its victims.

It's Alzheimer's, the disease that steals a person's mind, dignity and independence, and exacts costs of over \$88 billion a year for institutional and home care.

Leading the way in the search for relief from this mysterious disease is the pharmaceutical industry, which is making the nation's largest investment in drug research.

Since 1988 alone, the industry has been conducting research on over 40 medicines for Alzheimer's, and currently has 13 in test. While these efforts hold hope for breakthroughs, the process is long and difficult, with only a few of the thousands of compounds developed ever achieving success.

This exhaustive, high risk research increases the industry's cost of doing business, and in turn the price of drugs. But it also leads to the kind of discoveries that can break the grip of an enormously costly disease like Alzheimer's.

To learn more about pharmaceutical research, and the critical role it plays in health care, call or write for our free booklet, "Good Medicine."

The Pharmaceutical Manufacturers Association, 1100 15th St, NW, Box I, Wash., DC 20005. 1-800-538-2692.

PHARMACEUTICALS
Good Medicine For Containing Health Costs.

48



Mind and Brain *Gerald D. Fischbach*

The human brain is the most complex structure in the known universe. Genes and experience have jointly shaped its machinery; its design is the result of millions of years of evolution. Our survival depends on a deeper understanding of the marvelous biochemical happening that arises from it: the mind.

60



The Developing Brain *Carla J. Shatz*

Remarkably precise connections between more than 100 billion neurons account for all the properties of the mind. Yet in the fetus this intricate wiring is only an approximation of that in the mature brain. The final linkages are shaped by stimulation of the newborn through such sensory experiences as touch, speech and images.


68



The Visual Image in Mind and Brain *Semir Zeki*

There is a great deal more to vision than light meeting the eye. The world we see is literally an invention of the brain, actively constructed from a constantly changing flood of information. Seeing and understanding occur simultaneously through the synchronized activities of specialized areas in the visual cortex.

78



The Biological Basis of Learning and Individuality *Eric R. Kandel and Robert D. Hawkins*

Learning and memory—the acquisition of knowledge and the retention of that information over time—are being studied on the cellular and molecular level. These processes, which connect us to our past and are the key to individuality, engage a simple set of rules to strengthen connections between nerve cells.

88



Brain and Language *Antonio R. Damasio and Hanna Damasio*

In the beginning, there were no words. Language arose and persisted because it is a superb means of communication. Three sets of interacting neural structures process language in the brain. One supports nonlanguage concepts, another assembles words and sentences, and a third mediates between the first two.

110




Working Memory and the Mind *Patricia S. Goldman-Rakic*

Working memory has been called the blackboard of the mind. Its ability to combine moment-to-moment awareness and instantaneous retrieval of archived information is fundamental to language comprehension, learning and reason. Experiments with monkeys are identifying the key neural structures.

118 **Sex Differences in the Brain**

Doreen Kimura



The different ways in which men and women approach intellectual problem solving have often been attributed to variations in experience. Evidence indicates that reproductive hormones alter brain function permanently during fetal development. Further research may explain the evolutionary significance of these changes.

126 **Major Disorders of Mind and Brain**


Elliot S. Gershon and Ronald O. Rieder



Schizophrenia and mania—the most devastating maladies of the mind—each afflict 1 percent of the population. Impressive advances in neuroscience and genetics are revealing the anatomic, biochemical and hereditary bases of these disorders. Research has already shaped the development of new therapies.

134 **Aging Brain, Aging Mind**


Dennis J. Selkoe



As we age, the neural structures involved in learning, memory and reasoning undergo a number of physical changes. Yet such alterations do not necessarily signal an inevitable slow march to mindlessness and mortality. Elderly individuals who remain in good health show only a subtle decline in cognitive function.

144 **How Neural Networks Learn from Experience**


Geoffrey E. Hinton



Networks of artificial neurons modeled on conventional computers are helping explain the ability of the brain to process and retain information. These neural-network simulations have already ruled out many theories. They are now beginning to reveal how the brain accomplishes the remarkable feat of learning.

152 **The Problem of Consciousness**

Francis Crick and Christof Koch



Can consciousness, the most profound and puzzling facet of the mind, be probed experimentally? Are elusive mental events explainable as the behavior of interacting neurons? The authors argue that existing evidence already provides a glimpse of the nature of visual consciousness that can guide experimenters.

DEPARTMENTS

18 **Science and the Citizen**

Quantum gravity.... Population politics.... Gold bugs.... The evidence for dark matter.... First steps for artificial life.... PROFILE: SSC architect Roy F. Schwitters.

160 **Science and Business**

Ag biotech moves out of the lab and into the kitchen.... Virtual reality meets the real world.... THE ANALYTICAL ECONOMIST: Options to the tyranny of economic growth.


12 **Letters to the Editors**

16 **50 and 100 Years Ago**


170 **The Amateur Scientist**

174 **Book Reviews**

180 **Essay: Jonathan Miller**



NO ONE TRIES TO UNDERSTAND YOU MORE THAN WE DO.



It doesn't take the mind of a brain surgeon to understand people want to live longer, healthier lives. That's why Toshiba builds Magnetic Resonance Imagers that allow physicians to see deep within the human body without X-rays or exploratory surgery. Even abnormalities in cell structure, like cancer, can be discovered.

But, it did take the mind of a brain surgeon, as well as other medical experts at our research and development center in San Francisco, to help us design an even better MRI—our new MRT-150A. It's just one example of our pioneering role in super-conductive magnet technology and computer-image processing.

It's also an example of how much Toshiba cares about making the world a healthier place to live.

In Touch with Tomorrow
TOSHIBA

For more information write: Toshiba America Medical Systems, Inc., 2441 Michelle Drive, Tustin, CA 92681 or call: 1-800-421-1968.

© 1992 SCIENTIFIC AMERICAN, INC

THE ILLUSTRATIONS

THE COVER painting is by Carol Donner.

Small drawing on the opening page of each article is by Patricia J. Wynne.

Page	Source	Page	Source
49	<i>Hide-and-Seek</i> (1940-42), by Pavel Tchelitchew, courtesy of the Museum of Modern Art, N.Y.; Mrs. Simon Guggenheim Fund	94	Marc Skinner
50	Janet Robbins, Harvard Medical School	95	Patricia J. Wynne
51	Carol Donner	110	Eric Hartman/Magnum
52	Tomo Narashima	112-113	Patricia J. Wynne
53	Patricia J. Wynne; courtesy of Clement Fox, Wayne State University, © Williams & Wilkins Co.	114	Patricia J. Wynne (<i>top</i>); Harriett Friedman and Patricia S. Goldman-Rakic (<i>bottom</i>)
54-55	Marcus E. Raichle, Washington University School of Medicine	115-116	Patricia J. Wynne
56	Apostolos P. Georgopoulos; © Society for Neuroscience	118	<i>The Old Oaken Bucket</i> , by Grandma Moses, © 1987 Grandma Moses Properties Co., N.Y.
60	<i>Behold Man</i> , by Lennart Nilsson, © 1973 Little, Brown and Company	120-121	Jared Schneidman
62-63	Dana Burns-Pizer (<i>top</i>), Carla J. Shatz (<i>bottom</i>)	123-125	Jared Schneidman
64-65	Dana Burns-Pizer	126	Anonymous; courtesy of Edward Adamson and John Timlin, © 1990
66-67	Guilbert Gates/JSD (<i>top</i>), Carla J. Shatz (<i>bottom</i>)	128	Nancy C. Andreasen, University of Iowa
68	Photograph by Robert Prochnow; painting by Richard Tobias, courtesy of Brooke Alexander, N.Y.	129	Monte S. Buchsbaum, University of California at Irvine
70	Carol Donner (<i>top left and bottom</i>), Semir Zeki (<i>top right</i>)	130	Ian Worpole
71	Guilbert Gates/JSD and Jared Schneidman	132	Carol Donner
72	Guilbert Gates/JSD	134	FPG International
74	Semir Zeki (<i>top</i>), Lawrence Erlsbaum Associates, © 1987 (<i>bottom</i>)	136	Carol Donner (<i>drawing</i>); Johnny Johnson (<i>graph</i>)
75	Courtesy of Semir Zeki (<i>top</i>), D. F. Benson, Archives of Neurology (<i>bottom</i>)	137	Dorothy G. Flood and Paul D. Coleman, University of Rochester (<i>photographs and drawings</i>); Johnny Johnson (<i>graph</i>)
78	Dan Wagner	138	Johnny Johnson
80-81	Ian Worpole; Patricia J. Wynne (<i>inset</i>)	139	Dennis J. Selkoe
82	Patricia J. Wynne	140	Robert P. Friedland, Case Western Reserve University; courtesy of <i>Clinical Neuroimaging</i> , © 1988 John Wiley and Sons, Inc.
83-84	Ian Worpole	142	Johnny Johnson (<i>graphs</i>); Ralph Warren Landrum, University of Kentucky (<i>photograph</i>)
85	<i>Mechanics of the Mind</i> , by Colin Blakemore, © 1977 Cambridge University Press (<i>top</i>); Jared Schneidman (<i>bottom</i>)	144	Tomo Narashima
88	FPG International	146	Laurie Grace
90	Carol Donner	147	Geoffrey E. Hinton
91	Department of Neurology, PET Facility and Image Analysis Facility, University of Iowa	149-151	Laurie Grace
92	Carol Donner	152	<i>Reproduction Prohibited</i> (<i>Portrait of Edward James</i>), by René Magritte, © 1992 Herscovici/Artist Rights Society
		155	Bettmann Archive
		156-157	Jason Goltz
		158	Johnny Johnson
		159	Jason Goltz

SCIENTIFIC AMERICAN®

Established 1845

EDITOR: Jonathan Piel

BOARD OF EDITORS: Alan Hall, *Executive Editor*; Michelle Press, *Managing Editor*; Timothy M. Beardsley; Elizabeth Corcoran; Deborah Erickson; Marguerite Holloway; John Horgan, *Senior Writer*; Philip Morrison, *Book Editor*; Corey S. Powell; John Rennie; Philip E. Ross; Ricki L. Rusting; Russell Ruthen; Gary Stix; Paul Wallich; Philip M. Yam

ART: Joan Starwood, *Art Director*; Edward Bell, *Art Director, Graphics Systems*; Jessie Nathans, *Associate Art Director*; Nisa Geller, *Photography Editor*; Johnny Johnson

COPY: Maria-Christina Keller, *Copy Chief*; Nancy L. Freireich; Daniel C. Schlenoff

PRODUCTION: Richard Sasso, *Vice President Production & Distribution*; *Managers:* Carol Albert, *Prepress*; Tanya DeSilva, *Projects*; Carol Hansen, *Composition*; Madelyn Keyes, *Systems*; Leo J. Petruzzzi, *Manufacturing & Makeup*; William Sherman, *Quality Assurance*; Carl Cherebin; John Mousot

CIRCULATION: Lorraine Leib Terlecki, *Circulation Director*; Cary Zel, *Circulation Manager*; Rosa Davis, *Fulfillment Manager*; Katherine Robold, *Newsstand Manager*

ADVERTISING: Robert F. Gregory, *Advertising Director*. OFFICES: NEW YORK: Michelle Larsen, *Director, New Business Development*; William Buchanan; Peter Fisch; Meryle Lowenthal; Elizabeth Ryan. CHICAGO: 333 N. Michigan Avenue, Chicago, IL 60601; Patrick Bachler, *Advertising Manager*. DETROIT: 3000 Town Center, Suite 1435, Southfield, MI 48075; Edward A. Bartley, *Detroit Manager*; William F. Moore. WEST COAST: 1554 S. Sepulveda Blvd., Suite 212, Los Angeles, CA 90025; Kate Dobson, *Advertising Manager*; Lisa K. Carden, Lianne Bloomer, San Francisco. CANADA: Fenn Company, Inc. DALLAS: Griffith Group

MARKETING SERVICES: Laura Salant, *Marketing Director*; Diane Schube, *Promotion Manager*; Mary Sadlier, *Research Manager*; Ethel D. Little, *Advertising Coordinator*

INTERNATIONAL: EUROPE: GWP International, Düsseldorf; Roy Edwards, GWP International, London; Kurt Bühlmann, GWP International, Zürich. SEOUL: Bisco, Inc. TOKYO: Nikkei International Ltd. SPECIAL PROJECTS: Barth David Schwartz, *Director*

ADMINISTRATION: John J. Moeling, Jr., *Publisher*; Marie M. Beaumonte, *Business Manager*

SCIENTIFIC AMERICAN, INC.

415 Madison Avenue
New York, NY 10017
(212) 754-0550

PRESIDENT AND CHIEF EXECUTIVE OFFICER: John J. Hanley

CORPORATE OFFICERS: *Executive Vice President and Chief Financial Officer*, R. Vincent Barger; *Senior Vice President*, Linda Chaput; *Vice Presidents:* Jonathan Piel, John J. Moeling, Jr.

CHAIRMAN OF THE BOARD: Dr. Pierre Gerckens

CHAIRMAN EMERITUS: Gerard Piel

Mathematica

THE DEFINITIVE SYSTEM FOR TECHNICAL COMPUTATION

"Not merely a product but a revolution"

— Macworld


"The importance of the program cannot be overlooked"

— New York Times


Basic function: Integrated environment for numerical, symbolic, graphical computation, interactive programming.

Users: Scientists, engineers, mathematicians, programmers, financial analysts, students. Over 150,000 worldwide. Includes all 50 largest U.S. universities.

Numerical computation: Arbitrary-precision arithmetic, complex numbers, special functions (hypergeometric, elliptic, etc.), combinatorial and integer functions. Matrix operations, root finding, function fitting, Fourier transforms, numerical integration, numerical solution of differential equations, function minimization, linear programming.



Symbolic computation: Equation solving, symbolic integration, differentiation, powerseries, limits. Algebraic operations, polynomial expansion, factorization, simplification. Operations on matrices, tensors, lists, strings.




Graphics and sound: 2D, 3D plots of functions, data, geometrical objects. Contour, density plots. 3D rendering with intersecting surfaces, lighting models, symbolic descriptions. Color PostScript output, combining and labeling, publication quality graphics, animation (most versions). Sound from waveforms and data (most versions).

Programming: High-level, interactive, symbolic language. Procedural and functional programming



constructs. Transformation rules and pattern matching. Fully compatible on all platforms. No built-in limitations on computation size.



External interface: Input of data (numbers, records, text) from files, programs. Output in TeX, C, Fortran, PostScript. Calling of external programs and functions. General *MathLink* interprocess communication mechanism.

User interface: Electronic book interactive documents mixing text, graphics, animations, calculations. Graphics, animation, sound interapplication compatibility. Style sheets, hierarchical outlining. Computation kernel can run on remote computer (most versions).

Additional material: Journal, newsletters, more than 30 books. Add-on packages, free *MathSource* electronic resource.

Versions: Macintosh • Microsoft Windows • MS-DOS • NEC PC • DEC RISC, VAX • HP • IBM RISC • MIPS • NeXT • SGI • Sony • SPARC, Sun-3 • CONVEX and others • Network licensing available. Student versions. Now shipping Version 2.1.

For the latest information call Wolfram Research at:

1-800-441-MATH



Wolfram Research

Wolfram Research, Inc.

100 Trade Center Drive, Champaign, IL 61820-7237, USA
217-398-0700; fax: 217-398-0747; email: info@wri.com

Wolfram Research (UK) Ltd.

Evenlode Court, Main Road, Long Hanborough, Oxon OX8 2LA, UK
+44-(0)993-883400; fax: +44-(0)993-883800; email: info-uk@wri.com

For the growth of your business, the prosperity of your business, there is no more powerful tool than the telecommunications network.

And at GTE, there is nothing vague about the

power we are putting into your hands when we design that network for you.

For one of the nation's leading retailers, we are now implementing one of the largest satellite-based private

THE POWER OF A NETWORK DESIGNED
FOR YOUR BUSINESS ALONE.

data and video communications networks in the world.

A network built around the advanced technology of GTE
Spacenet. A network like no other because it was created
to power the success strategies of that company alone.

The power of a personalized telecommunications
network. It is an unmistakable advantage.

We've given it to others. And we can give it to you.

Because at GTE, the power is on.



GTE

THE POWER IS ON



LETTERS TO THE EDITORS

All about Eve

There is distressing apparent arrogance on both sides of the debate on modern human origins ["The Recent African Genesis of Humans," by Allan C. Wilson and Rebecca L. Cann; and "The Multiregional Evolution of Humans," by Alan G. Thorne and Milford H. Wolpoff, *SCIENTIFIC AMERICAN*, April], although it has been more prominent in the pronouncements of the students of mitochondrial DNA than in those of the paleoanthropologists. Recently, using the very same mitochondrial DNA material available to Wilson and Cann, Alan R. Templeton of Washington University has shown that there are 100 possible genealogies that are more parsimonious than the one the authors presented. S. Blair Hedges of Pennsylvania State University and others further demonstrated that 50,000 possible trees are statistically "better" than the one advocated by Wilson and Cann. The overwhelming conclusion is that the available genetic evidence not only cannot provide a time scale for the emergence of modern human form but it also does not provide any hint that Africa was the locale where this took place.

C. LORING BRACE
Museum of Anthropology
University of Michigan

Both papers in your April issue Eve debate seem logically flawed. The Thorne and Wolpoff scenario involves too much special pleading to be taken seriously. They would have characters marking specific population lineages surviving over hundreds of thousands of years, which implies that these features must confer great selective advantages. At the same time, there is enough gene flow involving advantageous new mutations to keep all the populations evolving toward modern *Homo sapiens* in lockstep. So selection and gene flow are making human populations more like one another, while selection and lack of gene flow are keeping them differentiated, and this strange balance is maintained for almost a million years. I think not.

Finally, the opening paragraph of the Thorne and Wolpoff article is blatantly self-serving revisionist history. Concerning the hominid status of *Ramapithecus*, Wolpoff writes, "Molecular biologists entered that discussion and sup-

ported the minority position held by one of us...." No one, to my knowledge, credits Wolpoff with any meaningful contributions to the issue until after the original presentations that Wilson and I made in 1966 and 1967.

VINCENT SARICH
Department of Anthropology
University of California at Berkeley

Cann replies:

Advances in the analysis of large data sets are necessary to help evaluate just how probable certain branch points in the human mitochondrial DNA trees really are. Every computer program makes assumptions about the biology of mitochondria, and not all assumptions are valid for every region of the molecule.

The molecular trees are only one line of evidence in a hypothesis of African origins. Two additional findings are important. First, the highest levels of mitochondrial and nuclear genetic diversity for any living human group are found in populations from sub-Saharan Africa. Most evolutionary biologists would accept that finding as strong support for Africa as our center of origin.

Second, the oldest human fossils judged to be anatomically modern in appearance come from Africa. Important blade technologies associated with modern humans are also found first in Africa, not Asia. As I have stated before, this field can move no further without the evaluation of DNA extracted from a multitude of human fossils less than 200,000 years old.

Thorne and Wolpoff reply:

We do not envision regional continuity markers as necessarily conferring selective advantage. Rather the great selective advantage belongs to the features that change across the world over the course of human evolution. We have sought continuity markers that do not respond to unique local selection because only these can address the issue of ancestry rather than adaptation as a potential explanation for their persistence. Balances between local selection (or genetic drift) and gene flow for features create spatial gradients that are sensitive to the magnitudes of each, but neither can predominate over the other.

Sarich misunderstands the main point

of our *Ramapithecus* reference: to refute the claim by Wilson and Cann that a "molecular clock" ultimately resolved the dispute. The arguments about *Ramapithecus* began with the first publication on the genus in 1934. The debate ended because of new fossil remains, better geologic information and anatomic studies, not a "clock."

Drinking and Genes

In "D₂ or not D₂," by John Horgan [*"Science and the Citizen,"* *SCIENTIFIC AMERICAN*, April], the statement that we found a gene linked to alcoholism is incorrect. Rather we claimed an association, not a linkage, of the D₂ gene with alcoholism. The statement attributed to one of us that job applicants will be tested for the A1 allele is also untrue. We are strongly opposed to such use of a diagnostic test for alcoholism.

ERNEST P. NOBLE
Department of Psychiatry
and Biobehavioral Sciences
University of California at Los Angeles

KENNETH BLUM
Department of Pharmacology
University of Texas
Health Science Center
San Antonio

Credit for Catalysts

In my article "Solid Acid Catalysts" [*SCIENTIFIC AMERICAN*, April], I wrongly attributed the synthesis of the zeolite Rho to Richard Barrer of Imperial College, London. Although Barrer was the first, in 1943, to synthesize a zeolite (KFI) that does not occur naturally, the credit for Rho belongs to Harry Robson, who produced it in 1965 at the Exxon Research and Development Center in Baton Rouge, La. The structure of Rho is similar to that of KFI, hence the source of, but not the justification for, my lapse in memory. I apologize to Robson and to David Shoemaker, Robert Ogilvie and Phillip Manor of the Massachusetts Institute of Technology, who solved the structure of Rho in 1973.

SIR JOHN MEURIG THOMAS
Royal Institution of Great Britain
London

Life is unpredictable. Thankfully, the Accord is not.

Will the stock market make a meteoric rise? Will bell-bottom pants make a comeback? Does the guy driving next to you know what he's doing?

Maybe.

At Honda, we concentrate on the third question. Which is why the Accord is a safe car to drive.

Settle in behind the wheel of the Accord and the first safety feature will be right in front of you. The driver's side airbag Supplemental Restraint System (SRS). It's standard on all 1992 Accords.

Take to the highway and there's a direct line of communication between you and the road, thanks to the 4-wheel double wishbone suspension. The result is extraordinary handling and control.

And, if life should toss you an unexpected turn in your Accord EX, underfoot are anti-lock brakes: one of the safest, most effective ways to stop a car. In panic situations, even under adverse conditions, they help prevent the wheels from locking up.

We could go on. The Accord's safety features are numerous. Rest assured, all are designed to give you peace of mind as you travel through life.

Which means that whatever tomorrow may bring, you're prepared in an Accord. A soothing fact that we know you'll be thankful for.  **HONDA**



Save big on some mind-expanding software. And help stop a mind-destroying disease.

About this time last year, we carried out an interesting experiment.

Knowing a high percentage of the readers of this magazine to be PC users, we decided to advertise our award-winning graphics software in these pages. To sweeten the deal, we offered it at an unheard-of price. And promised a part of the proceeds to the World Wildlife Fund.

Well, it was a roaring success. We welcomed lots of new people to the exciting world of PC graphics. They got great software at a great price. And the World Wildlife Fund received a healthy check.

This year, the deal's even better. Once again, we're offering our best software at a great price. But in honor of this special issue on the brain, we're upping our charity donation. Sending a full 10% of the proceeds from your purchase to the Alzheimer's Association, to help stop one

of the brain's most formidable enemies.

This opportunity may not come around again. So please, if you use a PC (or have access to one), read on.

Put a little art in your work with Windows Draw:™ America's #1-selling graphics package.

When Windows Draw was introduced last year, it started a popular revolution. Combining unprecedented ease of use and affordability, Windows Draw was the first graphics package designed to join word processing as a "must-have accessory" (*PC Magazine*).

Just about *anyone* can benefit from Windows Draw. It lets you easily add visual impact to newsletters, presentations, memos, reports, charts, invitations – you name it.

In fact, because it's compatible with your favorite PC programs, you can use Windows Draw to dress up just about *everything* you do.

Easy-to-use drawing and color tools are right on the screen. Pull-down menus list an enticing selection of options available to you – including special effects. And help information is available at the touch of a key.

If you don't want to do your own drawings, you can choose from a dazzling library of 2,600 ready-made, professionally drawn images.

Including everything from jet planes and golfers to fancy borders and technical symbols. And you can edit them just like your own drawings.


You can even make your text look as good as your graphics.


You'll be able to curve words, wrap them around pictures, shade them, color them and more. And you'll enjoy using Windows Draw in combination with PageMaker or your favorite word processing package.

COMPUTE Magazine called Windows Draw "a beautifully

Use Windows Draw to add illustrations, borders, fancy text and more to everything you produce.

Windows Draw and Designer are terrific tools for desktop publishing and presentations.





For high-precision illustration, Micrografx Designer simply has no equal.

designed draw program." "An outstanding bargain," said *PC Sources*. And *PC World* declared Windows Draw "an ideal package for the non-professional artist."

For precision illustration, Micrografx Designer™ is the choice of experts and non-experts alike.

For everyday graphics projects, you just can't beat Windows Draw. But for precision illustration, you need the celebrated power of Micrografx Designer.

We like to call Designer "a drawing board for the imagination." It lets you easily turn the most intricate visual concept into a brilliant, incredibly precise, professional-quality illustration.

Designer is the choice of



Someone you know has Alzheimer's. A vicious destroyer of memory, personality, and ultimately, the brain itself.

The Alzheimer's Association has been working desperately to find a treatment for this frightening disease. And today, there are glimmers of hope. But much more must be done. That's why, in honor of this special issue on the brain, we'll donate 10% of the proceeds from your purchase to the Alzheimer's Association.

engineers, scientists, educators and other demanding professionals. But it's also a favorite of non-technical people who simply want to know they're working with the best PC illustration tools available.

Designer features accuracy to 1/3,000 of an inch. Up to 64 layers in a single illustration. A huge 132" by 132" drawing area, and many more high-end features that have put it in a class by itself.

Designer is also a terrific *presentation* package. With over \$5,000 worth of type and text-handling software built-in, it lets you easily turn your text and illustrations into powerful hard-copy, slide or PC presentations.

In awarding Designer its highest rating, the universally respected *Software Digest* credited it with "state-of-the-art

features, good performance and unmatched ease of learning and use in a sophisticated package."

Windows Magazine said: "Designer's flexibility and precision make it the best choice for any serious user." And Designer took home *InfoWorld's* prestigious "Product of the Year" award for 1991.

A no-risk, no-question-about-it terrific offer.


Here, in a nutshell, is our offer:

A terrific price. We'll send you *Windows Draw*, for everyday graphics projects, for only **\$89** (reg. \$149.95). Or *Designer*, for precision illustration, for only **\$299** (reg. \$695). Shipping and handling are \$10 for each product ordered (U.S. only).

A terrific cause. As mentioned, we'll donate 10% of the proceeds from your purchase to the Alzheimer's Association to help research efforts.

A terrific help. We'll also include unlimited 24-hour technical support by telephone (weekend hours too!) for any product you purchase via this offer.

To take advantage of this offer, call us directly at the number below and mention this ad. Please, since this is a limited-time offer, *don't wait to call*. Do something good for yourself *and* for others less fortunate. *Call today!*




1-800-653-1980

We accept VISA, MasterCard and American Express. Offer expires October 31, 1992. Limit five per person.

M I C R O G R A F X ®

Micrografx, Inc., 1303 Arapaho, Richardson, TX 75081 (214) 234-1769. Micrografx has offices in Toronto, Paris, London, Munich, Milan, Sydney, Copenhagen and Tokyo. Copyright ©1992, Micrografx, Inc. All rights reserved. Micrografx is a registered trademark and Micrografx Designer and Windows Draw are trademarks of Micrografx, Inc. All other products are trademarks or registered trademarks of their respective owners.

© 1992 SCIENTIFIC AMERICAN, INC



SEPTEMBER 1942

"A few years ago, the statement that the Orion nebula was partly composed of iron would have appeared to be incredible. Now, 10 forbidden lines of *Fe* II, and about 15 of *Fe* III, have been definitely observed by Wyse, so that there can be no doubt about it. In the meantime, all surprise has been removed by the even more remarkable discovery that atoms of iron form part of the gas which is distributed throughout interstellar space. Year by year, the evidence for the general uniformity of composition of matter, throughout the regions of space accessible to our observation, becomes more and more impressive."

"Although scanning microscopes of low resolution were successfully constructed even with the image observed directly on the fluorescent screen of a kinescope, high-resolution instruments required a slower method of recording. A facsimile receiver registering an image in eight minutes proved to be a satisfactory solution."

"Many persons bleed excessively from minor wounds because their blood contains too little of the clotting principle (prothrombin). No satisfactory method has been known before to control bleeding in persons thus afflicted and even

pulling their teeth has involved serious hazard to their lives. The new clotting globulin supplies the constituent in which the blood of 'bleeders' is deficient and places in the hands of the medical and dental professions an effective means for stopping the flow of blood from minor wounds in a few seconds."

"A curious sensation sometimes experienced by pilots, and termed the 'leans,' has recently been described. Here's what leads up to it: During instrument flight, a slow lateral tilting of the aircraft remains unnoticed. When finally the tilting has become pronounced enough for the pilot to be aware of it, he corrects the attitude of the plane in a much shorter period of time than that which was required for the original tipping. Now the human ear is so constructed that it readily notes rapid acceleration but not slow acceleration. The pilot thus ends up with the sensation that the plane is now tipping down on the opposite side."




SEPTEMBER 1892

"In order to test the power of gripping in the young infant, Dr. Louis Rob-

inson placed his fingers against the palm. The contact at once caused the hand to close apparently by pure reflex action, since it made little difference whether the child was asleep or awake. He then slowly, but with a slight jerking motion, lifted his fingers, and found to his surprise that the child tightened its grasp and allowed him to raise it from the bed. In many cases a newly-born child would hang and support its weight with ease for a minute. Still more surprising was the fact that in most instances it would make no objection to the experiment whatever. Among the newly-born offspring of the human race these faculties were of no use. It seemed, therefore, legitimate to infer that the astonishing prehensile power in the hands of the modern infant was a vestige of the habits which for many epochs saved their arboreal forefathers during their tender youth from destruction."

"The recent lowering of the equine trotting record by Nancy Hanks to 2 minutes 5 $\frac{1}{4}$ seconds is a remarkable verification of a mathematical law that has been followed to a fraction of a second for the past sixty-two years. This law is that the time to trot a mile is reducing at the rate of eleven twenty-sixths of a second a year. This guide shows that in the year 2047 the mile a minute gait will be reached, and that 297 years hence it will be in order to race trotters with the lightning's flash."

"Abbot Rousselot, professor at the Carmelite School, has succeeded in creating the series of apparatus necessary for registering, one by one, the motions whose *ensemble* constitutes a word or a phrase. Evidently, every time that, for any cause whatever, the air contained in the rubber tube enters into vibration, the vibrations will be communicated to the air of the drum, and after this the rubber and then the plate and lever will enter into motion. If the cylinder is revolving at the same time, the line that will be inscribed thereon by the point of the lever, instead of being straight, will become a tracing of the vibrations. It will be possible hereafter to note the pronunciation of any language, dialect or idiom whatever, without relying upon the testimony of the ear, which distinguishes but slight differences between the modes of speaking of several individuals."



Abbot Rousselot's phonetic recorder

W H A T M A K E S A M O M E N T




A M E M O R Y



**Special Offer for
Scientific American readers...**

SAVE 20% ON HARDCOVER BOOKS

**New From the Book Division of
Scientific American**



LIFE

The Science of Biology Third Edition

William K. Purves, Gordon H. Orians, and H. Craig Heller bring you this up-to-date and accessible introduction to biology. Includes new chapters on conservation biology and recombinant DNA. Cloth (ISBN 2276): \$59.95. **Your price: \$47.96**

RECOMBINANT DNA Second Edition

THE authoritative introduction to recombinant DNA has been revised and updated. James D. Watson and his team report on the latest developments in polymerase chain reactions, gene transfer, oncogenes and anti-oncogenes, AIDS, and more. Paper (ISBN 2282): **\$32.95** Cloth (ISBN 1994): \$49.95. **Your price: \$39.96**

IMMUNOLOGY

This remarkable new textbook presents immunology in an experimental context. Professor Janis Kuby focuses on the cellular and molecular levels of immune response, including T cell receptors, immunodeficiency diseases, and AIDS. Cloth (ISBN 2257): \$46.95. **Your price: \$37.56**

DNA REPLICATION Second Edition

Arthur Kornberg and Tania A. Baker Cloth (ISBN 2003): \$64.95. **Your price: \$51.99**

MOLECULAR CELL BIOLOGY Second Edition

James Darnell, Harvey Lodish, and David Baltimore Cloth (ISBN 1981): \$56.95. **Your price: \$45.56**

FOUNDATIONS OF COMPUTER SCIENCE

Alfred V. Aho and Jeffrey D. Ullman integrate models, concepts, logic and techniques in computer science with principles of discrete mathematics. The result is a new approach to computer science theory that clearly explains how mathematical abstractions become working programs. Cloth (ISBN 8233): \$56.95. **Your price: \$45.56**

THE SCIENCE OF MUSICAL SOUND

John R. Pierce Paper (ISBN 6005): **\$19.95**

BIOCHEMISTRY Third Edition

Lubert Stryer Cloth (ISBN 1843): \$56.95. **Your price: \$45.56**


THE ELEMENTS OF PHYSICAL CHEMISTRY

Peter Atkins, the author of the most widely adopted physical chemistry textbook, brings you a brief, but sound overview of physical chemistry principles and phenomena. Cloth (ISBN 2364): \$42.95. **Your price: \$34.36**

SENSORY ECOLOGY How Organisms Acquire and Respond to Information

How do organisms obtain and use information from their environments? Covering the full spectrum of life, David B. Dusenbery brings together the latest research in physics, microbiology, botany, animal behavior, and psychology to explore this fascinating topic. Cloth (ISBN 2333): \$34.95. **Your price: \$27.96**

Now in Paperback



VISIONS OF SYMMETRY

Notebooks, Periodic Drawings, and Related Works of M.C. Escher Doris Schattschneider Paper (ISBN 2352): \$24.95 Cloth (ISBN 2126): \$44.95. **Your price: \$35.96**

SHADOWS OF CREATION


Dark Matter and the Structure of the Universe Michael Riordan and David Schramm Paper (ISBN 2366): \$11.95 Cloth (ISBN 2157): \$19.95. **Your price: \$15.96**


OUR UNIVERSE

An Armchair Guide Michael Rowan Robinson Paper (ISBN 2359): \$19.95 Cloth (ISBN 2156): \$29.95. **Your price: \$23.96**

FRACTALS, CHAOS, POWER LAWS

Minutes From an Infinite Paradise Manfred Schroeder Paper (ISBN 2357): \$19.95 Cloth (ISBN 2136): \$32.95. **Your price: \$26.36**





SPACETIME PHYSICS

Introduction to Special Relativity Second Edition

In this thoroughly updated edition of an undisputed classic, Edwin Taylor and John Archibald Wheeler present a highly accessible overview of relativity, covering microgravity, collider accelerators, satellite probes, neutron detectors, radioastronomy, and pulsars. Paper (ISBN 2327): **\$19.95** Cloth (ISBN 2326): **\$29.95** **Your price: \$23.96**

ANOTHER FINE MATH YOU'VE GOT ME INTO...

Scientific American columnist Ian Stewart brings you 16 brain-teasing puzzles, games, and fables that challenge, entertain, and inspire while they demonstrate compelling mathematical ideas. Paper (ISBN 2341): **\$13.95** Cloth (ISBN 2342): **\$21.95** **Your price: \$17.56**

THINKING, PROBLEM SOLVING, COGNITION Second Edition

Why do some people come up with creative solutions, while others get mentally paralyzed? Psychologist Richard Mayer uses dozens of probing exercises to explore all aspects of higher cognitive processes. Paper (ISBN 2215): **\$22.95**

HOW TO THINK ABOUT STATISTICS Revised Edition

Using compelling examples, Professor John L. Phillips, Jr. explores how to understand, interpret, and apply statistics. Covers everything from polls, surveys, and economic indicators, to advertising claims and research findings. Paper (ISBN 2287): **\$14.95**. Cloth (ISBN 2288): **\$24.95** **Your price: \$19.96**

NEW BOOK OF PUZZLES

101 Classic and Modern Puzzles to Make and Solve

Jerry Slocum and Jack Botermans, the creators of *Puzzles Old and New*, are back with an all-new treasure chest of puzzles. You get over 100 of the most tantalizing, ingenious, beautiful puzzles ever designed, all reproduced in magnificent color with complete, easy-to-follow instructions. Cloth (ISBN 2356): **\$19.95** **Your price: \$15.96**

ARCHAEOLOGY

The Science of Once and Future Things

How did complex societies develop? What caused them to change or collapse? What can past societies tell us about our society and its future? Bryan Hayden takes a compelling look at archaeology, exploring the basic concepts and tools to the complex archaeological puzzles. Paper (ISBN 2307): **\$19.95**

CODES, PUZZLES, AND CONSPIRACY

A New Mathematical Thriller From Dr. Ecco

Dennis Shasha
Paper (ISBN 2314): **\$11.95**
Cloth (ISBN 2275): **\$17.95**
Your price: \$14.36

THE EPICUREAN LABORATORY

Exploring the Science of Cooking

Tina L. Seelig
Paper (ISBN 2162): **\$15.95**

CAN YOU WIN?

The Real Odds for Casino Gambling, Sports Betting, and Lotteries

Michael Orkin
Paper (ISBN 2155): **\$11.95**

HELPLESSNESS

On Development Depression & Death

Martin Seligman
Paper (ISBN 2328): **\$14.95**

✂ cut here and mail in an envelope today

Save on Videos Too!

FRACTALS

An Animated Discussion
with Edward Lorenz and
Benoit B. Mandelbrot

VHS tape (ISBN 2213):
\$59.95. Your price: \$47.99

POWERS OF TEN

The Films of Charles
and Ray Eames
Volume One

VHS tape (ISBN 5029):
\$39.95. Your price: \$35.00

Mail This Today and Save 20% on all Hardcover Books

W.H. FREEMAN AND COMPANY
4419 West 1980 South
Salt Lake City, UT 84104

Name _____

Address _____

City/State/Zip _____

☐ Check enclosed

☐ VISA

☐ Master Card

Account # _____

Exp. Date _____

Signature _____

Qty.	Title	ISBN	Price

Sub-Total	\$ _____
Sales Tax (NY, CA, UT residents only)	\$ _____
Shipping & Handling (\$1.95 for 1st book, \$1.25 for each add'l book)	\$ _____
Total	\$ _____

Payment in U.S. dollars must accompany all orders. Prices valid in North America only and are subject to change without notice. Allow four weeks for delivery. For Canadian orders, add \$2.25 shipping and handling for the first book and \$1.50 for each additional book, and please enclose 7% GST. N.Y. residents, please compute sales tax based on book total and shipping.



Gravity Quantized?

A radical theory of gravity weaves space from tiny loops

Like cats and dogs, round holes and square pegs, and polka-dotted ties and striped shirts, general relativity and quantum mechanics just do not go well together. Einstein's theory of general relativity depicts gravity as warps in the fabric of space and time, whereas quantum mechanics describes the other forces ruling matter—namely, electromagnetism and the strong and weak nuclear forces—as interactions of energy packets called quanta.

Each theory works well in its sphere: quantum mechanics primarily addresses microscopic effects and general relativity large-scale ones. But physicists have long hoped to fuse the two world-views into a unified theory of all physical phenomena, from quarks to quasars. For the past decade, by far the dominant candidate for a so-called quantum gravity theory has involved superstrings, infinitesimal loops of energy whose wriggles supposedly give rise to all the particles and forces of nature.

In the past few years, however, string theorists seem to have lost their bearings in the theory's labyrinthine mathematics. Meanwhile another, more modest approach to quantum gravity has begun to attract attention. The theory bears at least a superficial resemblance to string theory, in that its fundamental features are also tiny loops. Yet these loops correspond not to objects or events but to space itself.

The model has been called the loop-space theory or the Ashtekar theory, after Abhay Ashtekar of Syracuse University, who took the first step toward the concept by tinkering with the equations of general relativity. (Ashtekar is not fond of either name, since the loop-space concept is just part of the theory and he is only one of several

contributors. He is open to suggestions for a new name.) Since the theory was conceived seven years ago, more than 200 papers by dozens of theorists have been devoted to it. "It's the greatest thing to happen in years," says John A. Wheeler of Princeton University, a pioneer in quantum gravity research. "They still have a lot to understand, but this is a way to gain new insights."

Formulating gravity in quantum mechanical terms once seemed to be a fairly straightforward task, Ashtekar notes. General relativity itself predicts that gravity, like the other forces of nature, propagates across space at the speed of light. In fact, astronomers are already searching for intense gravitational waves generated by pairs of black holes and other exotic phenomena [see "Catching the Wave," by Russell Ruthen; *SCIENTIFIC AMERICAN*, March]. To "quantize" gravity, physicists needed simply to show that gravitational waves actually consist of quantum particles, or gravitons, with properties

similar to those of photons, gluons and other force-carrying particles.

Beginning in the 1960s, theorists seemed to have developed the mathematical tools to accomplish this feat. The tools, called perturbation methods, helped to eliminate inconsistencies—such as predictions whose likelihood exceeded 100 percent—that arose when physicists tried to derive precise predictions from quantum theories of electromagnetism and the nuclear forces. Perturbation methods allow workers to avoid the inconsistencies by deriving a series of approximate rather than exact solutions to the theories.

But perturbation failed to eliminate the inconsistencies arising from quantized models of gravity. The reason for this failure, Ashtekar explains, has to do with a fundamental difference between gravity and the other forces of nature. Perturbation methods assume that the force-transmitting quanta career, collide and pirouette on a fixed, unchanging stage of space and time.

But according to general relativity, gravitons alter the geometry of space and time, and that alteration in turn affects the behavior of the gravitons. This self-referential, nonlinear property of gravity distinguishes it from all other forces and makes it impossible to tame with perturbation methods.

In the mid-1980s Ashtekar began searching for a "non-perturbative" quantum gravity theory, one that could generate consistent solutions without approximations. He decided that the key was to replace the metric, a variable representing the geometry of space and time in general relativity, with some more malleable variable. He eventually chose a mathematical function originally developed by Amitabha Sen, a physicist now at Motorola.

The Sen variable splits time and space, which are unified into a single, four-dimensional continuum under Einstein's metric, into two distinct entities subject to quantum uncertainty and analogous to position and momentum in




TOO LOOPY? Lee Smolin of Syracuse University illustrates the loop-space concept at conferences with a model made of interlinked key rings. Photo: Frank Veronsky.



THE ONLY BRAKE PEDAL THAT MAKES A SPORTS CAR GO FASTER.

The brake pedal in the new Mazda RX-7 may be the lightest in the world. We made it out of aluminum, then drilled holes in it. Why go to such extremes? We know what a handful of enthusiasts have always known. In a pure sports car every ounce counts. That's why we started with a sequential twin turbo rotary engine that's 200 pounds lighter than a comparable piston engine. Which gave us a head start on a weight reduction program nothing short of obsessive (we even trimmed the spark plug leads). The result is something that you may never see again. 2789 lbs. 255 horsepower. 0-60 in 4.9 seconds. And exclusive rights to the label pure sports car.



THE 1993 MAZDA RX-7

Sequential twin turbo rotary engine. Independent double-wishbone suspension. 11.6-inch, ventilated four-wheel disc brakes with ABS. Driver's-side air bag. 36-month/50,000-mile, no-deductible, "bumper-to-bumper" warranty. 24-hour roadside assistance. See your Mazda Dealer for limited-warranty details. To receive a free RX-7 brochure, call 1-800-639-1000.

© 1992 Mazda Motor of America, Inc.

mazda
IT JUST FEELS RIGHT.®

© 1992 SCIENTIFIC AMERICAN, INC

conventional quantum mechanics. Altered in this way, Einstein's equations suddenly resembled those used in quantum theories of electromagnetism. The equations also became much simpler and easier to manipulate.

Shortly after Ashtekar reported on his work in 1985, Theodore A. Jacobson of the University of Maryland, Carlo Rovelli of the University of Pittsburgh and Lee Smolin of Syracuse University determined that Ashtekar's new formulation could indeed generate exact solutions for certain quantum gravitational states. Borrowing from the mathematics of knot theory, Rovelli and Smolin then showed that each of these quantum states could be considered as a different knot, of which the simplest form was a loop. These loops are akin to the lines of force generated by an electric field.

Ashtekar, Smolin and Rovelli have been trying to demonstrate how these knots and loops can be woven together, like the links in a chain-mail coat, to form the apparently seamless fabric of space. The loops are so tightly knitted

that they cannot be distinguished except at a distance of 10^{-35} meter, a length as small in comparison to an atom as an atom is in comparison to the solar system. General relativity predicts that at precisely this scale, known as the Planck scale, quantum gravitational effects should become significant. This is also the scale at which superstrings are thought to operate.

One crucial difference between loop-space theory and superstring theory is that the latter purports to explain not just gravity but all the forces and particles of nature. Such a theory is sometimes called a theory of everything. The loop-space theory is currently less ambitious, generally describing only the passage of gravitons through otherwise empty space. The gravitons appear as "embroidery" knitted into the loops.

Ashtekar and his colleagues have just begun inserting particles corresponding to other forces and to mass, such as photons or protons, into the loop-space model. Although the model correctly describes the interactions of gravity and matter, Smolin concedes that it may

never "explain" all the forces of nature from first causes in the way superstring theory is intended to do. "We don't have some overreaching ambition to have a complete theory of nature," Smolin says. "Our idea is to have a quantum mechanical theory of gravity."

That in itself would be a tremendous achievement. Gravity is by far the weakest of all the forces, and under most circumstances its effects on small-scale, quantum interactions can be ignored. But a quantum gravity theory is crucial for understanding what happens in the first instant of the big bang or toward the center of a black hole, where energies and densities surge toward infinity and Planck-scale effects predominate.

Some physicists have suggested that the loop-space theory is based on flawed assumptions and will not lead to any significant breakthrough. "It's my opinion, and the opinion of some other superstring theorists, that this is just wrong," says David Gross of Princeton University, a prominent superstrings advocate. Eventually, he predicts, the loop-space theorists will founder on the same mathematical problems that have blocked similar efforts.

This assessment is too pessimistic, says Gary T. Horowitz of the University of California at Santa Barbara. He suggests that some string theorists, because they were trained as particle physicists, are overly skeptical of loop-space theory and any other approach to quantum gravity rooted in general relativity. While the loop-space workers "have not proved their case," says Horowitz, who describes himself as a string theorist, "I don't think their problems are insurmountable." After all, he adds, the jury is still out on superstrings, too.

Smolin considers superstrings and loops to be complementary rather than competing approaches to a more fundamental theory of physics. "We should be able to learn a lot from each other," he says. "Neither approach is likely to be the entire truth. People will learn more and more about how to solve different parts of the problem, and then someone will come along with an entirely new theory that will incorporate both of these views."

A consistent theory of quantum gravity, Smolin adds, may also help resolve some of the nagging philosophical questions that quantum mechanics injected into physics early in this century. Is uncertainty intrinsic to reality? Does reality exist independent of observation? "I don't think you can solve these problems with philosophy or just by talking about them," Smolin says. "I think it will require some new physics."

—John Horgan

It Came from Within

Thomas Gold is a true scientific iconoclast. He is perhaps best known for his claim that oil and natural gas stem not from decayed organisms, as virtually all other experts maintain, but from nonbiological processes occurring deep inside the earth. Now Gold, a physicist at Cornell University, is once again flouting conventional wisdom. In the *Proceedings of the National Academy of Sciences*, he argues that life first began billions of years ago not at or near the earth's surface, as most biologists believe, but in rocky fissures far underground.

Gold speculates that some of the primitive microorganisms migrated to the surface, where they evolved into amoebas, anchovies and actors. But most still dwell underground, sustained not by light and oxygen as most organisms are but by sulfur compounds, methane or other energy-rich chemicals. Moreover, similar subterranean communities may exist within other planets (and moons) in the solar system.

Gold points out that the most primitive of all living organisms are anaerobic, thermophilic microbes called archaeobacteria, which thrive in hot (up to 120 degrees Celsius), airless environments. These organisms have been found in boreholes more than 500 meters deep, in hot springs on the seafloor and in other unlikely locations. Not surprisingly, the most dramatic borehole evidence that Gold cites was collected at a drilling project in Sweden that he himself initiated six years ago to find nonbiological oil and gas. Although that effort has been a bust—according to most experts, if not to Gold—he says the drills have turned up microbes five kilometers down.

Gold argues, moreover, that the earth's interior would have provided a much more hospitable environment for proto-life four billion years ago than the surface would have, ravaged as it was by asteroids and cosmic radiation. And if life emerged within the earth, then why not within other planets? "Deep, chemically supplied life," Gold says, "may be very common in the universe."

Carl R. Woese of the University of Illinois, an authority on archaeobacteria, applauds Gold's speculation—or its terrestrial component, at least. "All the theories in the microbiology world and the origin of life are in flux," he says, "so it's time to widen the realm of the possible." Gold's suggestion that life might also dwell within other planets, however, "goes beyond what I'm comfortable with," Woese says.

—John Horgan

Now schools without French teachers can still have lessons in French.



Students at three high schools in northwestern suburbs of Chicago now can take classes their schools might not have been able to afford otherwise — thanks to a system provided by Ameritech.

Called Distance Learning, the system is a new application for video conferencing, a service originally intended for business use. By video, a teacher can combine students from three schools to form a class and teach them simultaneously even though the schools are twenty miles apart. It's only one of many educational efforts Ameritech has under way. Our Buddy System, for example, is linking students and home computers with their schools and a whole world of information. We're also providing grants to students doing graduate work in telecommunications

and supporting teachers writing curricula using our Ameritech PagesPlus® directories as learning tools.

At Ameritech, we're developing innovative products and services of all kinds to help make sure today's students aren't just ready to face the future, but to shape it.

For more information on our contributions to the quality of life in communities we serve, call 1-800/242-8580.

THE COMMUNICATIONS COMPANIES OF AMERITECH ARE:

Illinois Bell • Indiana Bell • Michigan Bell • Ohio Bell • Wisconsin Bell
Ameritech Audiotex Services • Ameritech Credit • Ameritech Development
Ameritech Information Systems • Ameritech International
Ameritech Mobile Communications • Ameritech Publishing
Ameritech Services • The Tigon Corporation.

AMERITECH

Connecting You With Tomorrow

4,000,007 reasons to upgrade to System 7.

1. Rave reviews.

MacUser gave System 7 its Editors' Choice Award. Byte, its Award of Excellence. Software Publishers Association named it a Best New Business Software Product of the Year.

2. One-step installation.

The "Before You Install" disk confirms the compatibility of your software (including INITs). Then, simply click the mouse to install System 7. Any questions? 90 days' toll-free phone assistance included.

3. It's developed quite a following.


Developers have shipped hundreds of applications that take advantage of System 7 power. Microsoft, Claris, Lotus, Aldus, Adobe and many more.

4. Runs on your Macintosh today.

System 7 works on all Macintosh computers. To run multiple programs, four megabytes of memory are recommended. Two megabytes are required.

System Software for the Apple® Macintosh®

System 7 brings a new world of power, ease of use, and versatility to every Macintosh personal computer.



5. Powerful software utilities.

System 7 will let you run the latest software utilities from Apple, including Macintosh PC Exchange, QuickTime™ and AppleTalk® Remote Access.

6. More power out of the box.

Macworld says, "The new Finder...is worth the cost alone." System 7 improves the way you organize, view and work with your information. Built-in file sharing lets you share files over a network, simply and easily.

7. Keep your investment current.

The best way to keep up with the latest technology is to stay ahead with System 7. You'll need it to take full advantage of today's applications, hardware and software. And it will help you grow with the future as new technology innovations come along.

Macintosh System 7. If the first seven reasons didn't convince you to upgrade, just ask the four million Macintosh® users who have already discovered the power of System 7.™ Call us at **1-800-538-9696, ext. 185.** We'll give you the name of an Apple software reseller near you. And you'll discover the power of System 7. The power to be your best.*



Noisy Nucleotides

DNA sequences show fractal correlations

Commuter traffic, earthquakes and the selection of presidential candidates usually seem to take place in random ways. But investigators of chaos theory who turn to patterns called fractals manage to find order in the midst of such unpredictable events. Now add DNA to the catalogue of things fractallike. "There is some magical phenomenon going on that we just do not understand," says H. Eugene Stanley, a physicist at Boston University.

The orderly patterns of fractals emerge because an incident in the apparently chaotic system is actually correlated with a previous occurrence, much as a long-lasting pocket of slowly moving vehicles can result from one rubbernecking motorist briefly hitting

the brake. Calculations by Stanley's research group and by Wentian Li of the Rockefeller University and Richard F. Voss of the IBM Thomas J. Watson Research Center have shown that the position of nucleotides—adenine, guanine, cytosine and thymine—in a DNA sequence depends to some extent on the nucleotides that preceded it.

The patterns in nucleotide sequences are similar to flicker, or $1/f$, noise (pronounced "one over eff"). These fluctuations are time-scale analogues to the shapes of fractals, such as snowflakes and coastlines, which have the property of self-similarity: the component parts resemble the structure as a whole. The patterns of $1/f$ noise are just as prevalent in nature as their geometric counterparts; they can be found in such diverse phenomena as electric circuits and flood records. "It is a special form of correlations found in natural phenomena and in human behavior," says Voss, who found its presence in music.

If signals are completely random, as are the outcomes of flipping a coin, the results would show a "white noise" signature. Unless the coin is weighted, the probability of heads will be one half regardless of what came before. The outcome would be a collection of random signals similar to those that produce the rushing sounds between FM stations.

In fact, "if you want to store as much information as possible over some length, the best storage method will be like white noise," notes Chung-Kang Peng, a member of Stanley's group. That is because every signal would be completely independent and carry its own message. If genetic data were stored as white noise, the probability of finding the same information along a strand of DNA would decay exponentially the further along the sequence one looked. "If the second nucleotide knows 50 percent of what the first knows, then the third knows only 25 percent," Peng says.

But base pairs in DNA do not seem to

Hot Ants

They live on the edge. Saharan silver ants eat dried corpses, love extreme heat and are given to dramatic, but short-lived, outbursts. When the temperature climbs above 45 degrees Celsius in the desert and the lucky creatures that have not been fried by the sun have bedded down for siesta, the desert floor explodes. Hundreds of ants from a *Cataglyphis bombycina* colony rush out of their burrow and, for several minutes, forage for the desiccated remains of less fortunate arthropods.

Like sunstruck travelers who pause frequently to mop their brows, the scavenging ants rest on sparse vegetation. Some spend as much as 75 percent of their time staying cool this way. Those unable to find a blade of dried grass and climb above ground level into cooler climes may perish for the cause and become food for their colony.

Their remarkable heat adaptation allows the ants to survive in a rough environment. Not only do they feed when others sleep, but by charging out of their holes at the hottest point of the day, Saharan silver ants avoid their principal predator—a lizard that begins to retire just at midday.

Similar thermophilic behavior has been observed in two other genera of desert ant: *Ocyrmymex* in Namibia and *Melophorus* in Australia. "Each real desert in the world has them," says Rüdiger Wehner, professor of zoology at the University of Zurich, whose study of the insects recently appeared in *Nature*. "It is an example of parallel evolution."

All species of *Cataglyphis* are unusual in that they do not leave scent marks. Like colonies of olfactory Hansels and Gretels, most ants leave a trail of pheromones on the way to their food source. The insects then use smell to

guide them between nest and food. If need be, some species can navigate using patterns of polarized light.

Most desert ants also forage at night and during the cooler parts of the day. But at peak heat, "it would not make sense to use pheromones," Wehner explains. Pheromones have high vapor pressure and would evaporate quickly, leaving the ants stranded. In addition, the chemicals are affixed to sand particles, which are easily blown away. It is thought that thermophilic ants such as *C. bombycina* rely mainly on celestial cues, particularly polarized light, as a means of orientation. "They are all individuals," Wehner notes proudly. "They don't follow each other—it is like the explosion of a star."

Wehner's discovery that temperature triggers the ant exodus was accidental. After a long, hot day in the Sahara collecting ants and observing their behavior, he and his colleagues would drive around in a Land Rover, returning the ants to their respective lairs. "We were being very kind to them," he explains. One evening the researchers put some ants back in a tunnel network and "suddenly there was an outburst," Wehner describes.


Because the team had observed these explosions only at midday, Wehner concluded that the heat of the Land Rover had warmed up the ants. Their heightened temperature then prompted the emergence of other foragers. The ants, however, were not fooled for long: they went about 20 meters and then turned back.

Despite burning sun and sand, Wehner will stand and sweat by his ants. "I will work until the end of my life with these creatures," he says. "They are too fascinating." —Marguerite Holloway



SOME DESERT ANTS forage (in this case, for a researcher's cheese) at temperatures above 45 degrees Celsius. Photo: Rüdiger Wehner.

INFORMATION IN




The Henry Holt Handbook of Current Science & Technology

Bryan Bunch

Q: How many times should you shuffle a deck of cards? Find the answer in this new work from Henry Holt Reference (A: in 1990, Columbia & Harvard mathematicians proved a deck is completely random after seven shuffles—anything more is pointless). A comprehensive review and analysis of recent developments, the *Handbook* examines important advances in topics ranging from Astronomy through Mathematics to Technology. Read about The Human Genome Project, new solutions to the Traveling Salesman Problem, Buckyballs, Scanning Tunneling Microscopes and whales with feet. Each topic contains a background section explaining the basic principles underlying the discipline, and then explores specific events in the early 1990s. In addition, each topic also includes tables, charts, and compilations of data such as the worst corporate polluters and complete lists of Nobel Prize winners.

Available Now • \$50.00 • 708 pages • ISBN: 0-8050-1829-8

INFORMATION OUT



Science and Technical Writing A Manual of Style

Philip Rubens (General Editor)

Finally, the first truly comprehensive guide specifically designed for writers of technical material. With sections such as *Audience Analysis*, *Grammar*, *Specialized Terminology* and *Referencing Source Material*, this accessible new manual provides directions and suggestions for providing clear, persuasive prose—text that will make your point and impress your readers. In addition, extensive sections on creating illustrations, tables, charts and diagrams will guide you in presenting information visually, supplementing and reinforcing your writing. *Science and Technical Writing* will help everyone, from beginning students to experienced writers, clarify their expression and structure to create accurate, effective documents. As a style guide, a grammatical reference and a source for ideas, it will be an invaluable resource for all successful technical writers.

Available Now • \$40.00 • 544 pages • ISBN: 0-8050-1831-X

Available from your bookseller, or use the order form below:

Please send me: _____ copies of *The Henry Holt Handbook of Current Science & Technology* (\$50.00 each)
_____ copies of *Science & Technical Writing* (\$40.00 each)

NY, MD, and UT customers please add appropriate sales tax

Name _____

Institution _____

Address _____

City _____

State _____

Zip _____

☐ I have enclosed a check for the correct amount (no postage charge)

☐ Please bill the following credit card: ☐ Visa ☐ MasterCard ☐ American Express

Account # _____

Exp. _____

Signature _____

Mail to: Henry Holt and Company, Inc.
Reference Books Department
115 West 18th Street
New York, NY 10011



Credit Card customers please add:
\$1.95 postage for first book
\$1.45 each additional book

Or call: 1-800-628-9658 (M-F, 9-5 EST)

SRC: SA

occur in a completely random fashion. The researchers applied different statistical techniques to DNA sequences catalogued in GenBank, a storehouse of genetic codes at Los Alamos National Laboratory. They found that the decay is much slower than exponential decrease. Specifically, the sequences show approximate $1/f$ patterns. Roughly speaking, the f here represents the number of bases over which a particular nucleotide repeats.

Along with Stanley's group, Li, a physicist, found that correlations exist in the intron sequences of DNA. Molecular biologists have sometimes referred to introns as "junk DNA" because they do not encode structural information. The real information-carrying regions are located in sequences called exons. Yet unlike introns, exons lack long-range correlation and resemble white noise.

Exactly why intron but not exon sequences show the correlations is not entirely known. The researchers think the long-range correlations—which extend over thousands of base-pair positions—represent a trade-off between efficient information storage and protection against error in the genetic code. Because changing one part forces a change in other parts, there is some redundancy in the code. Thus, the correlations "would give some immunity to error" during transcription, Voss says. Exons, which hold the crucial data, would not exhibit long-range correlations because they need to carry as much information as possible.


Indeed, Voss has turned up some intriguing results based on evolutionary classification. He found that the sequences for organisms lowest on the evolutionary scale (bacteria and bacteriophages) were the least correlated. The correlations increased for higher organisms, reaching perfect $1/f$ patterns in invertebrates. Then the scaling correlations decreased for vertebrates, mammals, rodents and finally primates.

Stanley and his colleagues will soon publish a slightly different result. The correlations increased as they moved up the entire evolutionary ladder. The group discovered that "as you evolve, the long-range correlations become stronger and stronger," Stanley says.

The results indicate that simpler organisms (that is, those with short sequences) would not need the error prevention required to maintain the more complex DNA sequences. "There seems to be a general principle involved," Voss observes. Nature has these fractal and $1/f$ -type fluctuations, so "when systems evolve in nature, they have to build in the same correlations," he remarks.

That may explain in part why music

An \$82.90 Value!



Get the 3-volume QUANTUM MECHANICS LIBRARY

for
only **\$2.95**

as your introduction to the
Library of Science

You simply agree to buy three more books—
at significant savings—within the next 12 months.

No scientific library is complete without this 3-book set covering quantum physics. From conceptual and philosophical problems of quantum mechanics, to practical applications, it's all here in the QUANTUM MECHANICS LIBRARY. Perhaps best of all, these three extraordinary volumes are yours for only \$2.95.



The Principles of Quantum Mechanics, 4th Edition

P.A.M. Dirac.

Here, Nobel Prize winner Dirac concentrates on the theoretical grounding needed to solve problems, and introduces the author's now-standard method of notation. "... the classic text on quantum mechanics."

—W.C. Schieve
University of Texas

Publisher's Price: \$29.95



Speakable and Unspeakable in Quantum Mechanics

J.S. Bell.

Contains all of John Bell's essays on the conceptual and philosophical problems of quantum mechanics, including previously unpublished works, written from 1964-1986. Covers 'impossibility proofs'... nonrelativistic quantum mechanics... 'nonlocality' related to the pilot-wave theory.

Publisher's Price: \$17.95

Quantum Mechanics

P.J.E. Peebles.

This book focuses on applying quantum theory to current problems in experimental physics. Peebles traces the discovery of wave mechanics, shows you detailed computations from perturbation theory, atomic and molecular structure, scattering theory, and reviews the art of numerical estimates.

Publisher's Price: \$35.00

THE LIBRARY OF SCIENCE is the nation's oldest and most respected science book club. Dedicated to serving both professionals and serious amateurs, it surveys the entire field of published sources on scientific subjects to make the very finest works available to club members at substantial savings, with prices starting as low as \$9.95.

MEMBERSHIP BENEFITS: In addition to getting the QUANTUM MECHANICS LIBRARY for only \$2.95 when you join, you'll also receive discounts on books you choose to purchase. Discounts generally range from 15% to 30% off the publishers' prices, and occasionally even more. • Also, you will immediately become eligible to participate in our Bonus Book Plan, with savings of more than 50% off the publishers' prices. • Moreover, our books are always identical to the publisher's editions. You'll never receive an "economy edition" with inferior paper and bindings from us. • At 3-4 week intervals (15 times per year), you will receive the Library of Science News, describing the coming Main Selection and Alternate Selections, together with a dated reply card. • In addition, up to three times a year, you may receive offers of Special Selections which will be made available to a group of select members. • If you want the Main Selection, do nothing, and it will be sent to you automatically. • If you prefer another selection, or no book at all, simply indicate your choice on the card and return it by the date specified. • You will have at least 10 days to decide. If, because of late mail delivery of the News, you should receive a book you do not want, we guarantee return postage.

©Newbridge Communications, Inc.


If reply card is missing, please write to The Library of Science, Dept. 2-DL6/00195, 3000 Cindel Drive, Delran, NJ 08370-0001, for membership information and an application.

Scientific American 9/92

Somebody in the lab

has made an amazing

new discovery.



By bringing affordable, next-generation 3D visualization power to the chemistry desktop, IRIS Indigo gives your lab a decisive, competitive advantage.



IRIS Indigo delivers 30 MIPS, 4.2 MFLOPS, 26 SPECmarks, CD-quality audio, 8-24 bit color, ACE-compatibility, and comes complete with color monitor, keyboard and mouse. Copyright ©1992 Silicon Graphics, Inc. Silicon Graphics, the Silicon Graphics logo and IRIS are registered trademarks, and IRIS Indigo is a trademark of Silicon Graphics, Inc. All other registered and unregistered trademarks are the sole property of their respective owners.

Desktop discovery: IRIS Indigo RISC/PC.

Discover the workstations that run the applications you know, render in 3D, and do it all affordably.

They run popular chemistry software – from chemical structure drawing to word processing to molecular modeling programs. And most importantly, IRIS Indigo™ workstations allow chemists to easily see, study and *understand* the 3-dimensional structural and dynamic features of their discoveries.

IRIS Indigo's award-winning, affordable technology enables application developers to deliver next-generation software to chemistry desktops today. Which means you reduce the time to market, the number of experiments required, and the expenses involved – while building a decisive, competitive advantage.

At prices (starting under \$10,000) that put these powerful, easy-to-use, Macintosh® and PC network-compatible machines on more desks, the IRIS Indigo family enhances a precious scientific resource – the lab chemist – by extending the promise of computer-aided discovery to every desktop.

Call for our free white-paper, "Competitive Benefits of 3D Chemical Modeling," and discover the power of IRIS Indigo for your team.

Call 1 (800) 800-7441. Ext. 4500



SiliconGraphics®
Computer Systems

is pleasing. "One speculation is that music is trying to imitate nature and builds in $1/f$ noise," Voss explains. But a cohesive model to account for the ubiquity of $1/f$ and fractal phenomena eludes researchers. Like many a scientist and harried office worker before him, Li laments: "There is still a lot of work to be done."

—Philip Yam

Bug in a Gilded Cage

All that glitters is sometimes bacterial

Panning for gold in frigid Alaskan streams, prospectors mused about mysterious outcroppings of ore, the mother lodes from which they presumed the precious flecks and nuggets came. If John R. Watterson of the U.S. Geological Survey in Denver is right, the immediate origin of the prospectors' gold has as much to do with biology as with geology. He has found evidence that most of the stream-borne gold particles in Alaska, and possibly elsewhere, were formed by bacteria that accrete a thin skin of pure gold.

Under the electron microscope, many of the gold dust particles Watterson has examined resolve into a lacy mesh of micron-diameter balls and thinner struts. Often the structures are confined to the surface of the gold particles, but some particles are solid masses of them. Watterson has proposed that the structures are the 24-karat castings

of bacteria belonging to the genus *Pedomicrobium*. "The fact that you can find these residual morphologies in the majority of particles I examined suggests that most of it is secondary gold that dissolved and reprecipitated on the cells," Watterson says. "That is quite contrary to the current model."

As far back as Aristotle, earth scientists have generally believed that placer gold, as such particles are called, is the result of mechanical weathering: wind and water chipping away at veins of ore. During this century, geologists have become more aware of how chemical—and sometimes biological—processes can also shape mineral deposits.


Watterson's discovery follows by about 10 years his serendipitous observation that gold solutions are lethal to many soil bacteria. Thin coats of gold, he explains, tend to condense around the bacterial spores, clogging the narrow pores in their cell walls through which nutrients enter.

Those observations spurred Watterson to look for similar accumulations of gold around cells in natural soils. When he looked at placer gold particles from Lillian Creek in Alaska, he was immediately struck by the abundance of *Pedomicrobium* cell castings. "It looked like a Ping-Pong-ball truck had ruptured on the highway," he recalls. "They were all over the place." Many of the particles seemed to be solid masses of gilded cells. Some gold samples from China and South Africa showed similar bacterial shapes, which suggests that *Pedomicrobia* and related bacteria might



GOLD PARTICLE about 0.1 millimeter wide is a 24-karat mass of bacteria that accrete gold around themselves. Photo: John R. Watterson.

ISCAN[®] EYE MOVEMENT MONITORING SYSTEMS



ISCAN's non-invasive, real-time video eye tracking products measure pupil position, size and point-of-gaze. Complete hardware and software available for benchtop, head mounted and high speed data acquisition and analysis.

- Fixation Monitoring
- Neurologic Testing
- Vision/ Perception Research
- Human Factors Eval.
- Aids for the Disabled
- Toxicology/ Drug Abuse

ISCAN, Inc.

125 Cambridgepark Drive
Cambridge, MA 02140, U.S.A.
Tel: (617) 868-5353 Fax: (617) 868-9231



WIRELESS[®]

This star-splashed design embraces the true meaning of education and faith in tomorrow. An upbeat shirt from WIRELESS, the catalog filled with fun, hard-to-find gifts. Sweatshirts, T-shirts, watches, vintage radio shows, videos, memorabilia, cassettes and compact discs, books—everything for your gift list.

For your free one year catalog subscription mail coupon below.

☐ Yes, please start my **FREE** one year **WIRELESS** subscription.

Name

Address Apt.

City

State Zip

Mail to: **WIRELESS**
1000 Westgate Drive, Dept. 8016
St. Paul, MN 55114



**FREE
CATALOG**
Huge Selection
Lowest Prices

Bargain Books

Overstocks, Remainders, Imports and Reprints from all major publishers. Books recently priced at \$20, \$30, \$40—now as low as \$1.95, \$2.95, \$3.95.

Choose from up to 8,000 titles monthly including 600-1,500 new arrivals.

Over 40 subject areas: Science, Biography, Politics, Sports, Literature, Health, Nature, Gardening and more, from yesterday's best-sellers to titles you never knew existed!

Fast Shipment, normally within 48 hours, and a moneyback guarantee.

Please send me your Free Catalog of Bargain Books.

Name

Address

City

State

Zip

HAMILTON

Box 15-259, Falls Village, CT 06031



In the World's Developing Countries, we're planting trees for less than...

...A NICKEL EACH!

If you'd like to help,
please contact:

TREES for the FUTURE
11306 ESTONA DRIVE, P.O. BOX 1786
SILVER SPRING, MARYLAND 20915-1786

1(800) 643-0001

have been involved in the formation of placer gold around the world.

Pedomicrobia can survive the accumulation of gold better than most bacteria, Watterson thinks, because of their unusual reproductive habits. Most bacteria multiply by fission, splitting into separate cells. *Pedomicrobia*, however, often reproduce by budding. Because new cells bud off their parents, remaining attached by short stalks, the bacteria can grow fast enough to escape the gilded cages closing around them. The reliance on budding also explains why these bacteria are good at forming visible gold particles: masses of *Pedomicrobia* cohere to one another rather than becoming isolated cells.

Stephen Mann, a biomineralization expert at the University of Bath, points out that many bacteria can become encased in mineral coatings under the right circumstances. Quite aside from their proclivity for gathering gold, *Pedomicrobium* bacteria accumulate iron and manganese oxides around themselves inside pipes in hydroelectric plants and similar environments rich in dissolved minerals. In general, howev-

er, these mineral deposits are the formless products of chance interactions between metal ions and various molecules leaving the cells. The golden casings around the *Pedomicrobium* cells are extraordinary, Mann remarks, because they are almost perfect in their purity.

The specificity with which the bacteria are gathering gold suggests that some molecule on their cell wall attracts gold complexes selectively. Asked what the bacteria might be doing with the gold, Watterson replies, "That's still an absolute mystery to me." Mann has speculated that the precipitating gold complexes may serve as acceptors of electrons from some biochemical process within the cells.

If there is a cell-wall protein that turns *Pedomicrobia* into gold bugs, Mann continues, it should be possible eventually to clone its gene and transfer it into other bacteria for industrial applications. Many mining companies are already using bacteria to help extract metals from low-grade ores. Surely some 21st-century prospector will try to harness these microscopic Midases as well.

—John Rennie

Making Water Run Uphill

Water usually runs downhill—except in the laboratory of Manoj K. Chaudhury, a senior research specialist at Dow Corning Corporation in Midland, Mich. Along with Harvard University chemist George M. Whitesides, Chaudhury has coaxed liquids to skate almost a centimeter across the surface of a silicon wafer, defying the tug of gravity as they go.

In the past, scientists have shown that liquids can flow uphill driven by changes in the surface of the liquid—say, differences in temperature or evaporation. Chaudhury and Whitesides's work, in contrast, shows that a chemical gradient on the surface of a solid material can induce similar behavior.

A deft bit of chemical legerdemain provides the magic behind the moving droplets. Chemists have long known that some materials are hydrophobic, or repel water; others are hydrophilic, or attract it. By carefully diffusing a gradient of silane vapor over a polished silicon wafer, the researchers created a surface that becomes more hydrophilic as the layer of silane thins.

A milliliter or so droplet of water deposited on the hydrophobic end of a wafer is consequently drawn to the other side. Even when the wafer is tilted upward at a 15-degree angle, that pull overwhelms both gravity and the viscosity (or stickiness) of the liquid and draws the droplet uphill.

Although Chaudhury and Whitesides picked silicon and water for their first experiments, other combinations of solids and liquids perform the same trick. Acetonitrile, which has a low viscosity, is particularly speedy, moving at a rate of eight to nine millimeters per second in contrast to water's pace of one to three millimeters per second. If the researchers can lengthen the distance over which their droplets travel to several centimeters, the technique may have applications in sensor technology.


Chaudhury and Whitesides began the project to explore the fluid dynamics at the so-called three-phase boundary, the realm where liquids, solids and gases interact. (The droplets turn to vapor at the boundary between the liquid and solid substrate.)

The researchers' results nicely illustrate a theory proposed three years ago by Françoise Brochard-Wyart, a French physicist. Although Chaudhury had confidence in the theoretical explanations, he says, "it was really very gratifying when we saw the drop actually move."

—Elizabeth Corcoran


Electricity Powers A Lot Of Things. Not The Least Of Which Is The GNP.

If the U.S. economy is to continue to grow, electric energy efficiency must play a role. As electricity assumes a larger share of our total energy use, we are achieving greater economic output with greater energy efficiency. Since 1973, U.S. Gross National Product has increased by 51%, while energy consumption per dollar of GNP has declined by 28%. That's due largely to advances in electric technologies.



Electric infrared processes dry paint ten times faster and cut energy use by as much as 90%.

Consider what we're doing for the American auto industry. The introduction of



Advances in electric technologies are enabling us to be more energy efficient while achieving greater economic output.

electric infrared drying has allowed auto manufacturers to cut paint drying time from 4 minutes to 31 seconds, while cutting energy use by as much as 90%. This process has enabled them to do the job ten times faster while producing a more uniform, durable finish. And by switching from traditional methods, cut down the number of emissions that are

being released into the air.

We all stand to gain from this kind of progress. Many relevant technologies already exist. And, with more on the way, it only promises to get better. For more information, call Edison Electric Institute at 1-800-438-8334 or, in Washington, at (202)-508-5002.



ELECTRICITY
Taking you into the future.

Treat yourself (or your doctor)



to these medically friendly titles...

THE BEST OF MEDICAL HUMOR

A Collection of Articles, Essays, Poetry, and Letters Published in the Medical Literature
Compiled and edited by Howard J. Bennett, M.D.

Physicians poke fun at themselves, their patients, medical education, and the health care system.

"Smiles at you from the shelf..."
— American Journal of Ophthalmology
"... a wonderful new book."
— Journal of the American Medical Association

\$25.00 hardcover, 240 pages, illustrated

A LITTLE BOOK OF DOCTORS' RULES

Edited by Clifton K. Meador, M.D.

Over 400 rules, observations, and helpful tips for medical students and physicians about patients and the practice of medicine. A perfect gift.

\$9.95 paperback, 128 pages,
available November 1992

DICTIONARY OF MEDICAL ACRONYMS & ABBREVIATIONS, 2nd ed.

Compiled by Stanley Jablonski
Selected by the American College of Physicians for its "Library for Internists" as the best medical acronym book available. This new edition contains thousands of new entries.

\$15.95 paperback, 320 pages,
available September 1992

NEUROANATOMY A Functional Atlas of Parts and Pathways

by Ray Poritsky, Ph.D.

The author-artist illustrates the brain and spinal cord in three-dimensional drawings, allowing the reader to learn by coloring and labeling the structures.

\$27.95 paperback, 280 pages

HEART FAMILY HANDBOOK

A Complete Guide for the Entire Family of Anyone with Any Heart Condition—to Help Make It the Speediest, Most Complete Recovery Possible

by Jane Schoenberg and JoAnn Stichman

Dozens of medical experts in cardiac care, including world-renowned cardiologists and cardiac surgeons, were interviewed by the authors and share their expertise and wealth of experience in helping patients and families facilitate the recovery process.

\$16.95 hardcover, 240 pages

THE PORTABLE PEDIATRICIAN

by Howard Markel, M.D., Jane A. Oski, M.D., Frank A. Oski, M.D., and Julia A. McMillan, M.D.

The John Hopkins Children's Center Alphabetically arranged (from Abdominal masses to Zoonoses) charts, tables, discussions, lists, and tips on over 240 common pediatric problems.

"Things I forgot to remember..."

\$35.00 paperback, 400 pages

TO ORDER, CALL TOLL FREE 800-962-1892

HANLEY & BELFUS, INC.

MEDICAL PUBLISHERS

210 South 13th Street Philadelphia, Pennsylvania 19107

Bent Light

*Gravitational distortions
illuminate dark matter*

Imagine walking into a pitch-black room and being asked to describe the people standing silently inside. That is the kind of quandary faced by astronomers trying to learn more about dark matter, the unseen material thought to account for as much as 99 percent of the mass of the universe. Amazingly enough, they are having some success in this undertaking.

All knowledge of dark matter derives from the fact that it has mass and so produces a gravitational field. In the past few years several astronomers have utilized that property to "see" dark matter indirectly. Because gravity bends, or lenses, light (according to Einstein's theory of relativity), dark matter can be discerned when it distorts one's view of luminous matter. "The potential of this technique has barely been explored," says Scott D. Tremaine of the University of Toronto, a veteran of dark matter searches. "I'm very optimistic about it."

The existence of dark matter was posited during the 1970s. At the time, astronomers demonstrated that the outer parts of galaxies rotate much faster than theory would predict. If galaxies consisted only of luminous matter, they would quickly fly apart. Similarly, individual galaxies in clusters move at inexplicably high velocities, and yet the clusters have not dispersed. The only plausible explanation for such behavior is that galaxies and clusters contain a healthy dose of dark matter that provides the gravitational glue needed to hold them together.

Clumps of dark matter probably helped to instigate the coalescence of the visible galaxies and clusters of galaxies. The abundance of dark matter determines the mass, and hence the fate, of the universe. Current theories predict there should be just enough dark matter to overcome the present cosmic expansion.

A little astronomical sleuthing has at last begun to unravel the riddle of where the invisible matter resides and how much of it there is. Although galaxy motions cannot provide much information about where the dark matter is located, observations of gravitational bending of light can. To that end, J. Anthony Tyson and Richard Wenk of AT&T Bell Laboratories and Francisco Valdes of the National Optical Astronomy Observatories have focused their attention on faint, blue galaxies several billion light-years from the earth.

Effortlessly contour and colormap surfaces generated from random sample points.

Pull a lever to adjust a variable, as the revolutionary Graphics Equalizer™ animates changes to your curve-fits and formulas on-screen.

Automate your presentation with master slides, outliner, speaker & handout notes.

Pop up the extraordinary graph gallery of over 160 business, technical, and statistical graph types.

Access your numbers instantly from your favorite Windows or DOS spreadsheet.

Hot-link multiple data files directly to Stanford Graphics' proprietary 4-D 70-trillion cell spreadsheet.

Animate the 3-D rotation of your graph and its actual data.

Instantly identify & modify your data points with the Intelligent Data Cursor™. Simply drag a bar or line to change a cell in your spreadsheet!

Click on the Calculator for on-the-fly business, financial or scientific computations.

**NEW!
VERSION
1.01**

STANFORD GRAPHICS FOR WINDOWS

Finally...presentation graphics with numerical clout!

"Stanford is a breakthrough product that combines technical charting, data analysis, and presentation tools. Basically, it's fabulous!"

— PC Magazine
March, 1992

"Stanford Graphics offers the best of both worlds. It combines top-notch presentation features with unprecedented technical sophistication. We were greatly impressed."

— Infoworld,
Sept., 1991

"Stanford Graphics slams the competition..."

— Infoworld
April, 1992

Does your presentation software give you all beauty and no brains? Where is the substance behind those pretty pictures? What if your presentation program had enough data-handling muscle to meet the needs of business *and* statistical *and* scientific users?

With on-screen curve fitting, formula solving, statistics, and data manipulation, Stanford Graphics advances both the art and science of technical presentations.

All this and Windows too. Stanford Graphics takes full advantage of the latest



STANFORD GRAPHICS™
PRESENTATION & ANALYSIS

© 1992 SCIENTIFIC AMERICAN, INC

in Windows 3.0 and 3.1 technology, with sophisticated DDE, OLE, and TrueType font support.

Stanford Graphics for Windows:
Presentation and Analysis.

Finally, form *and* substance.

3D 3-D VISIONS
Call Toll Free for Details
1-800-729-4723

2780 Skypark Drive, Torrance, CA 90505
Tel: (310) 325-1339 FAX: (310) 325-1505

Each square degree of sky contains more than 300,000 blue galaxies, enough to provide a convenient backdrop against which to search for dark matter. Gravity from clusters of galaxies located closer to the earth deflects light from the more distant blue galaxies, altering their oval forms into elongated ellipses and arcs. The shapes and locations of the distorted blue galaxies indicate the abundance and distribution of matter in the nearer cluster.

Tyson finds that dark matter follows the overall pattern of visible matter but that its basic unit of "clumpiness" measures about 300,000 light-years across, distinctly larger than a typical galaxy. Vahé Petrosian of Stanford University, who has made gravitational-lens observations much like those done by Tyson, notes that his work shows dark matter to be "more highly peaked"—that is, more centrally concentrated—than the luminous matter in clusters. That finding underscores just how little is known about the origin of structure: most cosmological models predict that dark matter should be spread more evenly than the visible galaxies.

Dark matter may form halos that stretch far beyond the visible edges of clusters of galaxies. If so, it "could easily close the universe," Tyson says; in other words, its gravitational pull could eventually halt the expansion of the universe. Tremaine adds that "there is no reason why you can't have dark matter clusters with no visible frosting at all." Much of the mass of the uni-

verse could reside in unseen clumps that are devoid of luminous galaxies.

So far Tyson can only set a lower limit to the density of the universe of about 0.2 times the critical value needed to overcome the expansion. Improving that estimate will require measuring the total extent of the dark matter halos. The outer parts of clusters cover broad patches of sky, so surveying them demands the use of large light detectors that can make use of a telescope's entire field of view. In the past the available detectors were too small to do the job properly. Tyson plans to begin using a larger detector this fall.

If there is enough dark matter to close the universe, then dark matter halos must fill the space between clusters of galaxies. In that case, one could look at an apparently empty section of sky and still see signs of gravitational distortion. Such observations might also chance on clusters consisting solely of dark matter. Conversely, if no such distortion appears, one could conclude that the density of the universe is well below the critical value—a finding that would send a lot of theorists back to their blackboards.

Tyson has looked at 40 blank fields and claims he already has an answer, but he is not telling anyone until he double-checks his results. A follow-up search using the updated detectors will take about a year. Until then, astronomers will have to be content to believe that there is light at the end of the dark matter tunnel.

—Corey S. Powell

Population Pressure

The road from Rio is paved with factions

At times it seems there are as many opinions concerning what—if anything—to do about population growth as there are people on the planet. Bitter schisms often divide groups arguing for women's rights, family planning and health, environmental protection, reduced consumption of natural resources, economic development and population control. "We are hearing the same things over and over and over again. Nothing has changed in the past 20 years," laments Jodi L. Jacobson, senior researcher at the Worldwatch Institute. "There is a lot of controversy between groups that should not be at odds with each other."

Indeed, the topic almost proved too controversial even for delegates to the United Nations Conference on Environment and Development (UNCED) in Rio de Janeiro this past summer. At the official meeting, held some 30 miles outside the city, speakers, including executive director of the U.N. Environment Program Mostafa K. Tolba and Norwegian prime minister Gro Harlem Brundtland, urged immediate steps to address population growth. "Poverty, environment and population can no longer be dealt with, or even thought of, as separate issues," Brundtland warned.

Meanwhile in downtown Rio, participants on a panel at the Global Forum—a meeting attended by some 2,000 non-governmental organizations as a parallel summit to UNCED—viewed efforts to control population as an infringement on women's rights, as a way to avoid eradicating poverty in developing countries and as a means of silencing Third World poor. "It is like triage: kill off the weak," said Vandana Shiva, director of the Research Foundation for Science, Technology and Natural Resource Policy in Dehra Dun, India. "Fascism never sleeps permanently."

Such rancorous debates are hardly new. Since 1968, when Stanford University biologist Paul R. Ehrlich emphasized the link between population growth and environmental degradation, experts have been contesting the merits of population "control" versus encouraging economic development as the means of slowing population growth. In that time, the number of human beings has increased from 3.5 to 5.5 billion.

Although the growth rate has actually declined from about 2.0 to 1.7 percent during the past 25 years, some 97 million people will be added annually in this



FAINT BLUE ARCS (center regions) are distant galaxies whose light has been bent by the cluster of orange galaxies (foreground). The distortion reveals the location and quantity of dark matter in the cluster. Image: J. Anthony Tyson.

What's a Mercury Villager?



That's a Loaded Question.

Send for a FREE Preview Kit and we'll give you loads of answers.

The new Mercury Villager has comfort and convenience features you just won't find in most other minivans. Like its innovative rear seat which folds and slides forward in a flash for added cargo space. Or its standard four-wheel anti-lock brakes. You can add a host of luxury car items too, like leather seat trim, power moonroof, power driver and passenger

seats and even keyless entry. So, return the attached reply card or call 1 800 446-8888 today for a FREE Preview Kit. After you see how much Mercury Villager has to offer, other minivans may leave you feeling a little empty.




All This And The Quality Of A Mercury.

Just Released...

SCIENTIFIC AMERICAN

Cumulative Index

on Computer Disk



Order SciDex™ today and turn your *Scientific American* library into an invaluable reference tool. You can have 44 consecutive years of scientific and technological insight at your fingertips within seconds.

- **Easy to Use** ■ **Fully Cross-Referenced**
- **Article Abstracts** ■ **530 Issues**
- **Over 4,300 Feature Articles Indexed**
- **Over 5,000 Authors** ■ **Print Your Search**
- **Over 43,000 Topical Entries**
- **Annual Renewal at Discount Price**

See advertisement on page 173 for complete details and order form.

René Brundt

William Shakespeare

Le da Vinci

L.v. Beethoven

Mozart

E. H. H. H.

Vincenzo

Edgar Allan Poe

Elizabeth Barrett Browning

René

THEY'RE ENTITLED TO
GET CREDIT FOR
THEIR CONTRIBUTION
TO THE ARTS.
AND YOU MAY BE, TOO.

If your business contributed money, personnel, goods or services to the arts last year, you may be eligible to win a Business in the Arts Award.

To nominate a business, write: Business Committee for the Arts, Inc., 1775 Broadway, Suite 510, New York, NY 10019. Or call 212-664-0600.

B BUSINESS COMMITTEE
FOR THE ARTS, INC.



The sheep that came home for winter.

Winds here can scream up to 70 miles per hour. Temperatures plummet to 20 below. This is Stillwater Valley, Montana. The winter home of a small, fledgling herd of Bighorn Sheep. A herd that for years has been struggling to survive. But people who work nearby are providing medication and food to nurture, protect and save these majestic animals. Hopefully helping each generation become healthier. Each generation larger.

Do people look after their neighbors so nature can grow strong?

People Do.



For more information, write: People Do-S, P.O. Box 7755,
San Francisco, CA 94120

© 1992 Chevron Corporation



The United States Academic Decathlon

Learn about the challenge! The U.S. Academic Decathlon is a national scholastic competition which provides recognition and reward for solid academic endeavor. Individual high school students compete in their grade point category: A, B, or C and below. Teams compete across a range of events for important awards: scholarships, medals, pride in themselves, their teams and their schools.

The greatest award, for young people, is educational achievement. The U.S. Academic Decathlon seeks renewed respect for schoolwork at all levels, started and sustained during the important pre-college years, open not just to the intellectual elite but to a spectrum of able students. In 1991, 25,000 young people took part in Decathlon competitions culminating in an exciting National Finals.

Scientific American is pleased to join American Airlines, The Krausz Companies, Inc., GTE, D.C. Heath, the Lennox Foundation, Ronald McDonald's Children's Charities, Northrop, The Psychological Corporation, Raytheon, TRW and others in sponsoring the U.S. Academic Decathlon. All of us salute the many local companies and organizations which help their schools and their children participate. *We invite you.*

Please send for more information; your participation is needed, and your effort and support will have major impact. Thank you.

Yes, please tell me more!

Name

Address

City

State Zip

Mail to: **USAD**
c/o Scientific American
415 Madison Avenue
New York, NY 10017-1111

decade. Depending on rates of growth, current projections put global population between 10 and 12.5 billion by 2050. And 97 percent of this increase is anticipated to take place in the developing world, where 32 percent of the population is younger than 15 years—that is, just coming into childbearing years.

For the first time, organizations such as the National Academy of Sciences and the Royal Society of London have become outspoken about the need to focus on population growth. Before the Earth Summit, the institutions issued a joint statement warning against overpopulation. "Population control is a codeterminant with control of environmental damage," comments Nobel laureate Henry Kendall of the Union of Concerned Scientists, which is starting a population project. "People who take issue with control of population do not understand that if it is not done in a graceful way, nature will do it in a brutal fashion."

Yet so far many environmentalists have steered clear of population. Some have worried about losing members if they endorsed policies that touch even indirectly on abortion. Despite his outspoken support for family planning during the late 1960s, President George Bush cut all U.S. funding to the U.N. Population Fund in 1985. He also eliminated money for international family-planning programs that provided abortion counseling or services (even if U.S. money was not directly spent on abortion). "Whenever you mention family planning and contraceptives, you have pushed the abortion button," notes Eric Washburn of the Natural Resources Defense Council, which is beginning to incorporate population into its programs.

Environmental associations have also been loath to tackle immigration issues. Some forays into this area have been criticized. In California, for instance, Sierra Club joined other groups, including Population-Environment Balance, a nonprofit organization based in Washington, D.C., to support "sustainable development and quality of life." Population-Environment Balance endorses a "replacement level" policy: 200,000 people would be let into the U.S. every year, a number equivalent to the number who emigrate. The short-lived coalition was attacked as potentially racist.

In the aftermath of Rio, however, environmental organizations seem to recognize that population is an issue they can no longer ignore. "Everyone is coming to realize that it is a piece of the environmental problem," notes Deborah Moore, staff scientist at the Environmental Defense Fund (EDF). Some are concentrating on reducing consumption—for example, industrialized na-

tions consume 75 percent of all energy used but contain only 25 percent of the population, according to the EDF. Others are focusing on immigration, on carrying capacity (the population any region can support without degradation) and on lobbying to increase funding to family-planning programs.

An influx of money is catalyzing some population programs. Several foundations have set aside grants for population projects within environmental groups. The Pew Charitable Trusts, for example, recently invited several of the country's largest environmental associations, including Ducks Unlimited, as well as religious and population groups, to submit proposals. Pew may give out as many as 10 grants of \$400,000. "There is a rush on," Moore observes.

But Moore and others worry that too much haste may lead to an oversimplification of issues: "We don't even have a good understanding of cause and effect." High population density does not necessarily result in environmental degradation, Moore maintains. Deforestation in the Amazon rain forest is not so much related to an increasing population as it is to land ownership and demand for timber and beef, she says; in contrast, water pollution in urban areas such as São Paulo can be directly linked to overpopulation.

In addition, experts in the environmental, scientific and family-planning communities are concerned that human rights and economic development have been ignored by some population-control efforts. Reports of forced sterilization (in such countries as China and India), inadequate health services and the use of local populations as guinea pigs in contraceptive tests have made many women deeply distrustful of family-planning services. "In the past, governments have really run roughshod over the rights of women," agrees Alex Marshall of the U.N. Population Fund.

Many plans have also ignored family health, notes Jacobson of Worldwatch. Birth rate may be down in Thailand, she points out, but sexually transmitted disease is on the rise. And in Brazil, so few forms of birth control are available that 41 percent of women choose sterilization as their means of contraception. Family planning "will be moot unless it has support from the bottom up," observes Ellen Starbird of the U.S. Agency for International Development.

Starbird and others argue that family-planning projects must take into account the concerns of people using, or avoiding, the services—such as some of the women speaking in Rio. "I wish some of the Northern groups were more sensitive to the Third World women,"



UNCONVENTIONAL WISDOM

They offer counsel and comment on the week's events from Washington. They do it with wit and dissension. They're The McLaughlin Group.

The Group comprises (clockwise from left) Jack Germond, Clarence Page, John McLaughlin, Eleanor Clift, Morton Kondracke and Fred Barnes.

They're contentious and contagious. So, tune in to The McLaughlin Group.

Made possible by a grant from GE.

The *McLaughlin* Group

Check your local listing for station and time.



We bring good things to life.

GIVE IT SOME THOUGHT

*It collects your
thoughts. It
takes your notes
and memos.
Yet the unique
Olympus L400
Microcassette™*



*Recorder weighs
less than your
wallet and bides
in your hand.
So consider the
possibilities. All
from Olympus.*

N e v e r m i s s a n o t h e r O.

OLYMPUS

Call 1-800-221-3000, ext. 85 for our free taping tips brochure. Or write Olympus Corporation, Crossways Park, Woodbury, NY 11797-2087.

THE SCIENTIFIC AMERICAN LIBRARY
IS PROUD TO ANNOUNCE THE PUBLICATION OF
NOBEL LAUREATE HAROLD E. VARMUS AND
ROBERT A. WEINBERG'S

MUCH-ANTICIPATED NEW BOOK,

GENES AND THE BIOLOGY OF CANCER

AVAILABLE IN BOOKSTORES: SEPTEMBER, 1992

SPECIAL OFFER FOR
SCIENTIFIC AMERICAN READERS:
GET A FREE 15-DAY APPROVAL COPY OF
GENES AND THE BIOLOGY OF CANCER
(SEE PAGE 40 FOR DETAILS)

concur John Rowley of the International Planned Parenthood Federation.

Despite the means by which some population policies have been implemented, many such efforts have won significant support among Third World feminists who acknowledge that the services often improve women's economic status and health—particularly when they are integrated with family health clinics and education. "Contraception is one of the tools to economic development," comments Wanga G. Mumba, executive director of the Environment and Population Center in Lusaka, Zambia.

Even so, the U.N. estimates that as many as 300 million women have no access to family planning, although 150 million want to stop or delay childbearing. According to the Population Crisis Committee (PCC), population would become stable around nine billion by 2050 if 75 percent of couples used family planning. This goal could be met if the world's annual family-planning budget were \$10.5 billion by 2000, PCC reports. (The world now spends \$4.5 billion on such services; 85 percent of the money comes from developing nations.)

The passionate discussions at the Earth Summit may result in some steps in the right direction. Initially, population was absent from UNCED's agenda: developing countries said they would not discuss population unless developed countries discussed consumption.

One of the signed documents includes a \$7.1-billion figure in a much contested chapter on "demographic dynamics and sustainability." But critics say the wording in Agenda 21, the 800-page strategic environment and development outline for the next century, makes the issue trivial. They argue that interference by the Vatican and by some feminist groups weakened the language by deleting the word "contraception" and the phrase "family planning."

Others counter that the important ideas were preserved. "We were looking for agreed language that would not give away the substance, and we did that," explains Marie Coleman from Australia, a senior visiting fellow at Pennsylvania State University, who chaired the group that drafted the chapter. "For the first time, we have language that locks in the Muslim countries and the Vatican."

But Agenda 21 means little unless the money is spent, women's concerns are taken into account and diverse organizations collaborate. "It seems a ridiculous problem," comments Washburn of the National Resources Defense Council. "If there are any groups that should be working together, it is the environmentalists, the population groups and the feminists." —*Marquerite Holloway*

JOIN THE GLOBAL INTUITION NETWORK

The purpose of The Network is to promote the applied use of intuition in decision making; share new knowledge on how to use this brain skill as it becomes known; and to promote ongoing research on intuitive processes for practical use in organizations of all types.

The focus of the Network is global, interdisciplinary, practical as well as academic, and on all levels and types of education. If you would like to join and participate in this Network (which is free), contact: Dr. Weston H. Agor, Director, Global Intuition Network, P.O. Box 614, UTEP, El Paso, Tx. 79968 or FAX 915-747-5111. The Global Intuition Network is supported by a grant from The Charlson Research Foundation.




T CATCH A GLIMPSE OF THE FUTURE

Two world-renowned scientists explore the revolutionary developments — as well as the exciting challenges ahead — in science's momentous battle against cancer



Fluorescent dye shows exchange of chromosomal material




X-ray crystallographic analysis of human growth hormone protein

GENES AND THE BIOLOGY OF CANCER

HAROLD E. VARMUS AND ROBERT A. WEINBERG

EXAMINE IT
FREE
FOR 15 DAYS



Despite major research breakthroughs, the incidence of cancer is as high as ever. In fact, almost one in four people in developed nations fall victim to some form of cancer.

Nobel laureate Harold E. Varmus and Robert A. Weinberg take you on an in-depth look at the nature, origins, and mechanisms of cancer as they recount how the study of cancer has illuminated our understanding of normal biological functions.

Learn why there are so many types of cancer. Where abnormal cells come from. How cancer develops. Why the tendency toward cancer may be passed from generation to generation.

- Why do normal cells stop growing, while cancerous cells proliferate indefinitely?
- How do chemical and environmental carcinogens transform cells and induce cancer?
- Do abnormal cells come from within the organism or are they introduced, like an infection, from another organism?
- When a cancerous cell replicates, do all its descendent cells have the same genetic composition?
- How can you tell whether a tumor cell is caused by an overactive growth-promoting gene or a damaged growth-suppressing gene?

A NEW BOOK BY TWO PREEMINENT SCIENTISTS

Professors Varmus and Weinberg explain the strategies organisms use to govern cell growth and differentiation. They demonstrate how these strategies fail in cancerous cells. And they explore the clinical implications of these developments in growth control.

YOURS TO EXAMINE FREE FOR 15 DAYS

Best of all, you can examine **GENES AND THE BIOLOGY OF CANCER** in your home FREE for 15 days. Without risk or obligation. By doing so, you'll be introduced to the Scientific American Library. Every other month, you'll have the opportu-

nity to review — completely risk-FREE — an authoritative, hardcover volume like:

- **VIRUSES** by Arnold J. Levine
- **CHEMICAL COMMUNICATION: The Language of Pheromones** by William C. Agosta
- **JOURNEY INTO GRAVITY AND SPACETIME** by John Archibald Wheeler
- **FROM QUARKS TO THE COSMOS** by Nobel laureate Leon Lederman and David Schramm


Each Scientific American Library volume is written by a leading scientist in accessible, compelling prose. Each

includes dozens of full-color photos, illustrations, diagrams, and charts. Each covers a major scientific topic. And each is yours to examine FREE for 15 days. Of course, there is no minimum number of books to buy. And you can cancel at any time.

MAIL THE RESERVATION CARD TODAY

Don't delay. Return the completed reservation card and we'll send you **GENES AND THE BIOLOGY OF CANCER** for a FREE 15-day trial examination. Send no money now. Just mail the postage-paid reservation card today.

If the card is missing, call toll-free
1-800-345-8112



Secondary tumors, or metastases, appear as small colored "hotspots" against the background blue of a cancer patient's chest in this radioisotope scan.



or write: Scientific
American Library
GENES AND THE

BIOLOGY OF CANCER Offer
P.O. Box 646, Holmes, PA 19043

SCIENTIFIC AMERICAN LIBRARY



MARK TILDEN constructs tiny robots with parts from dead Walkmans, discarded calculators and other high-tech trash. Photo: Michael Messner.

Photovores

Intelligent robots are constructed from castoffs

When the lights turn on in Mark Tilden's office at the University of Waterloo in Ontario, visitors are sometimes startled: a roller edges forward here, and a hopper springs into the air over there. Tilden made these light-craving robots based on what he calls BEAM principles (for biology, electronics, aesthetics and mechanics). Lovingly polished brass and stainless steel frames embrace motors that once ran Vanilla Ice tapes, solar cells that powered pocket calculators and gears that drove printer spools.

In spite of his university setting, Tilden operates outside the academic territory of peer review and incremental thought. No anatomic charts of his elegant, spidery creations of brass and silicon grace journal pages. Yet the robots have invaded not only Tilden's office but also high school robot workshops and even the Third Artificial Life Conference, held in Santa Fe, N.M., in June.

By using only recycled electronic and mechanical equipment, Tilden forces himself and his cadre of disciples to be imaginative, to avoid falling back on the construction of special-purpose solutions that lack biological élan. BEAM robots must not cost more than a few dollars to build. Out the door go million-dollar grants and development con-

tracts. Occasionally, but only in the extremity of empty trash bins, will he resort to off-the-shelf components.

A hardware engineer at the university's Computing Facility by day and a robot designer by night, Tilden has built no less than 32 different robots from electromechanical castoffs. The earliest are the simplest; the most recent actually walk. Do they represent some kind of evolutionary process? "Evolve? Who wants to evolve? If you can't survive, you're Spam in a can!" Tilden exclaims.

Survival means continuing to operate year in and year out with no maintenance. It means being strong enough to withstand a fall from a shelf, and it means, above all, functioning with a minimum of brains. Among Tilden's simplest creatures are solarrollers, wheeled robots that roll forward under a desk-lamp "sun." Hoppers sit quiescently for long periods, steadily accumulating energy in a capacitor, then leap a foot or more into the air. Pointless now, perhaps, but potential pre-adaptive behavior for a later BEAM bug.

At the Santa Fe meeting, a light-seeking robot (Tilden calls them photovores) known as the turbot was a favorite. The tetrahedral turbot moves by rolling from one of its four triangular faces to another, pried up and over by a rotating arm. In strong light the turbot stays within a confined area, but in dimmer light it balances more frequently on its edges and develops chaotic movements that take it further afield.

Through high school robotics workshops, Tilden has inspired a new gener-

ation of students. The better students enter their latest creations in the annual Robot Olympics conducted by Tilden. Nonstudents from Japan and the U.S. also bring their BEAMish bugs to compete in high jumping, rope climbing and sumo wrestling, as well as in the micromouse maze and solaroller races. Conducted every spring, the Robot Olympics have created a flood of mail asking for Tilden's Olympic rules and guidelines booklet.

Reveling in the powers of the simplest circuits, Tilden wonders if his creatures will ever need anything like a brain. He tells the story of a fly that buzzed about a particular window in his apartment for too many days last winter. He moved his solar-powered window cleaner there to keep the fly continually busy. Then, after half an hour of breadboarding, he completed the basic circuit for a mechanical fly-trap of the Venus persuasion. Within a few hours, the BEAM window cleaner had teased the fly into the BEAM trap, where metal petals closed slowly over it. On the following day, Tilden found the fly's carcass below the window, where, in time, the BEAM floor cleaner would pick it up.

Tilden is suspicious of the usual approach to artificial intelligence. "You can't slam-dunk a brain into a robot body and expect it to do anything useful or interesting," he says. He accuses AI engineers of loading too many low-level control functions into software when simple circuits or even mechanisms might handle them.

Tilden is not alone in feeling that robotics is best tackled from the bottom up. Rodney Brooks, director of a micro-robotics laboratory at the Massachusetts Institute of Technology, shares that view. Brooks has pioneered simple walking robots that use only a few microchips in place of a complete computer. In fact, Tilden traces the development of his passion for lifelike mechanisms to a talk that Brooks gave in October 1989. Brooks described his notion of subsumption architecture. This view holds that complex behavior becomes possible in a robot only when there are simpler behaviors present that it may subsume.

Tilden finally met his mentor at the Santa Fe conference, where his menagerie of creatures unintentionally stole the show from the single large insect that Brooks brought to the conference. For the future, Tilden ponders an "ecology" of robots fulfilling their biological, electronic, artistic and mechanical mandates in concert. They will, one hopes, be better disposed to one another than to flies.

—A. K. Dewdney

"Nothing we can do can kill the thing off."



— Bobby Winston, Producer, San Francisco Shakespeare Festival

"In a heavy fund-raising period, we've run our LaserJet 24 hours a day for days at a time. It just will not die," says producer Bobby Winston. "It's truly a phenomenal piece of engineering."

For years, experts have applauded LaserJet performance. In 1991 alone, HP LaserJet printers won *Printer of the Year* from Byte and InfoWorld; *Product of the Year* from Datamation and Byte; *Most Valuable Product* from PC Computing; *World Class* and *Best*

Buy from PC World; and *Readers' Choice* from Publish.

All five models have received many accolades, from the 4 ppm personal LaserJet IIP to the 17 ppm LaserJet IIISi. With HP, you can get genuine PCL5 printer language, with scalable typefaces. And HP's exclusive Resolution Enhancement technology, for improved print quality. Prices range from \$1,249 to \$5,495.*



To hear what other customers have to say about HP LaserJet printers, call **1-800-752-0900, Ext. 2988** for a free video. You'll see why the LaserJet is a hard act to follow.

HP LaserJet Printers.



*Suggested U.S. list prices.
© 1992 Hewlett-Packard Company PE12251



The Frustrations of a Quark Hunter

Roy F. Schwitters teeters on the edge of his chair, glaring across the hearing room at Senator Dale Bumpers of Arkansas. As director of the Superconducting Super Collider (SSC), Schwitters is fighting for the life of the project. If the construction of the SSC is completed, it will be the world's most powerful particle accelerator and its most expensive scientific instrument. But in mid-June, the House of Representatives voted to kill the \$8.3-billion program. Today Schwitters must try to persuade 51 senators to restore the funding.

"Do you know the projected cost of the dipole magnets?" Bumpers asks, referring to the magnets that may someday guide energetic particles through the SSC's 87-kilometer-long circular tunnel under the prairie near Dallas.

"Roughly \$150,000 each," Schwitters replies.

"According to Dr. William Happer," Bumpers snarls, the Japanese think "the cost of building the magnets will be two to four times higher than [you] estimate."

"That was an early projection, sir," he exclaims in a firm voice steeped in anger. "It was made before we had the opportunity to talk things over with the experts who are now building the magnets." The source of the controversy, William Happer, is sitting in the audience. Bumpers asks him to stand, and Happer, a bit embarrassed, admits he now agrees with the figure of \$150,000.

For three hours, Schwitters parries the attacks of Bumpers and defends the SSC before a panel of nine senators. Schwitters is not battling alone. Five Nobel Prize winners attended the hearing to lend their weight to his cause. Later in the day Schwitters meets with President George Bush, who promises to get personally involved. Yet with the public complaining about government spending in an election year, can Congress resist the symbolic gesture of eliminating its most costly science program?

Schwitters, now 48, feels a deep obli-

gation to provide a productive future for particle physics by completing the giant accelerator. In the 1970s he was one of the key investigators in a discovery that confirmed the existence of fundamental particles called quarks. As a professor of physics at Harvard University during the 1980s, he directed the construction of the premiere particle



ROY F. SCHWITTERS is fighting congressional opposition to the construction of the Super Collider underneath the Texas prairie. Photo: David Sams.

detector for the world's highest-energy accelerator at the Fermi National Accelerator Laboratory (Fermilab). And if Schwitters succeeds in building the Superconducting Super Collider, he may just preside over the discovery of the factor that gives all particles mass.

Driving half an hour through Ellis County, Tex., from the SSC headquarters to the magnet-testing facility, Schwitters welcomes an opportunity to talk about himself instead of the project for a change. He grew up in Seattle, and like many children of the Sputnik era, he learned to enjoy launching model rockets, building radios and experi-

menting with chemistry sets. "My parents were very supportive," he says. "I was always blowing things up."

Although neither of his parents had a college education, they encouraged him to go to the Massachusetts Institute of Technology. It was the first time he had traveled east of the Mississippi. During his college years, Schwitters got the "world's greatest summer job" as a guide at Mount Rainier National Park in Washington State. There he met Karen Chrystal; they were married at the end of Schwitters's junior year. He remained at M.I.T. for a doctoral degree in physics and performed the experiments for his thesis at the Stanford Linear Accelerator Center (SLAC). In 1971, as Schwitters finished his dissertation, his wife gave birth to the first of their three children.


Schwitters joined SLAC as a postdoctoral student, wanting to continue his work with that "good, highly motivated group." Under the direction of Burton Richter, Schwitters, along with Martin Breidenbach, George H. Trilling and Gerson Goldhaber, began building a complex particle detector for the Stanford Positron-Electron Asymmetric Rings (SPEAR). The SPEAR collider smashed electrons into their antimatter counterparts, positrons, in collisions that released as much as four billion electron volts (GeV) of energy.

During the early 1970s, physicists were just beginning to accept the idea that all matter was composed of quarks and leptons. This theory held that the proton and neutron were composed of three quarks each; the electron and positron were classified as leptons. Yet physicists had barely an inkling that there might be several kinds, or families, of quarks and leptons.

In 1973 the SPEAR group noticed that it was producing something unusual at an energy of about 3.2 GeV. When electrons and positrons collided at that energy, the number of "events" recorded was slightly higher than the average at other energies. In June 1974 the team decided to take another look at the curious results. The initial analysis revealed nothing. But several weeks lat-

The New Standard of Excellence.

The most comprehensive, authoritative scientific dictionary now available, the **Academic Press Dictionary of Science and Technology** offers you, in one volume, the largest collection of scientific terms available in any single reference source. Containing complete, practical definitions for approximately 124,000 words (not counting abbreviations), the **Dictionary** covers key terminology in 124 fields of science and technology—including molecular biology, computer science, physics, engineering, chemistry, ecology, behavior, geophysics, veterinary science, mathematics, astronomy, biotechnology, and numerous other fields.



September 1992, 2,432 pp., \$115.00/ISBN: 0-12-200400-0

Prepublication Price: \$99.50*

*Expires on final day of month of publication.

Order from your local bookseller or directly from



ACADEMIC PRESS

HBJ Order Fulfillment Department #17915
6277 Sea Harbor Drive, Orlando, FL 32887

CALL TOLL FREE 1-800-321-5068 or FAX 1-800-336-7377

Prices subject to change without notice. ©1992 by Academic Press, Inc. All Rights Reserved. LK/MEH/WR—11092

Its Key Features Include

- Approximately 124,000 fully-defined entries, *not counting abbreviations*—more than any other scientific dictionary
- Complete coverage of 124 fields of science ranging from acoustics to zoology
- Single volume, large 8½" x 11", double-column format for easy access
- Cutting-edge definitions in such dynamic fields as artificial intelligence and biotechnology
- More medical terms than any other general scientific dictionary now available
- More coverage of subfields of medicine, including rapidly growing disciplines such as oncology, radiology, hematology, cardiology, and toxicology
- Nearly 2,000 detailed illustrations and technical photographs, including 24 pages of color plates
- Pronunciation guides for difficult or phonetically irregular terms
- Complete appendix of frequently consulted scientific data, including the periodic table, chronology of science, and standard weights and measures

Another distinctive feature of the **Dictionary** is its 124 "Windows." These boxed essays are written by leading scientists in the field, such as Stephen Jay Gould, Michael DeBakey, Rosalyn Yalow, and Linus Pauling, each of whom offers a brief introduction to his or her area of expertise. Presenting practical, concise information on each discipline, the "Windows" help make the terminology easier to understand.

The Only Source You Need for Scientific Terminology

If your library, institution, or corporation serves practicing scientists and professionals in any scientific field—engineers, computer professionals, and other technical personnel; high school, college, and graduate students; writers, researchers, and educators working with a scientific vocabulary; or general readers interested in science—then the **Academic Press Dictionary of Science and Technology** is the only scientific dictionary you need. The definitions are clear and accessible to the non-specialist, yet provide all the technical information that the specialist needs.

Set a new standard of excellence by adding the **Academic Press Dictionary of Science and Technology** to your reference collection.

It will become your reference of choice.

er the team realized that a computer glitch had skewed the results.

At home one mid-October night, Schwitters began to scrutinize the runs, one by one. He found two in which the number of events was three times higher than the average. "I thought this was great," Schwitters chuckles. "I had found the bad runs, and all we would have to do is find a reason to remove them." On October 22 he and his colleagues began a painstaking analysis of the two "bad" sets of data. Perhaps, they speculated, the inconsistencies were the result of a malfunctioning component of the detector or another computer error.

Then, while waiting for a bus on the afternoon of November 5, Schwitters "literally had one of these epiphanies that people talk about in science," he recalls. "I realized there was nothing wrong with the runs." Rather, he remembers thinking, the group may have actually discovered a new particle.

Because the evidence was preliminary, the group would need to perform another set of measurements. But the SPEAR collider had been upgraded to produce collisions at 5 GeV, and it was no easy feat to crank the machine back down to 3.2 GeV. "A few of us fought like crazy to go back and check," Schwitters says. Richter reluctantly decided the team could spend the weekend to take another look.

By Sunday, November 10, at an energy of 3.105 GeV, they were producing events at 100 times the normal rate. They had seen the signature of a new type of quark called charm. More important, the existence of charm established that the theory of quarks was correct beyond a shadow of a doubt. "Needless to say," Schwitters exclaims, "that was the most important point in my scientific career."

But the SPEAR team was not alone in its discovery. Using a particle accelerator, the Alternating Gradient Synchrotron at Brookhaven National Laboratory, Samuel Chao Chung Ting and his collaborators had also found evidence for the charm quark. On November 11 both teams announced the discovery of a new particle at 3.105 GeV. Ting had had good evidence as early as the end of October but did not have the confidence to publish his results.

During the next year, rumors circulated that the SLAC team had learned of Ting's work well before November 11. But in 1976 the Royal Swedish Academy of Sciences settled the issue of priority, for the most part, by awarding both Richter and Ting the Nobel Prize for the discovery of the charm quark. Since then, physicists have concluded that quarks come in six differ-

ent "flavors": up, down, strange, charm, bottom and top; all have been discovered except the top quark.

In 1979, excited by the prospects of teaching and living in Boston, Schwitters accepted a professorship in the physics department of Harvard University. During his tenure, he continued to apply his expertise in accelerator detectors. He teamed up with Alvin V. Tollestrup to lead the international

Schwitters feels a deep obligation to provide a productive future for particle physics.

team of 200 scientists designing and building the Collider Detector at Fermilab (CDF). The detector weighed 4,500 tons and could accommodate more than 100,000 particle collisions per second. For a decade, the CDF was the primary experimental apparatus of the laboratory, revealing insights into the size of quarks, the nature of the weak and electromagnetic forces and the properties of the bottom quark.

Throughout, one of Schwitters's goals has been the elusive top quark. He had hoped to detect it at SLAC; he had expected to see it at Fermilab. No such luck. "I feel," Schwitters says, "a little like the Captain Ahab of the top quark." Not surprisingly, he has had his hopes pinned on the SSC since he became involved in the project in 1986.

At the time, Maury Tigner of Cornell University was responsible for developing the initial design of the collider. As now conceived, the major components of the collider will be installed in a long circular tunnel, 50 meters below the grassy plains of Ellis County. The SSC is designed to accelerate two beams of protons in opposite directions and guide them into the center of one of two huge detectors. There the collision of one proton with another should release 40 trillion electron volts (TeV)—an energy scale that surpasses existing accelerators by a factor of 20.

Schwitters became the director of the laboratory in January 1989, just as Admiral James D. Watkins was being confirmed as secretary of energy. As head of the agency that provides the federal funding for U.S. accelerator programs, Watkins told senators he would not approve the construction of the SSC "if it went a penny above 5.9 billion." That year Congress appropriated the first \$225 million to start the project.

Yet in 1990, as the laboratory began the final design work, it became clear

that the SSC, as originally envisioned, could not be built for \$5.9 billion. To date, Schwitters's laboratory has spent more than \$1 billion of state and federal funds to develop the magnets and begin construction. "We had to soul-search," Schwitters explains. "Should we reduce the scientific scope, or should we ask the government for some more money? It was a very trying time."

Particle physicists objected strongly to scaling back the SSC. If the collider generated much less than 40 TeV, the project would have little chance of achieving its primary scientific goal: the discovery of the origin of mass. Theorists have calculated that a proton collider at 40 TeV will generate evidence for a particle known as the Higgs boson—or reveal a more complex mechanism—that accounts for the mass of all particles. "The whole motive," Schwitters declares, "is to do science and build a machine that works."

Congress was sympathetic to that argument in 1991, and it acceded to the fact that the collider might cost \$8.3 billion. But the legislators' largess apparently did not extend into 1992. If the SSC survives the current funding crisis, Schwitters's worries are far from over. First, his team must conduct the final tests on prototypes of the magnets before mass production can begin. Second, the laboratory is counting on foreign nations to contribute nearly \$1.5 of the \$8.3 billion, but intense negotiations, particularly with the Japanese, have failed so far to obtain a commitment.

Most important, Schwitters must create an environment for cutting-edge research while supervising a mammoth construction project. "The research initiatives must come from the youngest members of the staff, who will probably have the best new ideas," he comments. "We have to provide that opportunity in the framework of a major construction project that must have clear top-down responsibility for all of the taxpayers' money."

Through all the political wrangling, Schwitters is trying to keep his staff focused on events in Ellis County rather than on Capitol Hill. Yet lurking in the back of all of their minds is the thought that the laboratory may be shut down. "We convinced the government to go forward five years ago," Schwitters sighs. "A lot of money has been spent. People have been moved off their farms. To say, 'Sorry, we didn't really mean it'—that is exceedingly negative." Schwitters's choice of words would likely have been more flavorful were it not for his delicate position in Washington.

—Russell Ruthen



“I’D HAVE TO GO 30 YEARS WITHOUT MISSING A PUTT

“Let’s face it, on most days, I’m lucky to go three holes without missing a putt. So when I say 30 years, I haven’t gone off the deep end.

“I’m just making a point about the unbelievably high quality standards that Motorola Cellular sets for itself.

“The fact is, they’re among the highest in the world.

“And since Motorola cellular phones are made so well, it should come as no surprise that they’re the world’s best sellers, delivering the kind of customer satisfaction nobody else can match.

“Motorola. It’s an American company setting standards of quality the whole world can shoot for.

“Something that gets a big thumbs up from me.”

TO HIT THE QUALITY STANDARDS MOTOROLA SETS FOR ITS CELLULAR PHONES.”



Al D'Amico



MOTOROLA
Cellular Telephones

Mind and Brain


The biological foundations of consciousness, memory and other attributes of mind have begun to emerge; an overview of this most profound of all research efforts

by Gerald D. Fischbach

Ruth gave me a piece of her mind this morning. I am grateful, of course, but I don't know where to put it or, for that matter, what it is. I suppose that the imperatives belong in the limbic system and the geographic information in the hippocampus, but I am not sure. My problem also troubled René Descartes. Three centuries ago he described the mind as an extracorporeal entity that was expressed through the pineal gland. Descartes was wrong about the pineal, but the debate he stimulated regarding the relation between mind and brain rages on. How does the non-material mind influence the brain, and vice versa?

In addressing this issue, Descartes was at a disadvantage. He did not realize the human brain was the most complex structure in the known universe, complex enough to coordinate the fingers of a concert pianist or to create a three-dimensional landscape from light that falls on a two-dimensional retina. He did not know that the machinery of the brain is constructed and maintained jointly by genes and by experience. And he certainly did not know that the current version is the result of millions of years of evolution. It is difficult to understand the brain because, unlike a computer, it was not built with specific purposes or principles of design in mind. Natural selection, the engine of evolution, is responsible.

If Descartes had known these things, he might have wondered, along with modern neurobiologists, whether the brain is complex enough to account for the mystery of human imagination, of memory and mood. Philosophical inquiry must be supplemented by experiments that now are among the most



urgent, challenging and exciting in all of science. Our survival and probably the survival of this planet depend on a more complete understanding of the human mind. If we agree to think of the mind as a collection of mental processes rather than as a substance or spirit, it becomes easier to get on with the necessary empirical studies. In this context the adjective is less provocative than the noun.

The authors of the articles in this special issue of *Scientific American* and their colleagues have been pressing the search for the neural basis of mental phenomena. They assume that mental events can be correlated with patterns of nerve impulses in the brain. To appreciate the meaning of this assumption fully, one must consider how nerve cells, or neurons, work; how they communicate with one another; how they are organized into local or distributed networks, and how the connections between neurons change with experience. It is also important to define clearly the mental phenomena that need to be explained. Remarkable advances have been made at each level of analysis. Intriguing correlations have in fact begun to emerge between mental attributes and the patterns of nerve impulses that flare and fade in time and space, somewhere inside the brain.

The most striking features of the human brain are the large, seemingly symmetric cerebral hemispheres that sit astride the central core, which extends down to the spinal cord. The corrugated hemispheres are covered by a cell-rich, laminated cortex two millimeters in thickness. The cerebral cortex can be subdivided by morphological and functional criteria into numerous sensory receiving areas, motor-control areas and less well-defined areas in which associative events take place. Many observers assume that here, in the interface between input and output, the grand syntheses of mental life must occur.

It may not be that simple. Mind is often equated with consciousness, a subjective sense of self-awareness. A vigilant inner core that does the sensing and moving is a powerful metaphor, but there is no a priori reason to assign a particular locus to consciousness or even to assume that such global awareness exists as a physiologically unified entity. Moreover, there is more to mind than consciousness or the cere-

GERALD D. FISCHBACH is Nathan Marsh Pusey Professor of Neurobiology and chairman of the department of neurobiology at Harvard Medical School and Massachusetts General Hospital. After graduating from Colgate University in 1960, he earned his medical degree at Cornell University Medical School in 1965 and received an honorary M. A. from Harvard University in 1978. Fischbach is also a member of the National Academy of Sciences, the National Institute of Medicine and the American Academy of Arts and Sciences. He is a past-president of the Society for Neuroscience and serves on several foundation boards and university advisory panels.




HIDE-AND-SEEK (1940–42), by Pavel Tchelitchew, captures interplay between the mind and environment that influences the brain's development as well as its architecture. Hidden

forms are embedded figures, a delicate test of mental function. Roots, branches and vines suggest neuronal arborization and the ability of such structures to change.

bral cortex. Urges, moods, desires and subconscious forms of learning are mental phenomena in the broad view. We are not zombies. Affect depends on the function of neurons in the same manner as does conscious thought.

And so we return to the organ itself. The brain immediately confronts us with its great complexity. The human brain weighs only three to four pounds but contains about 100 billion neurons. Although that extraordinary number is of the same order of magnitude as the number of stars in the Milky Way, it cannot account for the complexity of the brain. The liver probably contains 100 million cells, but 1,000 livers do not add up to a rich inner life.

Part of the complexity lies in the diversity of nerve cells, which Santiago Ramón y Cajal, the father of modern brain science, described as “the mysterious butterflies of the soul, the beating of whose wings may some day—who knows?—clarify the secret of mental life.” Cajal began his monumental studies of adult and embryonic neurons about 100 years ago, when he came across Camillo Golgi’s method of staining neurons with silver salts. The great advantage of this technique, which led Cajal to his neuron doctrine, is that silver impregnates some cells in their entirety but leaves the majority untouched. Individuals thus emerged from the forest. Seeing them, Cajal realized immediately that the brain was made up of discrete units rather than a continuous net.



NEURONS, revealed by Golgi staining, carry nerve impulses. The cellular architecture of the brain was discovered by Santiago Ramón y Cajal. Janet Robbins in David H. Hubel's laboratory at Harvard Medical School made this preparation.

He described neurons as polarized cells that receive signals on highly branched extensions of their bodies, called dendrites, and send the information along unbranched extensions, called axons. The Golgi stain revealed a great variety of cell-body shapes, dendritic arbors and axon lengths. Cajal discerned a basic distinction between cells having short axons that communicate with neighbors and cells having long axons that project to other regions.

Shape is not the only source of variation among neurons. Diversity is even greater if molecular differences are considered. Whereas all cells contain the same set of genes, individual cells express, or activate, only a small subset. In the brain, selective gene expression has been found within such seemingly homogeneous populations as the amacrine cells in the retina, the Purkinje cells in the cerebellum and the motor neurons in the spinal cord. Beyond the structural and molecular differences, even more refined distinctions among neurons can be made if their inputs and projections are taken into account. Is it possible that each neuron is unique? This is certainly not the case in all but the most trivial circumstances. Yet the fact that the brain is not made up of interchangeable parts cannot be ignored.

In the face of this astounding diversity, it is a relief to learn that simplifications can be made. Several years ago Vernon B. Mountcastle, working on the somatosensory cortex, and David H. Hubel and Torsten N. Wiesel, working on the visual cortex, produced an important insight. They observed that neurons of similar function are grouped to-

gether in columns or slabs that extend through the thickness of the cortex. A typical module in the visual cortex whose component cells respond to a line of a particular orientation measures approximately one tenth of a millimeter across. The module could include more than 100,000 cells, the great majority of which participate in local circuits devoted to a particular function.

Another simplification is that all neurons conduct information in much the same way. Information travels along axons in the form of brief electrical impulses called action potentials, the beating wings of Cajal's butterflies. Action potentials, which measure about 100 millivolts in amplitude and one millisecond in duration, result from the movement of positively charged sodium ions across the surface membrane from the extracellular fluid into the cell interior, or cytoplasm.

The sodium concentration in the extracellular space is about 10 times the intracellular concentration. The resting membrane maintains a voltage gradient of -70 millivolts; the cytoplasm is negatively charged with respect to the outside. But sodium does not enter rapidly because the resting membrane does not allow these ions easy access. Physical or chemical stimuli that decrease the voltage gradient, or depolarize the membrane, increase sodium permeability. Sodium influx further depolarizes the membrane, thus increasing sodium permeability even more.


At a critical potential called the threshold, the positive feedback produces a regenerative event that forces the membrane potential to reverse in sign. That is, the inside of the cell be-

comes positive with respect to the outside. After about one millisecond the sodium permeability declines, and the membrane potential returns to -70 millivolts, its resting value. The sodium permeability mechanism remains refractory for a few milliseconds after each explosion. This limits to 200 per second or less the rate at which action potentials can be generated.

Although axons look like insulated wires, they do not conduct impulses in the same way. They are not good cables: the resistance along the axis is too high and the membrane resistance too low. The positive charge that enters the axon during the action potential is dissipated in one or two millimeters. To travel distances that may reach many centimeters, the action potential must be frequently regenerated along the way. The need to boost repeatedly the current limits the maximum speed at which an impulse travels to about 100 meters per second. That is less than one millionth of the speed at which an electrical signal moves in a copper wire. Thus, action potentials are relatively low frequency, stereotypical signals that are conducted at a snail's pace. Fleeting thoughts must depend on the relative timing of impulses conducted over many axons in parallel and on the thousands of connections made by each one.

The brain is not a syncytium, at least not a simple one. Action potentials cannot jump from one cell to another. Most often, communication between neurons is mediated by chemical transmitters that are released at specialized contacts called synapses. When an action potential arrives at the axon terminal, transmitters are released from small vesicles in which they are packaged into a cleft 20 nanometers in width that separates presynaptic and postsynaptic membranes. Calcium ions enter the nerve terminal during the peak of the action potential. Their movement provides the cue for synchronized exocytosis, the coordinated release of the neurotransmitter molecules.


Once released, transmitters bind to postsynaptic receptors, triggering a change in membrane permeability. The effect is excitatory when the movement of charge brings the membrane closer to the threshold for action-potential generation. It is inhibitory when the membrane is stabilized near its resting value. Each synapse produces only a small effect. To set the intensity (action-potential frequency) of its output, each neuron must continually integrate up to 1,000 synaptic inputs, which do not add up in a simple linear manner. Each neuron is a sophisticated computer.



The Brain: Organ of the Mind

For good reason, the human brain is sometimes hailed as the most complex object in the universe. It comprises a trillion cells, 100 billion of them neurons linked in networks that give rise to intelligence, creativity, emotion, consciousness and memory. Large anatomic subdivisions in the brain offer a rough map of its capabilities. At a very gross level, the brain is bilaterally symmetric, its left and right hemispheres connected by the corpus callosum and other axonal bridges. Its base consists of structures such as the medulla, which regulates the autonomic functions (including respiration, circulation and digestion), and the cerebellum, which coordinates movement. Within lies the limbic system (blue), a collection of structures involved in emotional behavior, long-term memory and other functions.

The highly convoluted surface of the cerebral hemispheres—the cortex (from the Latin word for bark)—is about two millimeters thick and has a total surface area of about 1.5 meters, approximately that of an office desk. The most evolutionarily ancient part of the cortex is part of the limbic system. The larger, younger neocortex is divided into frontal, temporal, parietal and occipital lobes that are separated by particularly deep sulci, or folds. Most thought and perception take place as nerve impulses, called action potentials, move across and through the cortex. Some brain regions with specialized functions have been studied in detail, such as the motor cortex (pink), the somatosensory cortex (yellow) and the visual pathway (purple). From the collective activity of all the brain regions emerges the most fascinating neurological phenomenon of all: the mind.




How Neurons Communicate

A neuron that has been excited (*turquoise*) conveys information to other neurons (*purple*) by generating impulses known as action potentials. These signals propagate like waves down the length of the cell's single axon and are converted to chemical signals at synapses, the contact points between neurons.

When a neuron is at rest, its external membrane maintains an electrical potential difference of about -70 millivolts (the inner surface is negative relative to the outer surface). At rest, the membrane is more permeable to potassium ions than to sodium ions, as indicated by the lengths of the dark arrows in the inset at the top right. When the cell is stimulated, the permeability to sodium in-

creases, leading to an inrush of positive charges (*a*). This inrush triggers an impulse—a momentary reversal (*b*) of the membrane potential. The impulse is initiated at the junction of the cell body and the axon and is conducted away from the cell body (*red arrows*).

When the impulse reaches the axon terminals of the presynaptic neuron, it induces the release of neurotransmitter molecules (*inset at bottom left*). Transmitters diffuse across a narrow cleft and bind to receptors in the postsynaptic membrane. Such binding leads to the opening of ion channels and often, in turn, to the generation of action potentials in the postsynaptic neuron. For the sake of clarity, several elements are drawn larger than scale.



STRUCTURAL VARIETY OF NEURONS (*shown as tracings from Golgi stains*) contributes to the vast capacity of the brain to store, retrieve, use and express information, as well as to experience emotion and control movement.

Many different kinds of transmitters have been identified in the brain, and this variety has enormous implications for brain function. Since the first neurotransmitter was identified in 1921, the list of candidates has grown at an increasing pace. Fifty is close to the mark. We have learned a great deal about how transmitters are synthesized, how they are released and how they activate receptors in the postsynaptic membrane.

This level of analysis is particularly relevant for psychiatric and neurological disorders that shed light on the workings of the mind [see "Major Disorders of Mind and Brain," by Elliot S. Gershon and Ronald O. Rieder, page 126]. For example, drugs that alleviate anxiety, such as Valium, augment the action of gamma-aminobutyric acid (GABA), an important inhibitory transmitter. Antidepressants, such as Prozac, enhance the action of serotonin, an indoleamine with a wide variety of functions. Cocaine facilitates the action of dopamine, whereas certain antipsychotics antagonize this catecholamine. Nicotine activates acetylcholine receptors, which are distributed throughout the cerebral cortex. Further insight into the chemical bases of thinking and behavior depends on obtaining more precise data regarding the sites of action of these potent agents and on the discovery of more selective ligands, molecules that bind to receptors.

The power of the molecule-to-mind approach can be illustrated by recent advances in the pharmacologic treatment of schizophrenia, the most common and the most devastating of all thought disorders. The classic antipsychotic drugs

include the phenothiazines (for example, Thorazine) and the butyrophenones (for example, Haldol). These agents ameliorate hallucinations, delusions, disorganized thinking and inappropriate affect—the "positive" symptoms of schizophrenia that are most evident during acute psychotic episodes. They are not as effective in treating autism and paucity of speech—"negative" symptoms that are prominent during interpsychotic intervals. Moreover, they all produce subtle, abnormal movements when administered to treat acute episodes of illness (hence the name "neuroleptics"). When administered for a long time, they often cause a devastating disorder called tardive dyskinesia. Involuntary and at times incessant writhing movements of the limbs and trunk characterize the disorder, which can persist long after the drug is discontinued.

Why would an agent that affects mental function also produce motor symptoms? The answer lies in the fact that conventional antipsychotics prevent the binding of dopamine to its receptors. To appreciate the importance of this insight, one must know that dopamine-containing nerve cell bodies, gathered deep in the midbrain in a region known as the ventral tegmentum, project their axons widely to the prefrontal cortex as well as to subcortical structures, including the basal ganglia, which are involved in many aspects of motor control. The prefrontal cortex is particularly relevant to schizophrenia because it contains circuits that are active during manipulation of symbolic information and in a type of short-term memory called working memory [see "Working


PET SCANS show the brain of a human subject performing a series of intellectual tasks related to words. The positron emission tomographic technique reveals that blood flow in the brain shifts to different locations, depending on which task is being performed. Marcus Raichle of the Washington University School of Medicine made the images.

Memory and the Mind," by Patricia S. Goldman-Rakic, page 110]. Neurons in this region may form a central processing unit of sorts.

A new drug, clozapine, affects the negative as well as the positive symptoms of schizophrenia. Most important, clozapine does not lead to tardive dyskinesia. The discovery of additional members of the dopamine receptor family may provide the explanation for the unique efficacy and selectivity of this antipsychotic.

Transmitter receptors can be grouped into two large (and growing) superfamilies based on their amino acid sequence and on presumptions about the shape that the molecules assume as part of the cell membrane in which they are embedded. A more detailed receptor classification scheme has emerged. It incorporates molecular architecture as well as the more traditional criteria of ligand binding and function. Based on the added molecular information, one receptor superfamily consists of ion channels, proteins that can form aqueous pores through which ions cross the membrane. They underlie the changes in permeability discussed above. The other superfamily, which includes the dopamine receptors, does not form channels. Instead its members interact with a neighboring membrane protein that cleaves a high-energy phosphate bond from guanosine triphosphate. This process initiates a cascade of biochemical reactions. Such G protein-mediated effects are slow in onset, and they last longer than directly gated receptor responses. It is therefore unlikely that they mediate rapid, point-to-point synaptic transmission in the brain. Rather they modulate the way ion channels respond to stimuli. They set the gain of the system much as the pedals on a piano modulate the action of the keys.

The first dopamine receptor gene was isolated four years ago. The search was based on the presumption that the receptor would resemble other receptors that were known to couple to G proteins. This powerful "homology" screening strategy led in short order to



the identification of four more dopamine receptors. One of the recent additions, imaginatively named D_4 , has attracted considerable attention. The receptor binds dopamine and clozapine with extraordinarily high affinity. Of equal importance, the D_4 gene is apparently not expressed in the basal ganglia, a finding that may explain the absence of tardive dyskinesia. Precise localization of the D_4 receptor within the prefrontal cortex may reveal the origin of hallucinations or at least a component of the neural machinery that has gone awry in schizophrenia.

The slow rate at which psychoactive drugs work presents a puzzle. Drug receptor interactions are immediate, yet symptoms of schizophrenia, depression and other disorders do not resolve for several weeks. The first consequences of drug binding cannot be the sole explanation for their efficacy. This issue leads to a more general consideration of mechanisms by which the environment might change the brain.

Investigation of dopamine synapses has also provided information about the curse of drug addiction. Cocaine, which binds to and inhibits a protein that transports dopamine away from its site of action, is one of the most powerful reinforcing drugs known. Recent studies point to a neural pathway that may be a target of all addictive substances—amphetamines, nicotine, alcohol and opiates. Within this pathway, the nucleus accumbens, a small subdivision of the basal ganglia, appears to be particularly important. Further studies of neurons in this region will certainly sharpen understanding of drug-seeking behavior. They may reveal mechanisms of motivation in general.


The structural, functional and molecular variety that has been described so far would seem to provide a sufficiently complete basis for mental function. Yet another dimension must be considered: plasticity, the tendency of synapses and

neuronal circuits to change as a result of activity. Plasticity weaves the tapestry on which the continuity of mental life depends. Action potentials not only encode information, their metabolic aftereffects alter the circuits over which they are transmitted.

Synaptic plasticity is the basis for the informative connectionist neural models that Geoffrey E. Hinton describes [see "How Neural Networks Learn from Experience," page 144]. More generally, plasticity multiplies the complexity provided by any fixed cast of molecular characters or cellular functions. Hence, it provides an even richer substrate for mental phenomena.

From the brief tour of synaptic biology presented above, you can imagine many ways that synaptic efficacy might be altered. For example, transmitter release can be enhanced by a small increase in the amount of calcium that enters a nerve terminal with each action potential. The probability of postsynaptic receptor activation can be changed, and on a longer time scale, variations in activity can alter the number of functional receptors. Increases or decreases in the number of receptors, which take time to occur, may account for the delayed effect of psychotherapeutic agents. Beyond changes in the function of synapses, activity may alter the number or location of synapses themselves. Axons sprout new endings when their neighbors become silent, and the terminal branches of dendritic arbors are constantly remodeled [see "Aging Brain, Aging Mind," by Dennis J. Selkoe, page 134].

In their discussion of plasticity and learning [see "The Biological Basis of Learning and Individuality," page 78], Eric R. Kandel and Robert D. Hawkins review evidence that short-term synaptic changes associated with simple forms of learning are accompanied by molecular modification of proteins. One



such modification is phosphorylation, the addition or attachment of a phosphate group. Phosphorylation has a profound effect on the function of proteins. It is commonly stimulated by transmitters and drugs that act via G protein-coupled receptors. But proteins are degraded on a time scale that ranges from minutes to days. Maintenance of memories that may last a lifetime requires more stable alterations, such as those associated with persistent changes in gene expression. A recently discovered family of genes called immediate early genes (IEGs), which are activated rapidly by brief bursts of action potentials, may provide a crucial link. As expected of master switches that initiate long-term changes in the brain, IEGs encode transcription factors, proteins that regulate the expression of other genes.

Some evidence has been obtained that impulse activity increases the expression of genes that encode trophic factors, proteins that promote the survival of neurons. The adage "use it or lose it" may soon have a specific biochemical correlate. The actions of each transcription factor and their relevance remain to be determined, however.

Another focus of inquiry into the basis of memory is the phenomenon of long-term potentiation (LTP), a persistent increase in synaptic efficacy that follows brief periods of stimulation. Attention has focused on synapses in the hippocampus because clinical and experimental data have implicated this region of the cortex in forms of memory that require conscious deliberation. At certain synapses in the hippocampus, LTP may last for weeks. At the same junctions, LTP meets the "Hebbian" criterion for learning in that it requires coincident presynaptic and postsynaptic activity. LTP does not occur if the postsynaptic neuron is rendered inactive during the priming, presynaptic stimulation. Donald O. Hebb suggested this relation in his 1949

book *The Organization of Behavior* as a basis for the formation of new neural ensembles during learning. It has been repeated often enough to have achieved the force of law.

Synaptic transmission in the hippocampus is mediated by glutamate, the most common excitatory transmitter in the brain, and LTP of the Hebbian type is blocked by aminophosphonovaleric acid (APV), a selective antagonist of one type of glutamate receptor. APV also diminishes the ability of rats to learn tasks that require spatial cues. This is probably not a coincidence, but it remains to be shown that these observations are causally related. The gene that encodes the APV-sensitive glutamate receptor has been cloned in recent months. We can therefore expect tests in transgenic mice bearing mutated receptors to be conducted in the near future. The work will not be straightforward. The plasticity of the brain and the likelihood that natural selection has provided alternative routes to such an important end may complicate matters.

Although the forces leading to plastic changes in the mature brain are ubiquitous and unrelenting, it is important to emphasize the precision and overall stability of the wiring diagram. We could not sense the environment or move in a coordinated manner, let alone think, if it were otherwise. All studies of higher brain function must take into account the precise way in which neurons are connected to one another.


Pathways in the brain have been traced by means of a variety of molecules that are transported along axons. Such reporter molecules can be visualized once the tissue is properly prepared. Connections have also been traced by fine-tipped microelectrodes positioned close enough to a nerve cell body or an axon to detect the small currents generated as an action potential passes by. Each technique

has revealed ordered, topographic maps in the cerebral cortex. The body surface is represented in the postcentral gyrus of the cerebral cortex even though the cortical neurons are three synapses away from sensory receptors in the skin. Likewise, a point-to-point map of the visual world is evident in the primary visual cortex at the occipital pole at the back of the brain. Order is evident at each of the early relays on route to the cortex, and topographic order has also been found in projections from the primary cortices to higher centers.


To appreciate just how precise the wiring diagram can be, we need only consider a fundamental discovery made about 30 years ago by Hubel and Wiesel. They determined that neurons in the primary visual cortex (V1) respond to line segments or edges of a particular orientation rather than to the small spots of light that activate the input neurons in the retina and lateral geniculate nucleus of the thalamus. The response implies that neurons in V1 are connected, via the lateral geniculate nucleus, to retinal ganglion cells that lie along a line of the preferred orientation.

We know the anatomy of the major sensory and motor systems in some detail. In contrast, the pattern of connections within the intervening association cortices and the large subcortical nuclei of the cerebral hemispheres is not clearly defined. Goldman-Rakic's experiments are designed to decipher the wiring diagram of the monkey's prefrontal cortex in order to provide a more complete anatomy of memory. Our lack of information about similar connections in the human brain is glaring. Unlike the molecular building blocks and the functions of individual neurons, it cannot be assumed that the intricacies of cortical connectivity will be conserved in different species. The intricacy of this network, after all, is what distinguishes *Homo sapiens* from all other forms of life. An effort akin to the genome project may be called for.

How does the specificity of synaptic connections come about during development? Carla J. Shatz reviews mechanisms by which axons are guided to their appropriate targets in the visual and other systems [see "The Developing Brain," page 60]. The initial stages of axon outgrowth and pathway selection are thought to occur independently of activity. The genetically determined part of the program is evident in the remarkably complete wiring diagram that forms during embryonic life. But once the advancing tips of the axons arrive in the appropriate region, the choices of particular targets are influenced by nerve impulses originating within



INTENT of a monkey to move its arm is revealed by electrical activity of neurons in the motor cortex. Microelectrodes recorded the impulses. Each line represents the rate of firing of individual neurons. Computer diagram at the left shows firing associated with the full range of arm movement. Diagram at



the right shows the firing of neurons controlling movement in one direction only (*long yellow line*). The direction of the population vector (*orange line*) is close to that of the actual movement. Apostolos P. Georgopoulos of Johns Hopkins University and his colleagues made the measurements.

the brain or stimulated by events in the world itself. Synapse formation during a critical period of development may depend on a type of competition between axons in which those that are activated appropriately are favored.

Steroid hormones also influence the formation of synapses during early development at least in certain regions of the brain [see "Sex Differences in the Brain," by Doreen Kimura, page 118]. Anatomic, physiological and behavioral data indicate that the brains of males and females are not identical.

The pattern of information flow in the brain during the performance of mental tasks cannot easily be determined by anatomic studies of the circuit diagram or by studies of plasticity. Neural correlates of higher mental functions are being sought directly in awake primates trained to perform tasks that require judgment, planning or memory, or all three capacities. This demanding approach requires sophisticated instrumentation, sophisticated experimental design and months of training until the monkey thinks the same thoughts as the investigator. All-night sessions spent listening to amplified action potentials generated by one or a few neurons followed by days of data analysis are the rule. Progress is slow, but important generalizations have emerged.

One of the most important principles is that sensory systems are arranged in a hierarchical manner. That is, neurons respond to increasingly abstract aspects of complex stimuli as the distance—measured in numbers of synapses from the source—grows. The fact that neurons in V1 respond to lines rather than spots makes the case. Another impor-

tant principle, discussed by Semir Zeki, is that information does not travel along a single pathway. Rather, different features of a single percept are processed in parallel pathways [see "The Visual Image in Mind and Brain," page 68]. A tennis player who wanders to the net from time to time will be alarmed to learn that the movement, color and shape of a tennis ball are processed in different cortical visual centers. Separation of these information streams begins in the retina; they remain segregated in the lateral geniculate nucleus and the primary visual cortex en route to the higher visual centers.

An analogous situation has been found in the auditory system. Mark Konishi and his colleagues at the California Institute of Technology have shown that the localization of sound sources by the barn owl depends on interaural phase and amplitude differences. Phase differences indicate location along the azimuth, whereas amplitude differences signal elevation. Phase and amplitude signals are processed in different pathways through three synaptic relays in the brain. It seems likely that this type of parallel processing characterizes other sensory systems, association cortices and motor pathways as well.

Where is the information reassembled? When does the subject become aware of the approaching ball? The receptive fields of neurons in higher centers are larger than those found in earlier relay stations, so they monitor a larger fraction of the external world. Zeki describes a model that depends on feedback connections from cells with large receptive fields to the cells in the primary visual cortex that have high spatial resolu-

tion. Such feedback circuits might coordinate the activity of cells in the primary cortex that have high spatial resolution and cells that respond to more abstract features of the stimulus no matter where it is located. Francis Crick and Christof Koch address the role in visual awareness of a 40-cycle-per-second oscillation in firing rate that is observed throughout the cortex [see "The Problem of Consciousness," page 152]. The oscillations, discovered by Wolf J. Singer and his colleagues at the Max Planck Institute for Brain Research in Frankfurt, may synchronize the firing of neurons that respond to different components of a perceptual scene and hence may be a direct neural correlate of awareness.

Konishi has identified the first neurons in the owl's brain that respond to a combination of interaural phase and amplitude differences but not to either parameter presented alone. These neurons, located deep in the animal's brain in a region called the inferior colliculus, activate a motor program that results in the owl's turning toward the sound source.

In the monkey's visual system, "face cells" located in the inferior temporal sulcus represent perhaps the highest level of abstraction yet identified. These neurons respond to faces but not to other visual stimuli. Similar cells may be present in our own brains. Lesions in the corresponding area of the temporal lobe result in prosopagnosia, a remarkably selective deficit in which the ability to recognize faces is lost. In the zebra finch's auditory system (birds again), a high level of abstraction is evident in neurons found in each male's brain that respond to the complex song of his father but not to pure tones

or to the songs of other males of the same species.

How many neurons must change their firing rate to signal a coherent percept or gestalt? The most extreme view holds that one cell may do the job. Is there one face cell per face? Such a supposition seems unlikely on first principles: we lose thousands of neurons every day, so overcommitment to one would be unwise. A more compelling argument comes from recent experiments that have shown face cells to be broadly tuned, responding to faces with similar features rather than to one face alone. The number of neurons that must be activated before recognition emerges is not known, but the data are consistent with a sparse coding rather than global or diffuse activation.

Face cells have their counterparts on the motor side. "Command" neurons have been identified in certain invertebrates that trigger all-or-none, fixed-action patterns, such as stereotypical escape behaviors. Apostolos P. Georgopoulos of Johns Hopkins University has found command neurons of a kind in the monkey's motor cortex (precentral gyrus) that encode the direction of forelimb movement. The firing of these neurons is not associated with the contraction of a particular muscle or with the force of the coordinated movement. Like face cells in the temporal lobe, individual motor cortex neurons are broadly tuned.

The vector obtained by summing the firing frequencies of many neurons is better correlated with the direction of movement than is the activity of any individual cell [see *illustration on opposite page*]. The vector becomes evident several milliseconds *before* the appropriate muscles contract and the arm actually moves. It must be a sign of motor planning. The vector is usually derived from less than 100 neurons, so sparse coding may be the rule in the motor cortex as it is in the temporal sulcus.

An important next step at this level of analysis is to produce mental phenomena by focal electrical stimulation. A beginning has been made by William Newsome and his colleagues at Stanford University. They trained monkeys to decide on the direction of movement of dots displayed in random positions on a screen. When the number of dots that showed net movement was set near the threshold for a consistent judgment about the population as a whole, focal stimulation of the V5 region in the cortex influenced the monkey's perceptual judgments.

Strokes and other unfortunate "experiments of nature" have also provided important insights regarding neural

correlates of mental phenomena. Antonio R. and Hanna Damasio continue a long tradition of research in their studies of language disorders among neurological patients [see "Brain and Language," page 88]. This work requires careful examination with a battery of tests designed to elicit the most subtle deficits. Here is an example of the pressing need to define the mental phenomena that need to be explained. The Damascios propose the view that language can be considered a three-part system: word formation, concept representation and mediation between the two. If, as they suggest, language has evolved as a tool to compress concepts and communicate them in an efficient manner, a clear view of its functional anatomy brings us to the crux of the mind matter.

The very real experience of phantom limbs cautions against quick acceptance of sparse coding or even of localization as a universal mechanism. Amputated limbs, experiments of nature of another sort, may be experienced as an integral part of the body or "self" [see "Phantom Limbs," by Ronald Melzack; *SCIENTIFIC AMERICAN*, April]. A deep and burning pain is a distressing component of the syndrome. It is impossible to find a local area in which such sensations are experienced. Attempts have been made to abolish phantom pain by cutting peripheral nerves, by destroying ascending pathways and by removing sensory regions of the brain. All attempts have failed to eliminate the perception of pain. It may be that the emotional response we call pain requires activation of neurons in widely dispersed regions of the brain.

The future of cognitive neuroscience depends on our ability to study the living human brain. Positron emission tomography (PET) and functional magnetic resonance imaging (MRI) hold great promise in this regard. These noninvasive imaging techniques depend on tight coupling between neuronal activity, energy consumption and regional blood flow. These relations were pointed out by Sir Charles Scott Sherrington in 1890 and later placed on a quantitative basis by Seymour S. Kety and Louis Sokoloff of the National Institute of Mental Health. The brain is never completely at rest. Furthermore, the increases in regional blood flow that MRI and PET detect are not large (they are on the order of 20 to 50 percent). So PET and MRI measurements depend on sophisticated subtraction algorithms that allow one to distinguish the pattern of blood flow during the mental task from the resting, or control, pattern. Assignment of the changes in blood flow to specific structures depends on accurate-

ly superimposing the computed images on precise anatomic maps.

At present, neither technique provides the spatial resolution to visualize single cortical columns. Moreover, the slow temporal resolution of both imaging techniques demands that mental tasks be repeated over and over again during the recording session. Technical advances, especially related to rapid MRI scanning, are sure to follow. Even with the current limitations, the advantages of working with humans, who can think on command, are overwhelming.

In sum, we can expect advances at an increasing rate on all levels of investigation relevant to the mind. We will soon know exactly how many transmitters and transmitter receptors there are in the brain and where each one is concentrated. We will also have a more complete picture of neurotransmitter actions, including multiple interactions of jointly released modulators. And we will learn much more about molecules that affect neuronal differentiation and degeneration. Molecules of the mind are not unique. Many of the neurotransmitters are common amino acids found throughout the body. Likewise, no new principles or molecules specific to the brain have emerged in studies of hormone regulation or of trophic factors that influence the survival and differentiation of neurons. The great challenge, then, is to determine how these molecules modulate the functional wiring diagram of the brain and how this functional nerve net gives rise to mental phenomena.

Ultimately, it will be essential to specify what exactly it means to say that mental events are correlated with electrical signals. Certainly, there is a need for theory at this level of analysis, and as emphasized by Crick and Koch, this effort has become one of the most exciting aspects of cognitive neuroscience.

Is the mind an emergent property of the brain's electrical and metabolic activity? An emergent property is one that cannot be accounted for solely by considering the component parts one at a time. For example, the heart beats because its pacemaker depends on the influx and efflux of certain ions. But the automaticity cannot be understood without considering the magnitude and kinetics of all the fluxes together. Once that is accomplished, what is left to explain in physiological terms? In an analogous manner, biological explanations of mental events may become evident once the component neural functions are more clearly defined. We will then have a more appropriate vocabulary for describing the emergent mind.



A DEPENDABLE, RELIABLE SOURCE of monthly income. It can be a source of comforting predictability. Like watching a sunset or taking a walk through a familiar stretch of woods. And we, at John Nuveen & Co. Incorporated, have always felt that it's the predictable things in life that are the most reassuring.  For as we see it, an investment de-

signed to provide you with a dependable source of income is something that can give you financial freedom. Not to mention the opportunity to live life on your own terms.

 And we're confident that your financial adviser will tell you that our Unit Trusts are designed to provide you with just that: a remarkably steady level of income, free from federal tax*  Because we've earned a reputation for working diligently to achieve that result, as well as for our ability

to provide investors with a variety of investment choices. Like the opportunity to receive your tax-free income in a regular monthly check.  And while investing

*Sometimes the
PREDICTABLE things
in life are the
most UNEXPECTEDLY
REASSURING.*

in these times of economic uncertainty can hardly be considered a walk in the park, it's nice to know that a Nuveen Tax-Free Unit Trust is something you can count on to be reassuringly

predictable.  So if you'd like to begin your journey on the road to attractive tax-free income, ask your broker, banker or financial adviser about Nuveen Tax-Free Unit Trusts. And rest assured, the return on your Nuveen investment will remain the same until the bonds in the portfolio mature, are called or are sold.† But please request a prospectus with more complete information (including charges and expenses), and read it carefully before you invest or send money. Or simply call us at...

1-800-262-3423.

NUVEEN

Quality Tax-Free Investments Since 1898.

*Income may be subject to state and local taxes, as well as to the alternative minimum tax. Save additional taxes with Nuveen state trusts which are also exempt from state and local income taxes. Capital gains, if any, will be subject to capital gains taxes. †Unit Trusts have a fluctuating redemption value.



The Developing Brain

During fetal development, the foundations of the mind are laid as billions of neurons form appropriate connections and patterns. Neural activity and stimulation are crucial in completing this process

by Carla J. Shatz

An adult human brain has more than 100 billion neurons. They are specifically and intricately connected with one another in ways that make possible memory, vision, learning, thought, consciousness and other properties of the mind. One of the most remarkable features of the adult nervous system is the precision of this wiring. No aspect of the complicated structure, it would appear, has been left to chance. The achievement of such complexity is even more astounding when one considers that during the first few weeks after fertilization many of the sense organs are not even connected to the embryonic processing centers of the brain. During fetal development, neurons must be generated in the right quantity and location. The axons that propagate from them must select the correct pathway to their target and finally make the right connection.

How do such precise neural links form? One idea holds that the brain wires itself as the fetus develops, in a manner analogous to the way a computer is manufactured: that is, the chips and components are assembled and connected according to a preset circuit diagram. According to this analogy, a flip of a biological switch at some point in prenatal life turns on the computer. This notion would imply that the brain's entire structure is recorded in a set of biological blueprints—presumably DNA—and that the organ begins to work only after the wiring is essentially complete.

Research during the past decade shows that the biology of brain development follows very different rules. The neural connections elaborate themselves from an immature pattern of wiring that only grossly approximates the adult pattern. Although humans are born with almost all the neurons they will ever have, the mass of the brain at birth is only about one fourth that of the adult brain. The brain becomes bigger because neurons grow in size, and the number of axons and dendrites as well as the extent of their connections increases.

Workers who have studied the development of the brain have found that to achieve the precision of the adult pattern, neural function is necessary: the brain must be stimulated in some fashion. Indeed, several observations during the past few decades have shown that babies who spent most of their first year of life lying in their cribs developed abnormally slowly. Some of these infants could not sit up at 21 months of age, and fewer than 15 percent could walk by about the age of three. Children must be stimulated—through touch,



speech and images—to develop fully. Based in part on such observations, some people favor enriched environments for young children, in the hopes of enhancing development. Yet current studies provide no clear evidence that such extra stimulation is helpful.

Much research remains to be done before anyone can conclusively determine the types of sensory input that encourage the formation of particular neural connections in newborns. As a first step toward understanding the process, neurobiologists have focused on


the development of the visual system in other animals, especially during the neonatal stages. It is easy under the conditions that prevail at that stage to control visual experience and observe behavioral response to small changes. Furthermore, the mammalian eye differs little from species to species. Another physiological fact makes the visual system a productive object of study: its neurons are essentially the same as neurons in other parts of the brain. For these reasons, the results of such studies are very likely to be applicable to the human nervous system as well.

But perhaps the most important advantage is that in the visual system, investigators can accurately correlate function with structure and identify the pathway from external stimulus to physiological response. The response begins when the rods and cones of the retina transform light into neural signals. These cells send the signals to the retinal interneurons, which relay them to the output neurons of the retina, called the retinal ganglion cells. The axons of the retinal ganglion cells (which make up the optic nerve) connect to a relay structure within the brain known as the lateral geniculate nucleus. The cells of the lateral geniculate nucleus then send the visual information to specific neurons located in what is called layer 4 of the (six-layer) primary visual cortex. This cortical region occupies the occipital lobe in each cerebral hemisphere [see illustration on next page].

Within the lateral geniculate nucleus, retinal ganglion cell axons from each eye are strictly segregated: the axons of one eye alternate with those from the other and thus form a se-

SEVEN-WEEK-OLD HUMAN FETUS is about an inch long. Eyes and limbs are visible, and the emerging brain is apparent. Stimulation is needed to complete development, a process that for many neural systems continues into neonatal life.

CARLA J. SHATZ is professor of neurobiology at the University of California, Berkeley, a position she took after many years at Stanford University. She graduated from Radcliffe College and received a master's degree in physiology from University College, London, and a Ph.D. in neurobiology from Harvard Medical School. Her studies of the development of connections in the mammalian visual system have gained her many honors, including, most recently, her election to the American Academy of Arts and Sciences.



VISUAL PATHWAY in the adult demonstrates the segregation of axons. The axons corresponding to the right eye are in red, and those corresponding to the left eye are in blue. Neighboring retinal ganglion cells in each eye send their axons to neighboring neurons in the lateral geniculate nucleus. Similarly, the neurons of the geniculate nucleus map their axons onto the visual cortex. The system forms a topographically orderly pattern that in part accounts for such characteristics as binocular vision.

ries of eye-specific layers. The axons from the lateral geniculate nucleus in turn terminate in restricted patches within cortical layer 4. The patches corresponding to each eye interdigitate with one another to form structures termed ocular dominance columns.

To establish such a network during development, axons must grow long distances, because the target structures form in different regions. The retinal ganglion cells are generated within the eye. The lateral geniculate neurons take shape in an embryonic structure known

as the diencephalon, which will form the thalamus and hypothalamus. The layer 4 cells are created in another protoorgan called the telencephalon, which later develops into the cerebral cortex. From the beginning of fetal development, these three structures are many cell-body diameters distant from one another. Yet after identifying one or the

other of these targets, the axons reach it and array themselves in the correct topographic fashion—that is, cells located near one another in one structure map their axons to the correct neighboring cells within the target.

This developmental process can be compared with the problem of stringing telephone lines between particu-

NEURAL DEVELOPMENT of the visual system is revealed here by a radioactive tracer injected into the vitreous humor of the left eye. The images correspond to a top view of the visual system of a cat. Only areas receiving input from the injected eye become labeled (*white areas*). In the lateral geniculate nucleus (*top left*), most of the left eye's input ends in layers in the right nucleus, although the wiring of the visual pathway places some label in the left nucleus. Similar segregation is seen in the ocular dominance columns in layer 4 of the visual cortex (*bottom left*); the gaps represent regions corresponding to axons from the uninjected eye. The adult patterns are in distinct contrast to their immature forms, shown to the right. The immature axons have yet to segregate: the label is uniformly distributed.



lar homes located within specific cities. For instance, to string wires between Boston and New York, one must bypass several cities, including Providence, Hartford, New Haven and Stamford. Once in New York, the lines must be directed to the correct borough (target) and then to the correct street address (topographic location).

Corey Goodman of the University of California at Berkeley and Thomas Jessel of Columbia University have demonstrated that in most instances, axons immediately recognize and grow along the correct pathway and select the correct target in a highly precise manner. A kind of "molecular sensing" is thought to guide growing axons. The axons have specialized tips, called growth cones, that can recognize the proper pathways. They do so by sensing a variety of specific molecules laid out on the surface of, or even released from, cells located along the pathway. The target itself may also release the necessary molecular cues. Removing these cues (by genetic or surgical manipulation) can cause the axons to grow aimlessly. But once axons have arrived at their targets, they still need to select the correct address. Unlike pathway and target selection, address selection is not direct. In fact, it involves the correction of many initial errors.

The first hint that address selection is not precise came from experiments using radioactive tracers. Injections of these tracers at successively later times in fetal development outline the course and pattern of axonal projections. Such studies have also

shown that structures emerge at different times in development, which can further complicate address selection.

For instance, Pasko Rakic of Yale University has shown that in the visual pathway in monkeys, the connections between the retina and the lateral geniculate nucleus appear first, followed by those between the lateral geniculate nucleus and layer 4 of the visual cortex. Other studies found that in cats and primates (including humans), the lateral geniculate nucleus layers develop during the prenatal period, before the rods and cones of the retina have formed (and thus before vision is even possible). When Simon LeVay, Michael P. Stryker and I were postdoctoral fellows at Harvard Medical School, we found that at birth, layer 4 columns in cats do not even exist in the visual cortex [see *illustrations below*]. I subsequently determined that even earlier, in fetal life, the cat has no layers in the lateral geniculate nucleus. These important visual structures emerge only gradually and at separate stages.

The functional properties of neurons, like their structural architecture, do not attain their specificity until later in life. Microelectrode recordings from the visual cortex of newborn cats and monkeys reveal that the majority of layer 4 neurons respond equally well to visual stimulation of either eye. In the adult, each neuron in layer 4 responds primarily if not exclusively to stimulation of one eye only. This finding implies that during the process of address selection, the axons must correct their early "mistakes" by removing


the inputs from the "inappropriate" eye.

In 1983 my colleague Peter A. Kirkwood and I found further evidence that axons must fine-tune their connections. It came from our work on the brains of six-week-old cat fetuses (the gestation period of the cat is about nine weeks). We removed a significant portion of the visual pathway—from the ganglion cells in both eyes to the lateral geniculate nucleus—and placed it in vitro in a special life-support chamber. (Inserting microelectrodes in a fetus is extremely difficult.) The device kept the cells alive for about 24 hours. Next we applied electrical pulses to the two optic nerves to stimulate the ganglion cell axons and make them fire action potentials, or nerve signals. We found that neurons in the lateral geniculate nucleus responded to the ganglion cells and, indeed, received inputs from both eyes. In the adult the layers respond only to stimulation of the appropriate eye.


The eventual emergence of discretely functioning neural domains (such as the layers and ocular dominance columns) indicates that axons do manage to correct their mistakes during address selection. The selection process itself depends on the branching pattern of individual axons. In 1986 David W. Sretavan, then a doctoral student in my laboratory, was able to examine the process in some detail. Experimenting with fetal cats, he selectively labeled single retinal ganglion cell axons in their entirety—from the cell body in the retina to their tips within the lateral geniculate nucleus—at successively later stages.

He found that at the earliest times in development, when ganglion cell axons have just arrived within the lateral geniculate nucleus (after about five weeks of gestation), the axons assume a very simple sticklike shape and are tipped with a growth cone. A few days later the axons arising from both eyes acquire a "hairy" appearance: they have short side branches along their entire length.

The presence of side branches at this age implies that the inputs from both eyes mix with one another. In other words, the neural regions have yet to take on the adult structure, in which each eye has its own specific regions. As development continues, the axons sprout elaborate terminal branches and lose their side branches. Soon individual axons from each eye have highly branched terminals that are restricted to the appropriate layer. Axons from one eye that traverse territory belonging to those from the other eye are smooth and unbranched [see *illustration on next page*].



The sequence of developmental changes in the branching patterns shows that the adult pattern of connections emerges as axons remodel by the selective withdrawal and growth of different branches. Axons apparently grow to many different addresses within their



AXONAL REMODELING in the lateral geniculate nucleus occurs largely before birth. At the earliest times in development (1), the axons from the left eye and right eye are simple and tipped with growth cones. The shaded region represents the intermixing of inputs from both eyes. After further development (2), the axons grow many side branches. The axons soon begin to lose some side branches and start to extend elaborate terminal branches (3). Eventually these branches occupy the appropriate territory to form eye-specific layers (4).

target structures and then somehow eliminate addressing errors.

One possible explanation for axonal remodeling is that specific molecular cues are arrayed on the surface of the target cells. Although this idea might seem conceptually attractive, it has very little experimental support. An alternative explanation appears to be stronger. It holds that all target neurons are fair game. Then, some kind of competition between inputs would lead to formation of specific functional areas.

An important clue concerning the nature of the competitive interactions between axons for target neurons has come from the experiments of David H. Hubel of Harvard Medical School and Torsten N. Wiesel of the Rockefeller University. In the 1970s, when both workers were at Harvard, they studied the formation of childhood cataracts. Clinical observations indicated that if the condition is not treated promptly, it can lead to permanent blindness in the obstructed eye. To emulate the effect, Hubel and Wiesel closed the eyelids of newborn cats. They discovered that even a week of sightlessness can alter the formation of ocular dominance columns. The axons from the lateral geniculate nucleus representing the closed eye occupy smaller than normal patches within layer 4 of the cortex. The axons of the open eye occupy larger than normal patches.

The workers also showed that the effects are restricted to a critical period. Cataracts, when they occur in adulthood and are subsequently corrected by surgery, do not cause lasting blindness. Apparently the critical period has ended long ago, and so the brain's wiring cannot be affected.

These observations suggest that the ocular dominance columns form as a consequence of use. The axons of the lateral geniculate nucleus from each eye somehow compete for common territory in layer 4. When use is equal, the columns in the two eyes are identical; unequal use leads to unequal allotment of territory claimed in layer 4.

How is use translated into these lasting anatomic consequences? In the visual system, use consists of the action potentials generated each time a visual stimulus is converted into a neural signal and is carried by the ganglion cell axons into the brain. Perhaps the effects of eye closure on the development of ocular dominance columns occur because there are fewer action potentials coming from the closed eye. If that is the case, blockage of all action potentials during the critical period of postnatal life should prevent

axons from both eyes from fashioning the correct patterns and lead to abnormal development in the visual cortex. Stryker and William Harris, then a post-doctoral fellow at Harvard, obtained this result when they used the drug tetrodotoxin to block retinal ganglion cell action potentials. They found that the ocular dominance columns in layer 4 failed to appear (the layers in the lateral geniculate nucleus were unaffected because they had already formed in utero).

Nevertheless, action potentials by themselves are not sufficient to create the segregated patterns in the cortex. Neural activity cannot be random. Instead it must be defined, both temporally and spatially, and must occur in the presence of special kinds of synapses. Stryker and his associate Sheri Strickland, who are both at the University of California at San Francisco, have shown that simultaneous, artificial stimulation of all the axons in the optic nerves can prevent the segregation of axons from the lateral geniculate nucleus into ocular dominance columns within layer 4. Although this result resembles that achieved with tetrodotoxin, an important difference exists. Here ganglion cell action potentials are present—but all at the same time. Segregation to form the columns in the visual cortex, on the other hand, proceeds when the two nerves are stimulated asynchronously.

In a sense, then, cells that fire together wire together. The timing of action-potential activity is critical in determining which synaptic connections are strengthened and retained and which are weakened and eliminated. Under normal circumstances, vision itself acts to correlate the activity of neighboring retinal ganglion cells, because the cells receive inputs from the same parts of the visual world.


What is the synaptic mechanism that strengthens or weakens the connections? As long ago as 1949, Donald O. Hebb of McGill University proposed the existence of special synapses that could execute the task. The signal strength in such synapses would increase whenever activities in a presynaptic cell (the cell supplying the synaptic input) and in a postsynaptic cell (the cell receiving the input) coincide. Clear evidence showing that such "Hebb synapses" exist comes from studies of the phenomenon of long-term potentiation in the hippocampus. Researchers found that the pairing of presynaptic and postsynaptic activity in the hippocampus can cause incremental increases in the strength of synaptic transmission between the paired cells. The strengthened state can last from hours to days.

Such synapses are now thought to be

Development and Neural Function

One of the characteristics of the developing visual system is segregation of inputs: each eye adopts its own territory in the visual cortex. The process, however, can be completed only if the neurons are stimulated. In experiments with cat eyes, for example, the axons of the left eye and of the right eye overlap in layer 4 of the visual cortex at birth (a). Visual stimuli will cause the axons to separate and

form ocular dominance columns in the cortex (b). Such normal development can be blocked with injections of tetrodotoxin; as a result, the axons never segregate, and the ocular dominance columns fail to emerge (c). Another way to perturb development is to keep one eye closed, depriving it of stimulation. The axons of the open eye then take over more than their fair share of territory in the cortex (d).



essential in memory and learning [see "The Biological Basis of Learning and Individuality," by Eric R. Kandel and Robert D. Hawkins, page 78]. Studies by Wolf Singer and his colleagues at the Max Planck Institute for Brain Research in Frankfurt and by Yves Fregnac and his colleagues at the University of Paris also suggest that Hebb synapses are present in the visual cortex during the critical period, although their properties are not well understood.

Just how coincident activity causes long-lasting changes in transmission is not known. There is general agreement among researchers that the postsynaptic cell must somehow detect the coincidence in the incoming presynaptic activity and in turn send a signal back to all concurrently active presynaptic inputs. But this cannot be the whole story. During the formation of the ocular dominance columns, inputs that are not active at the same time are weakened and eliminated.

Consequently, one must also propose the existence of a mechanism for activity-dependent synaptic weakening. This weakening—a kind of long-term depression—would occur when presynaptic action potentials do not accompany postsynaptic activity. Synapses that have

this special property (opposite to that of Hebb synapses) have been found in the hippocampus and cerebellum. The results of the Stryker and Strickland experiments suggest that such synapses are very likely to exist in the visual cortex as well.

A strongly similar process of axonal remodeling operates as motor neurons in the spinal cord connect with their target muscles. In the adult, each muscle fiber receives input from only one motor neuron. But after motor neurons make the first contacts with the muscle fibers, each muscle fiber receives inputs from many motor neurons. Then, just as in the visual system, some inputs are eliminated, giving rise to the adult pattern of connectivity. Studies have shown that the process of elimination requires specific temporal patterns of action-potential activity generated by the motor neurons.


The requirement for specific spatial and temporal patterns of neuronal activity might be likened to a process whereby telephone calls are placed from addresses in one city (the lateral geniculate nucleus in the visual system) to those in the next city (the visual cortex) to verify that connections have been made at the correct locations. When two near

neighbors in the lateral geniculate nucleus simultaneously call neighboring addresses in the cortex, the telephones in both those homes will ring. The concurrent ringing verifies that relations between neighbors have been preserved during the wiring process.

If, however, one of the neighbors in the lateral geniculate nucleus mistakenly makes connections with very distant parts of layer 4 or with parts that receive input from the other eye, the called telephone will rarely if ever ring simultaneously with those of its neighbors. This dissonance would lead to the weakening and ultimate removal of that connection.

The research cited thus far has explored the remodeling of connections after the animal can move or see. But what about earlier in development? Can mechanisms of axonal remodeling operate even before the brain can respond to stimulation from the external world? My colleagues and I thought the formation of layers in the lateral geniculate nucleus in the cat might be a good place to address this question. After all, during the relevant developmental period, rods and cones have not yet emerged. Can the layers

RETINAL ACTIVITY, recorded frame by frame every 0.5 second by a hexagonal array of microelectrodes (*black spots*), is locally synchronized. Each diagram represents the pattern and intensity of action-potential firing (*red*) of individual ganglion cells. The wave of retinal activity sweeps across from the lower left to the top right of the retina.



develop their specific territories for each eye even though vision cannot yet generate action-potential activity?

We reasoned that if activity is necessary at these early times, it must somehow be generated spontaneously within the retina, perhaps by the ganglion cells themselves. If so, the firing of retinal ganglion cells might be contributing to layer construction, because all the synaptic machinery necessary for competition is present. It should be possible to prevent the formation of the eye-specific layers by blocking action-potential activity from the eyes to the lateral geniculate nucleus.

To hinder activity during fetal development, Sretavan and I, in collaboration with Stryker, implanted special minipumps containing tetrodotoxin in utero just before the lateral geniculate nucleus layers normally begin to form in the cat (at about six weeks of fetal development). After two weeks of infusion, we assessed the effects on the formation of layers. Much to our satisfaction, the results of these in utero in-

fusion experiments showed clearly that the eye-specific layers do not appear in the presence of tetrodotoxin. Moreover, by examining the branching patterns of individual ganglion cell axons after the treatment, we reassured ourselves that tetrodotoxin did not simply stunt normal growth.


In fact, the branching patterns of these axons were very striking. Unlike normal axons at the comparable age, the tetrodotoxin-treated axons did not have highly restricted terminal branches. Rather they had many branches along the entire length of the axon. It was as if, without action-potential activity, the information necessary to withdraw side branches and elaborate the terminal branches was missing.

In 1988, at about the same time these experiments were completed, Lucia Galli-Resta and Lamberto Maffei of the University of Pisa achieved the extraordinary technical feat of actually record-

ing signals from fetal ganglion cells in utero. They demonstrated directly that retinal ganglion cells can indeed spontaneously generate bursts of action potentials in the darkness of the developing eye. This observation, taken together with our experiment, strongly suggests that action-potential activity is not only present but also necessary for the ganglion cell axons from the two eyes to segregate and form the eye-specific layers.


Still, there must be constraints on the spatial and temporal patterning of ganglion cell activity. If the cells fired randomly, the mechanism of correlation-based, activity-dependent sorting could not operate. Furthermore, neighboring ganglion cells in each eye somehow ought to fire in near synchrony with one another, and the firing of cells in the two eyes, taken together, should be asynchronous. In addition, the synapses between retinal ganglion cell axons and neurons of the lateral geniculate nucleus should resemble Hebb synapses in their function: they should be

ACTION-POTENTIAL READING



ACTION-POTENTIAL READINGS of the developing retina are recorded by microelectrodes (*black spots*). The electrodes detect the small, extracellular currents that flow when the gan-

glion cells (*stained purple*) fire. All the cells fire at about the same time and then become silent before firing again. The area shown represents about 3 percent of the entire retina.



able to detect correlations in the firing of axons and strengthen accordingly.

We realized that to search for such patterns of spontaneous firing, it would be necessary to monitor simultaneously the action-potential activity of many ganglion cells in the developing retina. In addition, the observation had to take place as the eye-specific layers were developing. A major technical advance permitted us to achieve this goal. In 1988 Jerome Pine and his colleagues at the California Institute of Technology, among them doctoral student Markus Meister, invented a special multielectrode recording device. It consisted of 61 recording electrodes arranged as a flat, hexagonal array. Each electrode can detect action potentials generated in one to several cells. When Meister arrived at Stanford University to continue postdoctoral work with Denis Baylor, we began a collaboration to see whether the electrode array could be used to detect the spontaneous firing of fetal retinal ganglion cells.

In these experiments, it was necessary to remove the entire retina from the fetal eye and place it, ganglion-cell-side down, on the array. (It is technically impossible to put the electrode array itself into the eye in utero.) Rachel Wong, a postdoctoral fellow from Australia visiting my laboratory, succeeded in carefully dissecting the retinas and in tailoring special fluids necessary to maintain the living tissue for hours in a healthy condition.

When neonatal ferret retinas were placed on the multielectrode array, we simultaneously recorded the spontaneously generated action potentials of as many as 100 cells. The work confirmed the *in vivo* results of Galli-Resta and Maffei. All cells on the array fired within about five seconds of one another, in a predictable and rhythmic pattern. The bursts of action potentials lasted several seconds and were followed by long silent pauses that persisted from 30 seconds to two minutes. This observation showed that the activity of ganglion cells is indeed correlated. Further analysis demonstrated that the activity of neighboring cells is more highly correlated than that of distant cells on the array.

Even more remarkable, the spatial pattern of firing resembled a wave of activity that swept across the retina at about 100 microns per second (about one tenth to one hundredth the speed of an ordinary action potential). After the silent period, another wave was generated but in a completely different and random direction. We found that these spontaneously generated retinal waves are present throughout the period when eye-specific layers take shape. They disappear just before the onset of visual function.

From an engineering standpoint, these waves seem beautifully designed to provide the required correlations in the firing of neighboring ganglion cells. They also ensure a sufficient time delay, so that the synchronized firing of ganglion cells remains local and does not occur across the entire retina. Such a pattern of firing could help refine the topographic map conveyed by ganglion cell axons to each eye-specific layer. Moreover, the fact that wave direction appears to be entirely random implies that ganglion cells in the two eyes are highly unlikely ever to fire synchronously—a requirement for the formation of the layers.

Future experiments will disrupt the waves in order to determine whether they are truly involved in the development of connections. In addition, it will be important to determine whether the correlations in the firing of neighboring ganglion cells can be detected and used by the cells in the lateral geniculate nucleus to strengthen appropriate synapses and weaken inappropriate ones. This seems likely, since Richard D. Mooney, a postdoctoral fellow in my laboratory, in collaboration with Daniel Madison of Stanford, has shown that long-term potentiation of synaptic transmission between retinal ganglion cell axons and the lateral geniculate nucleus neurons is present during these early periods of development. Thus, at present, we can conclude that even before the onset of function, ganglion cells can spontaneously fire in the correct pattern to fashion the necessary connections.

Is the retina a special case, or might many regions of the nervous system generate their own endogenous activity

patterns early in development? Preliminary studies by Michael O'Donovan of the National Institutes of Health suggest that the activity of motor neurons in the spinal cord may also be highly correlated very early in development. It would appear that activity-dependent sorting in this system as well might use spontaneously generated signals. Like those in the visual system, the signals would refine the initially diffuse connections within targets.

The necessity for neuronal activity to complete the development of the brain has distinct advantages. The first is that, within limits, the maturing nervous system can be modified and fine-tuned by experience itself, thereby providing a certain degree of adaptability. In higher vertebrates, this process of refinement can occupy a protracted period. It can begin in utero and, as in the primate visual system, continue well into neonatal life, where it plays an important role in coordinating inputs from the two eyes. The coordination is necessary for binocular vision and stereoscopic depth perception.

Neural activity confers another advantage in development. It is genetically conservative. The alternative—exactly specifying each neural connection using molecular markers—would require an extraordinary number of genes, given the thousands of connections that must be formed in the brain. Using the rules of activity-dependent remodeling described here is far more economical. A major challenge for the future will be to elucidate the cellular and molecular bases for such rules.

FURTHER READING

PRENATAL DEVELOPMENT OF THE VISUAL SYSTEM IN THE RHESUS MONKEY. P. Rakic in *Philosophical Transactions of the Royal Society of London*, Series B, Vol. 278, No. 961, pages 245-260; April 26, 1977.

OCULAR DOMINANCE COLUMN DEVELOPMENT: ANALYSIS AND SIMULATION. Kenneth D. Miller, Joseph B. Keller and Michael P. Stryker in *Science*, Vol. 245, pages 605-615; August 11, 1989.

COMPETITIVE INTERACTIONS BETWEEN RETINAL GANGLION CELLS DURING PRENATAL DEVELOPMENT. Carla J. Shatz in *Journal of Neurobiology*, Vol. 21, No. 1, pages 197-211; January 1990.

IMPULSE ACTIVITY AND THE PATTERNING OF CONNECTIONS DURING CNS DEVELOPMENT. C. J. Shatz in *Neuron*, Vol. 5, No. 6, pages 745-756; December 1990.

DEVELOPMENT. Edited by Corey S. Goodman and Thomas M. Jessell. Special issue of *Current Opinion in Neurobiology*, Vol. 2, No. 1; February 1992.



The Visual Image in Mind and Brain

In analyzing the distinct attributes of images, the brain invents a visual world. Unusual forms of blindness show what happens when specialized parts of the cortex malfunction


by Semir Zeki

The study of the visual system is a profoundly philosophical enterprise; it entails an inquiry into how the brain acquires knowledge of the external world, which is no simple matter. The visual stimuli available to the brain do not offer a stable code of information. The wavelengths of light reflected from surfaces change along with alterations in the illumination, yet the brain is able to assign a constant color to them. The retinal image produced by the hand of a gesticulating speaker is never the same from moment to moment, yet the brain must consistently categorize it as a hand. An object's image varies with distance, yet the brain can ascertain its true size.

The brain's task, then, is to extract the constant, invariant features of objects from the perpetually changing flood of information it receives from them. Interpretation is an inextricable part of sensation. To obtain its knowledge of what is visible, the brain cannot therefore merely analyze the images presented to the retina; it must actively construct a visual world. To do so, the brain has developed an elaborate neural mechanism, one so marvelously efficient that it took a century of study before anyone even guessed at its many components. Indeed, when studies of cerebral diseases began to reveal some secrets of the visual brain, neurologists initially dismissed the startling implications as improbable.

The hallmark of that machinery is a complex division of labor. It is manifested anatomically in discrete cortical areas and subregions of areas specialized for particular visual functions; it is manifested pathologically in an inability to acquire knowledge about some aspect of the visual world when the relevant machinery is specifically compromised. Paradoxically, none of this subdivision and specialization within the brain is normally evident at the perceptual level. The visual cortex thus presents us with the intellectual challenge of trying to understand how its components cooperate to give us a unified picture of the world, one that bears no trace of the division of labor within it. There is, to use an old phrase, a great deal more to vision than meets the eye.

This modern conception of the visual brain has evolved



only within the past two decades. The early neurologists, starting with those who worked during the late 19th century, saw it very differently. Laboring under the false notion that objects transmitted visual codes in reflected or emitted light, they thought that an image was “impressed” on the retina, much as it would be on a photographic plate. These retinal impressions were subsequently transmitted to the visual cortex, which served to analyze the contained codes. This decoding process led to “seeing.” Understanding what was



seen—making sense of the received impressions and resolving them into visual objects—was thought to be a separate process that arose through the association of the received impressions with similar ones experienced previously.

This view of how the brain operates, which persisted into the mid-1970s, was therefore also deeply philosophical, although neurologists never acknowledged it as such. It divided sensing from understanding and gave each faculty a separate seat in the cortex. The origin of this dualistic doctrine is obscure, but it bears a resemblance to Immanuel Kant's belief in the two faculties of sensing and of understanding, the former passive and the latter active.

Neurologists saw evidence for their supposition in the fact that the retina connects overwhelmingly to one distinct part of the brain, the striate or primary visual cortex, also known as area V1. This connection is made with high topographic precision: V1 effectively contains a map of the entire retinal field. The retina and V1 are linked through a subcortical structure called the lateral geniculate nucleus, which contains six layers of cells. The four uppermost layers contain cells with small cell bodies and are referred to as the parvocellular layers; the two lowest layers, which have large cell bodies, are the magnocellular layers. Many years ago the late neurologist Salomon E. Henschen of Uppsala University supposed that the function of the large cells was “collecting

WHEN VIEWING AN IMAGE, distinct areas in the cortex analyze it for different visual attributes, such as color, shape and motion. “Seeing” and “understanding” occur simultaneously through the synchronized activities of these cortical areas. The world that one sees is an invention of the visual brain.

SEMIR ZEKI is professor of neurobiology at the University of London. He obtained his doctorate from University College, London, and did his postdoctoral work at the National Institute of Mental Health in Washington, D.C., and at the University of Wisconsin at Madison. Zeki has also served as a visiting professor at several American and European universities. His particular interests center on the study of the anatomic and functional organization of the visual cortex in the monkey and, more recently, in the human brain.



ANATOMIC AND FUNCTIONAL DIVISIONS within the visual system are the physical foundation for vision. Most connections between the retina and the visual cortex at the back of the brain pass through the lateral geniculate nucleus. In cross section, this subcortical structure has six cell layers: two in the magnocellular pathway (*M*) and four in the parvocellular pathway (*P*).

light" and that of the small cells was registering colors; his basic insight, that the anatomic subdivisions had functional implications, has assumed increasing importance in recent years.



Neurologists at the time found that lesions anywhere along the pathway connecting the retina with V1 created a field of absolute blindness, the extent

and position of which corresponded precisely with the size and location of the lesion in V1. That observation led Henschen to conceive of area V1 as the "cortical retina"—the place where "seeing" occurred.

Moreover, the German psychiatrist Paul Emil Flechsig of Leipzig University had shown during the late 19th cen-

tury that certain regions of the brain, among them V1, had a mature appearance at birth, whereas others, including the cortical regions surrounding V1, continued to develop, as though their maturation depended on the acquisition of experience. For Flechsig and most other neurologists, this observation implied that V1 was "the enter-

MACAQUE MONKEY BRAIN



VISUAL CORTEX of the macaque monkey has been studied in detail. A cross section through the brain (*left*) at the level indi-

cated (*right*) shows part of the primary visual cortex (V1) and some of the other visual areas in the prestriate cortex (V2-V5).

ing place of the visual radiation into the organ of the psyche," and the areas around it were the repositories of higher "psychic" functions (*Cogitationzentren*) related to sight. Flechsig's theory found support in rather questionable evidence purporting to show that lesions in this so-called visual association cortex, unlike those in V1, might lead to "mind blindness" (*Seelenblindheit*), a condition in which subjects were thought to see but not to comprehend what they saw.

Surprisingly, it was research on the visual association cortex that ultimately compromised this dualistic concept of visual brain organization. Work undertaken in the 1970s by John M. Allman and Jon H. Kaas of the University of Wisconsin in the owl monkey and by me in the macaque monkey showed that the visual association cortex—now better referred to as the prestriate cortex—consists of many different cortical areas separated from V1 by another area, V2. A turning point in our understanding of how the brain constructs the visual image came subsequently, with my demonstration that these areas are individually specialized to undertake different tasks.

In my physiological studies, I presented macaques with a range of stimuli (colors, lines of various orientations and dots moving in different directions) and, using electrodes, monitored the activity of cells in the prestriate cortex. The results showed that all the cells in a prestriate area called V5 are responsive to motion, that most are directionally selective and that none is concerned with the color of the moving stimulus. These facts suggested to me that V5 is specialized for visual motion. (Neuroanatomic terminology is not always uniform; some investigators prefer the label MT to V5.)

In contrast, I found that the overwhelming majority of cells in another area, V4, are to some extent selective for specific wavelengths of light and that many are selective for line orientation, the constituents of form, as well. By far most of the cells in two further adjoining areas, V3 and V3A, are also selective for form but like the cells of area V5 are largely indifferent to the color of the stimulus.

These studies led me to propose in the early 1970s the concept of functional specialization in the visual cortex, which supposes that color, form, motion and possibly other attributes of the visible world are processed separately. Because the preponderance of input to the specialized areas comes from V1, a corollary of this finding was


that V1 must also show a functional specialization, as must area V2, which receives input from V1 and connects with the same specialized areas. These two areas must, in a sense, act as a kind of post office, parceling out different signals to the appropriate areas.

In recent years, new tissue-staining techniques in combination with physiological studies have provided a startling confirmation of that theory. They have also allowed us to trace these specializations from V1 throughout the prestriate cortex.

With the advent of positron emission tomography (PET), which can measure increases in regional cerebral blood flow when people perform specific tasks, my colleagues at the Hammersmith Hospital in London and I have begun to apply

these findings, which were derived from experiments on monkeys, to a direct study of the human brain. We found that when normal-seeing humans view a Land color Mondrian (an abstract painting containing no recognizable objects), the highest increase in regional cerebral blood flow occurs in a structure named the fusiform gyrus. By analogy with a similar region in macaque monkeys, we refer to this cortical area as human V4. The results are very different when subjects view a pattern of moving black-and-white squares: the highest cerebral blood flow then occurs in a more lateral area, quite separate from V4, which we call human V5.

This demonstration of the separation of motion and color processing constitutes direct evidence that functional



DISSIMILAR IMAGES stimulate different regions of the visual cortex. A brightly colored Mondrian causes area V4 to become highly active, as shown by tests of regional cerebral blood flow. Black-and-white moving images trigger activity in area V5. Both types of images lead to activity in areas V1 and V2, which have less specialized functions and distribute signals to other cortical areas.

specialization is also a feature of the human visual cortex. The PET studies reveal another interesting feature: under both conditions of stimulation, area V1 (and probably the adjoining area V2) also showed marked increases in regional cerebral blood flow. As in the monkeys, these regions, too, must be distributing signals to different areas of the prestriate cortex.

The key to the distribution system in these areas lies in their structural and functional organization. Area V1 is unusually rich in cell layers, yet it reveals an even richer architecture if one examines it with a staining technique first applied by Margaret Wong-Riley of the Medical College of Wisconsin in Milwaukee. The organelles known as mitochondria contain a metabolic enzyme called cytochrome oxidase that makes energy available to a cell. By staining a region of the brain for that enzyme, research-

ers can identify which cells have the greatest metabolic activity.


When so stained, the metabolic architecture of V1 is characterized by columns of cells that extend from the cortical surface to the underlying nerve tissue called white matter. If viewed in sections cut parallel to the cortical surface, these columns appear as heavily stained blobs or puffs, separated from one another by more lightly stained interblob regions. At Harvard Medical School, Margaret Livingstone and David H. Hubel found that wavelength-selective cells are concentrated in the blobs of V1, whereas form-selective cells are concentrated in the interblobs.

The columns are especially prominent in the second and third layers of V1, which receive input from the parvocellular layers of the lateral geniculate nucleus. The cells in those parts of the lateral geniculate nucleus respond

in a strong, sustained way to visual stimuli, and many of them are concerned with color.

A distinct set of structures can be seen in layer 4B of V1, which receives input from the magnocellular layers of the lateral geniculate nucleus, whose cells respond transiently to stimuli and are mostly indifferent to color. Layer 4B projects to areas V5 and V3. The cells in layer 4B that connect with V5 are clustered into small patches that are isolated by cells connected to other visual areas. In short, the organization of layer 4B in V1 suggests that certain parts of it are specialized for motion perception and are segregated from regions that handle other attributes.

Like V1, area V2 has a special metabolic architecture. In the case of V2, however, that architecture takes the shape of thick stripes and thin stripes separated from one another by more



lightly staining interstripes. As work done by Edgar A. DeYoe and David C. Van Essen of the California Institute of Technology, by Hubel and Livingstone, and by Stewart Shipp of University College, London, and me has shown, cells selective for wavelength congregate in the thin stripes, and cells selective for directional motion are found in the thick stripes. Cells sensitive to form are distributed in both the thick stripes and the interstripes.

V1 and V2 might therefore be said to contain pigeonholes into which the different signals are assembled before being relayed to the specialized visual areas. The cells in these pigeonholes have small receptive fields; that is, they respond only to stimuli falling on a small region of the retina. They also register information about only a specific attribute of the world within that receptive field. It is as though V1 and V2 were undertaking a piecemeal analysis of the entire field of view.

These facts allow us to delineate four parallel systems concerned with different attributes of vision—one for motion, one for color and two for form. The two that are computationally most distinct from each other are the motion and color systems. For the motion system, the pivotal prestriate area is V5; its inputs run from the retina, through the magnocellular layers of the lateral geniculate nucleus, to layer 4B of V1. From there the signals pass to V5, both directly and through the thick stripes of V2. The color system depends on area V4; its inputs pass through the parvocellular layers of the lateral geniculate nucleus to the blobs of V1, then proceed to V4 directly or through the thin stripes of V2.

Of the two form systems, one is intimately linked to color, and the other is independent of it. The first is based on V4 and derives its inputs from the parvocellular layers of the lateral geniculate nucleus by way of the interblobs of V1 and the interstripes of V2. The second is based on V3 and is more concerned with dynamic form—the shapes of objects in motion. It derives its inputs from the magnocellular layers of the lateral geniculate nucleus through layer 4B of V1; the signals then proceed to V3 both directly and through the thick stripes of V2.

Although these four systems are distinct, the anatomy of areas V1 and V2 offers many opportunities for the pigeonholes to communicate with one another, as do the direct connections between the specialized visual areas. Hence, there is an admixture of the parvocellular and magnocellular signals, which the prestriate areas use in different ways to execute their functions.

This remarkable segregation of functions is reflected in some of the pathologies afflicting the visual cortex. Lesions in specific cortical areas produce correspondingly specific visual syndromes that may be far less debilitating than total blindness yet still severe enough to drive patients to distraction and despair. Lesions in area V4 lead to achromatopsia, in which patients see only in shades of gray. This syndrome is different from simple color blindness: not only do such patients fail to see or know the world in color, they cannot even recall colors from a time before the lesion formed. Nevertheless, if their retinas and V1 regions are healthy, their knowledge of form, depth and motion remains intact.

Similarly, a lesion in area V5 produces akinetopsia, in which patients neither see nor understand the world in motion. While at rest, objects may be perfectly visible to them, but motion relative to them causes the objects to vanish. The other attributes of vision remain unscathed—a specificity that results directly from functional differentiation in the human visual cortex.

Given the separation of form and color in the cortex, it is perhaps a little surprising that no one has ever reported a complete and specific loss of form vision. A partial explanation is that such a deficit would require the obliteration of areas V3 and V4 to eliminate both form systems. Area V3 forms a ring around V1 and V2. Consequently, a lesion large enough to destroy all of V3 and V4 would almost certainly destroy V1 as well and thus cause total blindness.

Some patients with lesions in the prestriate cortex do suffer from a degree of form imperception, often coupled with achromatopsia. These people commonly experience far greater difficulty when identifying stationary forms

than when the same forms are in motion. They frequently prefer watching television to watching the real world, because television is dominated by moving images. When faced with stationary objects, these patients often resort to the strategy of moving their heads to simplify the task of identification. These observations suggest that they acquire their knowledge of forms through the dynamic form system based in area V3.

The functional specialization in the visual cortex also manifests itself in a syndrome that I have called the chromatopsia (“color vision”) of carbon monoxide poisoning. This condition has been described sporadically but not infrequently in the medical literature, although it was never taken seriously until the functional specialization was discovered. Some people who survive the lethal effects of smoke inhalation during fires often suffer diffuse cortical damage from carbon monoxide poisoning, which deprives tissues of oxygen. As a result, these patients often have vision that is severely compromised in all respects except one: their color vision is affected only mildly if at all. Because color is the only kind of visual knowledge available to them, the patients try—often unsuccessfully—to identify all objects solely on the basis of color. They may, for example, misidentify all blue objects as “ocean.”

The precise cause of this strange chromatopsia is unknown. The metabolically active blobs of V1 and the thin stripes of V2, both of which are concerned with color, do have unusually high concentrations of blood vessels nourishing them. It is therefore probable that these regions are relatively spared from damage because their rich blood supply renders them less vulnerable to oxygen deprivation.

In summary, then, we know that a total lesion in area V1 produces a complete inability to acquire any visual information and that a lesion in one of the specialized areas makes a corresponding attribute of the visual world inaccessible and incomprehensible. What would happen, we might ask, if signals from the lateral geniculate nucleus were routed directly to the specialized areas, thus bypassing V1 altogether? Nature has actually done that experiment for us, and the resulting phenomenon provides more important insights into the functioning of the visual cortex.

The phenomenon is known as blindsight; it was first described by Ernst Pöppel of the University of Munich and his colleagues and later studied in great detail by Lawrence Weiskrantz of the

FOUR PERCEPTUAL PATHWAYS within the visual cortex have been identified. Color is seen when wavelength-selective cells in the blob regions of V1 send signals to specialized area V4 and also to the thin stripes of V2, which connect with V4. Form in association with color depends on connections between the interblobs of V1, the interstripes of V2 and area V4. Cells in layer 4B of V1 send signals to specialized areas V3 and V5 directly and also through the thick stripes of V2; these connections give rise to the perception of motion and dynamic form.

University of Oxford and his colleagues. People with this condition are totally blind because of lesions in area V1. Yet if they are forced to guess, they can discriminate correctly among a wide variety of visual stimuli. They can, for example, distinguish between motion in different directions or between different wavelengths of light. Their abilities are imperfect and not completely reliable, but they are better than random guessing. Nevertheless, blindsight patients are not consciously aware of having seen anything at all, and they are often surprised that their "guesses" should have been so accurate.

The basis for this discrimination almost certainly resides in a small but direct connection between the lateral geniculate nucleus and the prestriate cortex, as uncovered by Masao Yuki of the Tokyo Metropolitan Institute for Neurosciences and by Wolfgang Fries of the University of Munich. Alternatively, some other as yet undiscovered subcortical connection to the specialized areas may be responsible. In any event, neurologists have good reason to suppose that in blindsight patients, visual signals reach the prestriate cortex.

Blindsight patients are people who "see" but do not "understand." Because they are unaware of what they have seen, they have not acquired any knowledge. In short, their "vision," which can be elicited only in laboratory situations, is quite useless. Thus, for the visual cortex to do its job of acquiring a knowledge of the world, a healthy V1 area is essential. V1 (and, by extension, V2) may be necessary because it begins to process information for further refinement by the specialized areas or because the results of the processing performed by the specialized areas are referred back to it.

The clinical literature holds many other examples that illuminate how the preprocessing in areas V1 and V2 may contribute directly and explicitly to perception. Damage to V5 can destroy the ability to discriminate the direction or coherence of motions. Yet as Robert F. Hess of the University of Cambridge and his colleagues found, such akinetopsic patients may still be aware that motion of some type is taking place, presumably because of signaling by cells in V1 and V2 (and possibly in other areas that receive magnocellular pathway signals). Similarly, an achromatopsic patient with a V4 lesion, whom I studied with Fries, could discriminate between different wavelengths because of his largely intact V1 even though he could no longer interpret the wavelength information as color.

Further insights come from a comparison of the residual form-vision capacities in two other patients who have cortical lesions that are, in a way, complementary. The first patient has a diffuse cortical lesion, caused by carbon monoxide poisoning, that affects area V1. He has terrific difficulty copying even simple forms, such as geometric shapes or letters of the alphabet, because the form-detecting system in his V1 area is so severely compromised.

The second patient has an extensive prestriate lesion from a stroke that has generally spared area V1. He can reproduce a sketch of St. Paul's Cathedral with greater skill than many normal people, although it takes him a great deal of time to do so. Yet this patient has no comprehension of what he has drawn. Because his V1 system is largely

intact, he can identify the local elements of form, such as angles and simple shapes, and accurately copy the lines he sees and understands. The prestriate lesion, however, prevents him from integrating the lines into a complex whole and recognizing it as a building. The patient sees and understands only what the limited capacity of his intact system allows.

The residual capacity in such patients unmasks an important feature of the organization of the visual cortex, namely, that none of the visual areas—not even "post office" areas V1 and V2—serves merely to relay signals to other areas. Instead each is part of the machinery that actively transforms the incoming signals and may contribute explicitly, if incompletely, to perception.

The profound division of labor with-

The World Seen through a Damaged Cortex

Damage to specialized regions of the cortex can cause strange types of blindness in which patients lose the ability to see just one attribute of the visual world, such as color, form or motion. Artwork produced by some of these patients offers glimpses into their view of the world, as well as into the workings of the visual cortex itself.



A patient with damage to the color pathways in the cortex lost all color vision. In his drawings, a banana, a tomato and green leaves all have similar colors.



This stroke patient suffered damage to the prestriate cortex that impaired his perception of form. He could copy a drawing with great skill, but he was unable to understand that the lines he had drawn produced an image of St. Paul's Cathedral.

in the visual cortex naturally raises the question of how the specialized areas interact to provide a unified image. The simplest way would be for all the specialized areas to communicate the results of their operations to one master area, which would then synthesize the incoming information. Philosophically, that solution begs the question, because one must then ask who or what looks at the composite image, and how it does so. That problem is beside the point, however, because the anatomic evidence shows no single master area to which all the antecedent areas exclusively connect. Instead the specialized areas connect with one another, either directly or through other areas.


For example, areas V4 and V5 connect directly and reciprocally with each other. Both of them also project to the pari-

etal and temporal regions of the brain, but as my work with my colleagues has shown, the outputs from each area occupy their own unique territory within the receiving region. Direct overlap between the signals from V4 and V5 is minimal. It is as if the cortex wishes to maintain the separation of the distinct visual signals—a strategy it also employs in memory and other systems [see “Working Memory and the Mind,” by Patricia S. Goldman-Rakic, page 110]. Any integration of the signals within the parietal or temporal regions must occur through local “wiring” that connects the inputs.

In fact, integration of the visual information is a monumental task that necessitates a vast network of anatomic links between the four parallel systems at every level, because each level con-


tributes explicitly to perception. Integration also creates some formidable problems. To understand coherent motion, for example, the brain must determine which features in the field of view are moving in the same direction and at the same speed. The motion-sensing cells in the specialized areas are able to make those comparisons because they have larger receptive fields than do their antecedent counterparts in V1.

Yet because their receptive fields are larger, these cells are inherently less efficient at pinpointing the position of any one stimulus within the visual field. For the brain to make spatial sense of the integrated information, the information must somehow be referred to an area that has a more precise topographic map of the retina and hence of the visual field. Of all the visual areas, the




When an achromatopsic (color-blind) patient was shown a Land color Mondrian (left) and asked to reproduce it, he


was able to copy the shapes in the painting successfully. The colors within the blocks eluded him (right).



This patient sustained damage from carbon monoxide poisoning and lost his perception of form. At the left, he was asked to choose which of four selections was identical



to a given shape; as the marks on his choices reveal, he could not identify them correctly. At the right, his inability to copy simple shapes is displayed.



KANIZSA TRIANGLE consists of illusory contours. A normal visual cortex sees a triangle even though interconnecting lines are missing. Such illusions show that the visual cortex must resolve conflicts between different functional areas.

one with the most precise map is area V1, followed by V2. The specialized areas must therefore send information back to V1 and V2 so that the results of the comparisons can be mapped back onto the visual field.

Reentrant connections, which allow information to flow both ways between different areas, are also essential for resolving conflicts between cells that have different capabilities and are responding to the same stimulus. A good example of such conflicts is found in the responses of cells in V1 and V2 to illusory contours, such as those of the Kanizsa triangle.

In this famous illusion, a normal observer perceives a triangle among the presented shapes even though the lines forming the triangle are incomplete; the brain creates lines where there are none. As Rudiger von der Heydt and Esther Peterhans of the Zurich University Hospital have demonstrated, the form-selective cells in V1 do not respond to the illusion and do not signal that a line is present. The V2 cells receive their inputs from V1, but because the V2 cells have larger receptive fields and more analytic functions, they do respond to the illusion by "inferring" the presence of a line. To settle the conflict, the V2 cells must have reentrant inputs to their counterparts in V1.

Another difficulty that arises from the process of integration is the binding problem. Cells responding to the same object in the field of view may be scattered throughout V1. Something must therefore bind together the signals from those cells so that they are treated as belonging to the same object and not separate ones. The problem becomes even more thorny when cells in two or

more visual areas respond to different attributes of the same object.

One way to resolve the problem is for the cells to fire in temporal synchrony. In practice, such synchrony does occur, at least to some extent, among cells that are anatomically connected to one another, as work by Wolf J. Singer of the Max Planck Institute for Brain Research in Frankfurt and his colleagues has shown. We must then face the problem, however, of who or what determines that the firing should be synchronous. Reentrant inputs provide at least a partial solution by linking the output of one area to other areas that sent it information.

These problems have led me and my colleagues to develop a theory of multistage integration. It hypothesizes that integration does not occur in a single step through a convergence of output onto a master area, nor does integration have to be postponed until all the visual areas have completed their individual operations. Instead the integration of visual information is a process in which perception and comprehension of the visual world occur simultaneously.

The anatomic requirement for multistage integration is immense, because it involves reentrant connections between all the specialized areas as well as with areas V1 and V2, which feed them signals. Our studies indicate that such a network of reentrant connections does exist.

The reentrant inputs to areas V1 and V2 differ fundamentally from their forward connections to the specialized areas. The forward projections are patchy and discrete, because segregated groups of cells in V1 and V2 send their outputs to specialized areas with corresponding visual attributes. The return projections, however, are diffuse and fairly nonspecific. For example, whereas V5 receives input only from select groups of cells in layer 4B of V1, the return input from V5 to layer 4B is diffuse and encompasses the territory of all the cells in the layer, including ones that project into V3. This reentrant system can thus serve three purposes simultaneously: it can unite and synchronize the signals for form and motion found in two different visual pathways; it can refer information about motion back to an area with an accurate topographic map; it can integrate motion information from V5 with form information on its way to V3.

Similarly, whereas the output from V2 to the specialized areas is highly segregated, the return input from those areas to V2 is diffuse. V4 projects back

not only to the thin stripes and the interstripes, from which it receives its input, but also to the thick stripes, from which it does not. This reentrant system can therefore help unite signals dealing with form, motion and color.



It is becoming increasingly evident that the entire network of connections within the visual cortex, including the reentrant connections to V1 and V2, must function healthily for the brain to gain complete knowledge of the external world. Yet as patients with blindsight have shown, knowledge cannot be acquired without consciousness, which seems to be a crucial feature of a properly functioning visual apparatus. Consequently, no one will be able to understand the visual brain in any profound sense without tackling the problem of consciousness as well [see "The Problem of Consciousness," by Francis Crick and Christof Koch, page 152].

The past two decades have brought neurologists many marvelous discoveries about the visual brain. Moreover, they have led to a powerful conceptual change in our view of what the visual brain does and how it accomplishes its functions. It is no longer possible to divide the process of seeing from that of understanding, as neurologists once imagined, nor is it possible to separate the acquisition of visual knowledge from consciousness. Indeed, consciousness is a property of the complex neural apparatus that the brain has developed to acquire knowledge.

Thus, our inquiry into the visual brain takes us into the very heart of humanity's inquiry into its own nature. This is not to say that understanding the workings of the visual brain will resolve the problem of consciousness—far from it. But it is a good beginning.

FURTHER READING

FUNCTIONAL SPECIALISATION IN THE VISUAL CORTEX OF THE RHESUS MONKEY. S. M. Zeki in *Nature*, Vol. 274, No. 5670, pages 423-428; August 3, 1978.
BLINDSIGHT: A CASE STUDY AND IMPLICATIONS. L. Weiskrantz. Oxford University Press (Clarendon Press), 1986.
SEGREGATION OF FORM, COLOR, MOVEMENT, AND DEPTH: ANATOMY, PHYSIOLOGY, AND PERCEPTION. Margaret Livingstone and David Hubel in *Science*, Vol. 240, pages 740-749; May 6, 1988.
THE FUNCTIONAL LOGIC OF CORTICAL CONNECTIONS. S. Zeki and S. Shipp in *Nature*, Vol. 335, No. 6188, pages 311-317; September 22, 1988.
THE REMEMBERED PRESENT: A BIOLOGICAL THEORY OF CONSCIOUSNESS. Gerald M. Edelman. Basic Books, 1990.
A VISION OF THE BRAIN. S. Zeki. Blackwell Scientific Publications (in press).



Announcing an interruption of tonight's regular programming

Now you can share with your family and friends your own enthusiasm for science with a holiday gift of the award-winning SCIENTIFIC AMERICAN Frontiers television series. This five-hour scan of the universe of science puts the viewer where the scientists are and tells their exciting stories in non-technical language. SCIENTIFIC AMERICAN Frontiers is guaranteed to kindle the flames of insight and understanding.

Go behind the scenes at the Moscow circus to the engineering department; take a quick course in the physics of roller coasters; get a progress report on the test-tube truffles that may one day compensate for the dramatic fall-off in the French harvests; meet the Russian mathematician who invented Tetris™ and the Alaskan ground squirrel that drops its body temperature to below freezing when it hibernates.

The premier season of SCIENTIFIC AMERICAN Frontiers earned positive reviews all across America. This wide acceptance assures your selection of SCIENTIFIC AMERICAN Frontiers as a "perfect gift." Take advantage of this special gift offer. Give five full hours of fascination with just one phone call (or use the order coupon). You'll be gratefully remembered.

SPECIAL GIFT-GIVING OFFER

Five 1-hour cassettes, handsomely packaged and gold stamped,

\$99.95

plus \$5 postage and handling.
U.S. funds. Please add sales tax in:
CA, IA, IL, MA, MI, NY.*



CALL TOLL-FREE
1-800-777-0444

Have your Visa or MasterCard number handy.



SCIENTIFIC AMERICAN Frontiers
Dept. SAF
415 Madison Avenue
New York, NY 10017

Produced by: Chedd-Angier Production Company in association with SCIENTIFIC AMERICAN Magazine.
SCIENTIFIC AMERICAN Frontiers PBS Series © 1990, 1991
Chedd-Angier Production Company, Inc. All rights reserved. Made possible by a grant from **GTB**

Tetris™ v/o ELECTRONORGTECHNICA ("ELORG").

GUARANTEE

If for any reason you are dissatisfied with the program, simply return it and a complete refund will be sent to you. If there is a problem with an individual cassette, return it and it will be replaced at no charge.

© 1992 SCIENTIFIC AMERICAN, INC

ORDER FORM

SAFB9

Please enter my order for _____ set(s) of SCIENTIFIC AMERICAN Frontiers VHS cassettes for \$99.95 plus \$5.00 postage and handling. Please add sales tax in CA, IA, IL, MA, MI, NY.*

Name _____ (please print)
Address _____ Apt. _____
City _____ State _____ Zip _____

GIFT ORDER FORM

Shipping address if different from above:


Name _____ (please print)
Address _____ Apt. _____
City _____ State _____ Zip _____

My check/money order for

\$ _____ is enclosed.
Charge my: ☐ Visa ☐ MasterCard

Card no. _____
Expires _____
Signature _____

*In Canada, please add GST (R127387652). Outside U.S. and Canada: \$200.00 U.S. funds drawn on U.S. bank.
Attach list with names and addresses of additional recipients.
Detach coupon and mail to: SCIENTIFIC AMERICAN Frontiers; Dept. SAF; 415 Madison Avenue; New York, NY 10017.



The Biological Basis of Learning and Individuality


Recent discoveries suggest that learning engages a simple set of rules that modify the strength of connections between neurons in the brain. These changes play an important role in making each individual unique

by Eric R. Kandel and Robert D. Hawkins

Over the past several decades, there has been a gradual merger of two originally separate fields of science: neurobiology, the science of the brain, and cognitive psychology, the science of the mind. Recently the pace of unification has quickened, with the result that a new intellectual framework has emerged for examining perception, language, memory and conscious awareness. This new framework is based on the ability to study the biological substrates of these mental functions. A particularly fascinating example can be seen in the study of learning. Elementary aspects of the neuronal mechanisms important for several different types of learning can now be studied on the cellular and even on the molecular level. The analysis of learning may therefore provide the first insights into the molecular mechanisms underlying a mental process and so begin to build a bridge between cognitive psychology and molecular biology.

Learning is the process by which we acquire new knowledge, and memory is the process by which we retain that knowledge over time. Most of what we know about the world and its civilizations we have learned. Thus, learning and memory are central to our sense of individuality. Indeed, learning goes beyond the individual to the transmission of culture from generation to generation. Learning is a major vehicle for behavioral adaptation and a powerful force for social progress. Conversely, loss of memory leads to loss of contact with one's immediate self, with one's life history and with other human beings.

Until the middle of the 20th century, most students of behavior did not believe that memory was a distinct mental function independent of movement, perception, attention and language. Long after those functions had been localized to different regions of the brain, researchers still doubted that memory could ever be assigned to a specific region. The first person to do so was Wilder G. Penfield, a neuro-



surgeon at the Montreal Neurological Institute.

In the 1940s Penfield began to use electrical stimulation to map motor, sensory and language functions in the cortex of patients undergoing neurosurgery for the relief of epilepsy. Because the brain itself does not have pain receptors, brain surgery can be carried out under local anesthesia in fully conscious patients, who can describe what they experience in response to electric stimuli applied to different cortical areas. Penfield explored the cortical surface in more than 1,000

patients. Occasionally he found that electrical stimulation produced an experiential response, or flashback, in which the patients described a coherent recollection of an earlier experience. These memorylike responses were invariably elicited from the temporal lobes.

Additional evidence for the role of the temporal lobe in memory came in the 1950s from the study of a few patients who underwent bilateral removal of the hippocampus and neighboring regions in the temporal lobe as treatment for epilepsy. In the first and best-studied case, Brenda Milner of the Montreal Neurological Institute described a 27-year-old assembly-line worker, H.M., who had suffered from untreatable and debilitating temporal lobe seizures for more than 10 years. The surgeon William B. Scoville removed the medial portion of the temporal lobes on both sides of H.M.'s brain. The seizure disturbance was much improved. But immediately after the operation, H.M. experienced a devastating memory deficit: he had lost the capacity to form new long-term memories.

Despite his difficulty with the formation of new memories, H.M. still retained his previously acquired long-term memory store. He remembered his name, retained a perfectly good use of language and kept his normal vocabulary; his IQ remained in the range of bright-normal. He remembered well the

MIRROR DRAWING EXPERIMENT in patients with temporal lobe lesions gave the first hint, in 1960, that there are two distinct types of learning systems. One form, which is spared by the lesions, involves tasks that have an automatic quality such as the skilled movements illustrated in this experiment. The subject, who can see his hand only in the mirror, tries to trace the shape of a star. The second type of learning depends on conscious awareness and cognitive processes and is abolished by the lesions.

ERIC R. KANDEL and ROBERT D. HAWKINS have collaborated on studies of the neurobiology of learning. Kandel is University Professor at the College of Physicians and Surgeons of Columbia University and senior investigator at the Howard Hughes Medical Institute. He received an A.B. from Harvard College, an M.D. from the New York University School of Medicine and psychiatric training at Harvard Medical School. Hawkins received a B.A. from Stanford University and a Ph.D. in experimental psychology from the University of California, San Diego. He is associate professor in the Center for Neurobiology and Behavior at Columbia.

events that preceded the surgery, such as the job he had held, and he remembered vividly the events of his childhood. Moreover, H.M. still had a completely intact short-term memory. What H.M. lacked, and lacked profoundly, was the ability to translate what he learned from short-term to long-term memory. For example, he could converse normally with the hospital staff, but he did not remember them even though he saw them every day.

The memory deficit following bilateral temporal lobe lesions was originally thought to apply equally to all forms of new learning. But Milner soon discovered that this is not the case. Even though patients with such lesions have profound deficits, they can accomplish certain types of learning tasks as well as normal subjects can and retain the memory of these tasks for long periods. Milner first demonstrated this residual memory capability in H.M. with the discovery that he could learn new motor skills normally [see illustration on page 78]. She, and subsequently Elizabeth K. Warrington of the National Hospital for Nervous Diseases in London and Lawrence Weiskrantz of the University of Oxford, found that patients such as H.M. can also acquire and retain memory for elementary kinds of learning that involve changing the strength of reflex responses, such as habituation, sensitization and classical conditioning.

It immediately became apparent to students of behavior that the difference between types of learning that emerged from studies of patients with temporal lobe lesions represented a fundamental

psychological distinction—a division in the way all of us acquire knowledge. Although it is still not clear how many distinct memory systems there are, researchers agree that lesions of the temporal lobes severely impair forms of learning and memory that require a conscious record. In accordance with the suggestion of Neal J. Cohen of the University of Illinois and Larry R. Squire of the University of California at San Diego and of Daniel L. Schacter of the University of Toronto, these types of learning are commonly called declarative or explicit. Those forms of learning that do not utilize conscious participation remain surprisingly intact in patients with temporal lobe lesions; they are referred to as nondeclarative or implicit.

Explicit learning is fast and may take place after only one training trial. It often involves association of simultaneous stimuli and permits storage of information about a single event that happens in a particular time and place; it therefore affords a sense of familiarity about previous events. In contrast, implicit learning is slow and accumulates through repetition over many trials. It often involves association of sequential stimuli and permits storage of information about predictive relations between events. Implicit learning is expressed primarily by improved performance on certain tasks without the subject being able to describe just what has been learned, and it involves memory systems that do not draw on the contents of the general knowledge of the individual. When a subject such as H.M. is asked why he performs a given



task better after five days of practice than on the first day, he may respond, "What are you talking about? I've never done this task before."

Whereas explicit memory requires structures in the temporal lobe of vertebrates, implicit memory is thought to be expressed through activation of the particular sensory and motor systems engaged by the learning task; it is acquired and retained by the plasticity inherent in these neuronal systems. As a result, implicit memory can be studied in various reflex systems in either vertebrates or invertebrates. Indeed, even simple invertebrate animals show excellent reflexive learning.

The existence of two distinct forms of learning has caused the reductionists among neurobiologists to ask whether there is a representation on the cellular level for each of these two types of learning process. Both the neural systems that mediate explicit memory and those that mediate implicit memory can store information about the association of stimuli. But does the same set of cellular learning rules guide the two memory systems as they store associations, or do separate sets of rules govern each system?

An assumption underlying early studies of the neural basis of memory systems was that the storage of associative memory, both implicit and explicit, required a fairly complex neural circuit. One of the first to challenge this view was the Canadian psychologist Donald O. Hebb, a teacher of Milner. Hebb boldly suggested that associative learning could be produced by a simple cellular mechanism. He proposed that associations could be formed by coincident neural activity: "When an axon of cell A...excite[s] cell B and repeatedly or persistently takes part in firing it, some growth process or metabolic change takes place in one or both cells such that A's efficacy, as one of the cells firing B, is increased." According to Hebb's learning rule, coincident activity in the presynaptic and postsynaptic neurons is critical for strengthening the connection between them (a so-called pre-post associative mechanism) [see illustration at left].

Ladislav Tauc and one of us (Kandel) proposed a second associative learning rule in 1963 while working at the Institute Mery in Paris on the nervous system of the marine snail *Aplysia*. They found that the synaptic connection between two neurons could be strengthened without activity of the postsynaptic cell when a third neuron acts on the presynaptic neuron. The third neuron, called a modulatory neuron, enhances




TWO CELLULAR MECHANISMS are hypothesized for associative changes in synaptic strength during learning. The pre-post coincidence mechanism, proposed by Donald O. Hebb in 1949, posits that coincident activity in the presynaptic and postsynaptic neurons is critical for strengthening the connections between them. The pre-modulatory coincidence mechanism proposed in 1963, based on studies in *Aplysia*, holds that the connection can be strengthened without activity of the postsynaptic cell when a third neuron, the modulatory neuron, is active at the same time as the presynaptic neuron. Stripes denote neurons in which coincident activity must occur to produce the associative change.

Classical Conditioning in *Aplysia*

The marine snail *Aplysia* (top left) is used in studies of the biological basis of learning because its simple nervous system consists of only 20,000 relatively large neurons. The diagram (bottom left) traces one of the pathways involved in classical conditioning of the gill-withdrawal reflex in *Aplysia*. An increase in the release of neurotransmitter due to activity-dependent facilitation is a mechanism that contributes to conditioning. The molecular steps in activity-dependent facilitation are shown in

the enlargement at the right. Serotonin released from the modulatory neuron by the unconditioned stimulus activates adenylyl cyclase in the sensory neuron. When the sensory neuron is active, levels of calcium are elevated within the cell. The calcium binds to calmodulin, which in turn binds to adenylyl cyclase, enhancing its ability to synthesize cyclic AMP. The cyclic AMP activates protein kinase, which leads to the release of a substantially greater amount of transmitter than would occur normally.



transmitter release from the terminals of the presynaptic neuron. They suggested that this mechanism could take on associative properties if the electrical impulses known as action potentials in the presynaptic cell were coincident with action potentials in the modulatory neuron (a pre-modulatory associative mechanism).


Subsequently, we and our colleagues Thomas J. Carew and Thomas W. Abrams of Columbia University and Edgar T. Walters and John H. Byrne of the University of Texas Health Science Center found experimental confirmation. We observed the pre-modulatory asso-

ciative mechanism in *Aplysia*, where it contributes to classical conditioning, an implicit form of learning. Then, in 1986, Holger J. A. Wigström and Bengt E. W. Gustafsson, working at the University of Göteborg, found that the pre-post associative mechanism occurs in the hippocampus, where it is utilized in types of synaptic change that are important for spatial learning, an explicit form of learning.

The finding of two distinct cellular learning rules, each with associative properties, suggested that the associative mechanisms for implicit and explicit learning need not require complex

neural networks. Rather the ability to detect associations may simply reflect the intrinsic capability of certain cellular interactions. Moreover, these findings raised an intriguing question: Are these apparently different mechanisms in any way related? Before considering their possible interrelation, we shall first describe the two learning mechanisms, beginning with the pre-modulatory mechanism contributing to classical conditioning in *Aplysia*.

Classical conditioning was first described at the turn of the century by the Russian physiologist Ivan Pavlov, who immediately appreciated that con-



HIPPOCAMPUS stores long-term memory for weeks and gradually transfers it to specific regions of the cerebral cortex. The diagram illustrates an example of this process involving a visual image. Neural input travels to the visual cortex and then to the hippocampus, where it is stored for several weeks before it is transferred back to the cortex for long-term memory. The hippocampus (*enlargement*) has three major synaptic pathways, each capable of long-term potentiation (LTP), which is thought to play a role in the storage process. LTP has different properties in the CA₁ and CA₃ regions of the hippocampus.

ditioning represents the simplest example of learning to associate two events. In classical conditioning, an ineffective stimulus called the conditioned stimulus (or more correctly, the to-be-conditioned stimulus) is repeatedly paired with a highly effective stimulus called the unconditioned stimulus. The conditioned stimulus initially produces only a small response or no response at all; the unconditioned stimulus elicits a powerful response without requiring prior conditioning.

As a result of conditioning (or learning), the conditioned stimulus becomes capable of producing either a larger response or a completely new response. For example, the sound of a bell (the conditioned stimulus) becomes effective in eliciting a behavioral response such as lifting a leg only after that sound has been paired with a shock to the leg (the unconditioned stimulus) that invariably produces a leg-lifting response. For conditioning to occur, the conditioned stimulus generally must be correlated with the unconditioned stimulus and precede it by a certain critical period. The ani-

mal is therefore thought to learn predictive relations between the two stimuli.

Because *Aplysia* has a nervous system containing only about 20,000 central nerve cells, aspects of classical conditioning can be examined at the cellular level. *Aplysia* has a number of simple reflexes, of which the gill-withdrawal reflex has been particularly well studied. The animal normally withdraws the gill, its respiratory organ, when a stimulus is applied to another part of its body such as the mantle shelf or the fleshy extension called the siphon. Both the mantle shelf and the siphon are innervated by their own populations of sensory neurons. Each of these populations makes direct contact with motor neurons for the gill as well as with various classes of excitatory and inhibitory interneurons that synapse on the motor neurons. We and our colleagues Carew and Walters found that even this simple reflex can be conditioned.

A weak tactile stimulus to one pathway, for example, the siphon, can be paired with an unconditioned stimulus (a strong shock) to the tail. The other

pathway, the mantle shelf, can then be used as a control pathway. The control pathway is stimulated the same number of times, but the stimulus is not paired (associated) with the tail shock. After five pairing trials, the response to stimulation of the siphon (the paired pathway) is greater than that of the mantle (the unpaired pathway). If the procedure is reversed and the mantle shelf is paired rather than the siphon, the response to the mantle shelf will be greater than that to the siphon. This differential conditioning is remarkably similar in several respects to that seen in vertebrates.

To discover how this conditioning works, we focused on one component: the connections between the sensory neurons and their target cells, the interneurons and motor neurons. Stimulating the sensory neurons from either the siphon or the mantle shelf generates excitatory synaptic potentials in the interneurons and motor cells. These synaptic potentials cause the motor cells to discharge, leading to a brisk reflex withdrawal of the gill. The unconditioned reinforcing stimulus to the tail activates many cell groups, some of which also cause movement of the gill. Among them are at least three groups of modulatory neurons, in one of which the chemical serotonin is the transmitter. (Neurotransmitters such as serotonin that carry messages between cells are called first messengers; other chemicals known as second messengers relay information within the cell.)

These modulatory neurons act on the sensory neurons from both the siphon and the mantle shelf, where they produce presynaptic facilitation, that is, they enhance transmitter release from the terminals of the sensory neurons. Presynaptic facilitation contributes to a nonassociative form of learning called sensitization, in which an animal learns to enhance a variety of defensive reflex responses after receiving a noxious stimulus [see "Small Systems of Neurons," by Eric R. Kandel; *SCIENTIFIC AMERICAN*, September 1979]. This type of learning is referred to as nonassociative because it does not depend on pairing between stimuli.

The finding that modulatory neurons act on both sets of sensory neurons—those from the siphon as well as those from the mantle—posed an interesting question: How is the specific associative strengthening of classical conditioning achieved? Timing turned out to be an important element here. For classical conditioning to occur, the conditioned stimulus generally must precede the unconditioned stimulus by a critical and often narrow interval. For condi-

tioning gill withdrawal by tail shock, the interval is approximately 0.5 second. If the separation is lengthened, shortened or reversed, conditioning is drastically reduced or does not occur.

In the gill-withdrawal reflex, the specificity in timing results in part from a convergence of the conditioned and unconditioned stimuli within individual sensory neurons. The unconditioned stimulus is represented in the sensory neurons by the action of the modulatory neurons, in particular the cells in which serotonin is the transmitter. The conditioned stimulus is represented by activity within the sensory neurons themselves. We found that the modulatory neurons activated by the unconditioned stimulus to the tail produce greater presynaptic facilitation of the sensory neurons if the sensory neurons had just fired action potentials in response to the conditioned stimulus. Action potentials in the sensory neurons that occur just after the tail shock have no effect.

This novel property of presynaptic facilitation is called activity dependence. Activity-dependent facilitation requires the same timing on the cellular level as does conditioning on the behavioral level and may account for such conditioning. These results suggest that a cellular mechanism of classical conditioning of the withdrawal reflex is an elaboration of presynaptic facilitation, a mechanism used for sensitization of the reflex. These experiments provided an initial suggestion that there might be a cellular alphabet for learning whereby the mechanisms of more complex types of learning may be elaborations or combinations of the mechanisms of simpler types of learning.


The next piece in the puzzle of how classical conditioning occurs was to discover why the firing of action potentials in the sensory neurons just before the unconditioned tail stimulus would enhance presynaptic facilitation. We had previously found that when serotonin is released by the modulatory neurons in response to tail shock, it initiates a series of biochemical changes in the sensory neurons [see illustration on page 81]. Serotonin binds to a receptor that activates an enzyme called adenylyl cyclase. This enzyme in turn converts ATP, one of the molecules that provides the energy needed to power the various activities of the cell, into cyclic AMP. Cyclic AMP then acts as a second messenger (serotonin is the first messenger) inside the cell to activate another enzyme, a protein kinase. Kinases are proteins that phosphorylate (add a phosphate group to) other proteins, thereby increasing the activity of some

and decreasing the activity of others.

The activation of the protein kinase in sensory neurons has several important short-term consequences. The protein kinase phosphorylates potassium channel proteins. Phosphorylation of these channels (or of proteins that act on these channels) reduces a component of the potassium current that normally repolarizes the action potential. Reduction of potassium current prolongs the action potential and thereby allows calcium channels to be activated for longer periods, permitting more calcium to enter the presynaptic terminal. Calcium has several actions within the cell, one of which is the release of transmitter vesicles from the terminal. When, as a result of an increase in the duration of the action potentials, more calci-

um enters the terminal, more transmitter is released. Second, as a result of protein kinase activity, serotonin acts to mobilize transmitter vesicles from a storage pool to the release sites at the membrane; this facilitates the release of transmitter independent of an increase in calcium influx. In this action, cyclic AMP acts in parallel with another second messenger, protein kinase C, which is also activated by serotonin.

Why should the firing of action potentials in the sensory neurons just before the unconditioned stimulus enhance the action of serotonin? Action potentials produce a number of changes in the sensory neurons. They allow sodium and calcium to move in and potassium to move out, and they change the membrane potential. Abrams and



IN LONG-TERM POTENTIATION the postsynaptic membrane is depolarized by the actions of the non-NMDA receptor channels. The depolarization relieves the magnesium blockade of the NMDA channel, allowing calcium to flow through the channel. The calcium triggers calcium-dependent kinases that lead to the induction of LTP. The postsynaptic cell is thought to release a retrograde messenger capable of penetrating the membrane of the presynaptic cell. This messenger, which may be nitric oxide, is believed to act in the presynaptic terminal to enhance transmitter (glutamate) release, perhaps by activating guanylyl cyclase or ADP-ribosyl transferase.

Kandel found that the critical function of the action potential for activity dependence was the movement of calcium into the sensory neurons. Once in the cell, calcium binds to a protein called calmodulin, which amplifies the activation of the enzyme adenylyl cyclase by serotonin. When calcium/calmodulin binds to the adenylyl cyclase, the enzyme generates more cyclic AMP. This capacity makes adenylyl cyclase an important convergence site for the conditioned and the unconditioned stimuli.

Thus, the conditioned and the unconditioned stimuli are represented within the cell by the convergence of two different signals (calcium and serotonin) on the same enzyme. The 0.5-second interval between the two stimuli essential for learning in the gill-withdrawal reflex may correspond to the time during which calcium is elevated in the presynaptic terminal and binds to calmodulin so as to prime the adenylyl cyclase to produce more cyclic AMP in response to serotonin.

Activity-dependent amplification of the cyclic AMP pathway is not unique to the gill- or tail-withdrawal reflexes of *Aplysia*. Genetic studies in the fruit fly *Drosophila* have implicated a similar molecular mechanism for conditioning. *Drosophila* can be conditioned, and sin-

gle-gene mutants have been discovered that are deficient in learning. One such mutant, called *rutabaga*, has been studied by William G. Quinn of the Massachusetts Institute of Technology and Margaret Livingstone of Harvard University and by Yadin Dudai of the Weizmann Institute in Israel. The gene encoding the defective protein in this mutant has now been shown to be a calcium/calmodulin-dependent adenylyl cyclase. As a result of the mutation in *rutabaga*, the cyclase has lost its ability to be stimulated by calcium/calmodulin. Moreover, Ronald L. Davis and his colleagues at Cold Spring Harbor Laboratory have found that this form of the adenylyl cyclase is enriched in the mushroom bodies, a part of the fly brain critical for several types of associative learning. Thus, both cell biological studies in *Aplysia* and genetic studies in *Drosophila* point to the significance of the cyclic AMP second-messenger system in certain elementary types of implicit learning and memory storage.

What about explicit forms of learning? Do these more complex types of associative learning also have cellular representations for associativity? If so, they must differ from the mechanisms for implicit learning,


because, unlike classical conditioning, explicit learning is often most successful when the two events that are associated occur simultaneously. For example, we recognize the face of an acquaintance most easily when we see that acquaintance in a specific context. The stimuli of the face and of the setting act simultaneously to help us recognize the person.

As we have seen, explicit learning in humans requires the temporal lobe. Yet it was unclear at first how extensive the bilateral lesion in the temporal lobe had to be to interfere with memory storage. Subsequent studies in humans and in experimental animals by Mortimer Mishkin of the National Institutes of Health and by Squire, David G. Amaral and Stuart Zola-Morgan of the University of California at San Diego help to answer the question. They suggest that one structure within the temporal lobe particularly critical for memory storage is the hippocampus. And yet lesions of the hippocampus interfere only with the storage of new memories: patients like H.M. still have a reasonably good memory of earlier events. The hippocampus appears to be only a temporary depository for long-term memory. The hippocampus processes the newly learned information for a period of weeks to months and then transfers the information to relevant areas of the cerebral cortex for more permanent storage [see "Brain and Language," by Antonio R. Damasio and Hanna Damasio, page 88]. As discussed by Patricia S. Goldman-Rakic, the memory stored at these different cortical sites is then expressed through the working memory of the prefrontal cortex [see "Working Memory and the Mind," page 110].


In 1973 Timothy Bliss and Terje Lømo, working in Per Andersen's laboratory in Oslo, Norway, first demonstrated that neurons in the hippocampus have remarkable plastic capabilities of the kind that would be required for learning. They found that a brief high-frequency train of action potentials in one of the neural pathways within the hippocampus produces an increase in synaptic strength in that pathway. The increase can be shown to last for hours in an anesthetized animal and for days and even weeks in an alert, freely moving animal.

Bliss and Lømo called this strengthening long-term potentiation (LTP). Later studies showed that LTP has different properties in different types of synapses within the hippocampus. We will focus here on an associative type of potentiation that has two interrelated characteristics. First, the associativity is of the Hebbian pre-post form: for facilitation to occur, the contributing presyn-

ACTIVITY-DEPENDENT FACILITATION IN APLYSIA



LONG-TERM POTENTIATION IN THE HIPPOCAMPUS



ASSOCIATIVE PROCESSES believed to contribute to learning in *Aplysia* and in the hippocampus of mammals may share similar mechanisms. Both may involve a modulatory substance that produces activity-dependent enhancement of transmitter release from the presynaptic neuron. Stripes denote neurons in which coincident activity must occur to produce the associative change.

aptic and postsynaptic neurons need to be active simultaneously. Second, and as a result, the long-term potentiation shows specificity: it is restricted in its action to the pathway that is stimulated.


Why is simultaneous firing of the presynaptic and postsynaptic cells necessary for long-term potentiation? The major neural pathways in the hippocampus use the amino acid glutamate as their transmitter. Glutamate produces LTP by binding to glutamate receptors on its target cells. It turns out that there are two relevant kinds of glutamate receptors: the NMDA receptors (named after the chemical *N*-methyl *D*-aspartate, which also binds to these receptors) and the non-NMDA receptors. Non-NMDA receptors dominate most synaptic transmission because the ion channel associated with the NMDA receptor is usually blocked by magnesium. It becomes unblocked only when the postsynaptic cell is depolarized. Moreover, optimal activation of the NMDA receptor channel requires that the two signals—glutamate binding to the receptor and depolarization of the postsynaptic cell—take place simultaneously. Thus, the NMDA receptor has associative or coincidence-detecting properties much as does the adenylyl cyclase. But its temporal characteristics, a requirement for simultaneous activation, are better suited for explicit rather than implicit forms of learning.

Calcium influx into the postsynaptic cell through the unblocked NMDA receptor channel is critical for long-term potentiation, as was first shown by Gary Lynch of the University of California at Irvine and by Roger A. Nicoll and Robert S. Zucker and colleagues at the University of California at San Francisco. Calcium initiates LTP by activating at least three different types of protein kinases.

The *induction* of LTP appears to depend on postsynaptic depolarization, leading to the influx of calcium and the subsequent activation of second-messenger kinases. For the *maintenance* of LTP, on the other hand, several groups of researchers have found that enhancement of transmitter from the presynaptic terminal is involved. These workers include Bliss and his colleagues, John Bekkers and Charles Stevens of the Salk Institute and Roberto Malinow and Richard Tsien of Stanford University.

If the induction of LTP requires a postsynaptic event (calcium influx through the NMDA receptor channels) and maintenance of LTP involves a presynaptic event (increase in transmitter release), then, as first proposed by Bliss, some message must be sent from the postsynaptic to the presynaptic neurons—and that poses a problem for neuroscien-


Representation of the Surface of the Body in the Cortex



The homunculus ("little man") is a traditional way of illustrating how the surface of the body is represented in the somatosensory cortex. Larger areas of the cortex are devoted to parts of the body that have greater sensitivity, such as the fingers and lips.

Recently the effects of sensitivity training have been shown in the owl monkey. The monkey's digits are represented in areas 3b and 1 of

the somatosensory cortex (*a*). The diagrams (*b* and *d*) outline the regions that map the surface of the digits of an adult monkey (*c*) before and after training. During training the monkey rotated a disk for one hour a day, using only digits 2, 3 and occasionally 4. After three months of this activity, the area representing the stimulated fingers in the brain had increased substantially.



tists. Ever since the great Spanish anatomist Santiago Ramón y Cajal first enunciated the principle of dynamic polarization, every chemical synapse studied has proved to be unidirectional. Information flows only from the presynaptic to the postsynaptic cell. In long-term potentiation, a new principle of nerve cell communication seems to be emerging. The calcium-activated second-messenger pathways, or perhaps calcium acting directly, seem to cause release of a retrograde plasticity factor from the active postsynaptic cell. This retrograde factor then diffuses to the presynaptic terminals to activate one or more second messengers that enhance transmitter release and thereby maintain LTP [see illustration on page 83].

Unlike the presynaptic terminals, which store transmitter in vesicles and release it at specialized release sites, the postsynaptic terminals lack any special release machinery. It therefore seemed attractive to posit that the retrograde messenger may be a substance that rapidly diffuses out of the postsynaptic cell across the synaptic cleft and into the presynaptic terminal. By 1991 four groups of researchers had obtained evi-

dence that nitric oxide may be such a retrograde messenger: Thomas J. O'Dell and Ottavio Arancio in our laboratory, Erin M. Schuman and Daniel Madison of Stanford University, Paul F. Chapman and his colleagues at the University of Minnesota School of Medicine and Georg Böhme and his colleagues in France. Inhibiting the synthesis of nitric oxide in the postsynaptic neuron or absorbing nitric oxide in the extracellular space blocks the induction of LTP, whereas applying nitric oxide enhances transmitter release from presynaptic neurons.

In the course of studying the effects of applying nitric oxide in slices of hippocampus, we and Scott A. Small and Min Zhuo made a surprising finding: we discovered that nitric oxide produces LTP only if it is paired with activity in the presynaptic neurons, much as is the case in activity-dependent presynaptic facilitation in *Aplysia*. Presynaptic activity, and perhaps calcium influx, appears to be critical for nitric oxide to produce potentiation. These experiments suggest that long-term potentiation uses a combination of two independent, associative, synaptic learning mechanisms: a Hebbian NMDA receptor mechanism and a

non-Hebbian, activity-dependent, presynaptic facilitating mechanism. According to this hypothesis, the activation of NMDA receptors in the postsynaptic cells produces a retrograde signal (nitric oxide). The signal then initiates an activity-dependent presynaptic mechanism, which facilitates the release of transmitter from the presynaptic terminals.

What might be the functional advantage of combining two associative cellular mechanisms, the postsynaptic NMDA receptor and the activity-dependent presynaptic facilitation, in this way? If presynaptic facilitation is produced by a diffusible substance, that substance could, in theory, find its way into neighboring pathways. In fact, studies by Tobias Bonhoeffer and his colleagues at the Max Planck Institute for Brain Research in Frankfurt indicate that LTP initiated in one postsynaptic cell spreads to neighboring postsynaptic cells. Activity dependence of presynaptic facilitation could be a way of ensuring that only specific presynaptic pathways—those that are active—are potentiated. Any inactive presynaptic terminals would not be affected [see illustration on page 84].

The changes in synapses that are thought to contribute to these instances of implicit and explicit learning raise a surprising reductionist possibility. The fact that associative synaptic changes do not require complex neural networks suggests there may be a direct correspondence between these associative forms of learning and basic cellular properties. In the cases that we have reviewed, the cellular properties seem to derive in turn from the properties of specific proteins—the adenylyl cyclase and the NMDA receptor—that are capable of responding to two independent signals, such as those from the conditioned stimulus and the unconditioned stimulus. Of course, these molecular associative mechanisms do not act in isolation. They are embedded in cells that have rich molecular machinery for elaborating the associative process. And the cells, in turn, are embedded in complex neural networks with considerable redundancy, parallelism and computational power, adding substantial complexity to these elementary mechanisms.

The finding that LTP occurs in the hippocampus, a region known to be significant in memory storage, made researchers wonder whether LTP is involved in the process of storing memories in this area of the brain. Evidence that it is has been provided by Richard Morris and his colleagues at the University of Edinburgh Medical School by means of a spatial memory task. When NMDA receptors in the hippocampus are blocked, the experimental animals

fail to learn the task. These experiments suggest that NMDA receptor mechanisms in the hippocampus, and perhaps LTP, are involved in spatial learning.

Having now considered the mechanisms through which learning can produce changes in nerve cells, we are faced with a final set of questions. What are the mechanisms whereby the synaptic changes produced by explicit and implicit learning endure? How is memory maintained in the long term?

Experiments in both *Aplysia* and mammals indicate that explicit and implicit memory storage proceed in stages. Storage of the initial information, a type of short-term memory, lasts minutes to hours and involves changes in the strength of existing synaptic connections (by means of second-messenger-mediated modifications of the kind we have discussed). The long-term changes (those that persist for weeks and months) are stored at the same site, but they require something entirely new: the activation of genes, the expression of new proteins and the growth of new connections. In *Aplysia*, Craig H. Bailey, Mary C. Chen and Samuel M. Schacher and their colleagues at Columbia University and Byrne and his colleagues at the University of Texas Health Science Center have found that stimuli that produce long-term memory for sensitization and classical conditioning lead to an increase in the number of presynaptic terminals. Similar anatomic changes occur in the hippocampus after LTP.

If long-term memory leads to anatomic changes, does that imply that our brains are constantly changing anatomically as we learn and as we forget? Will we experience changes in our brain's anatomy as a result of reading and remembering this issue of *Scientific American*?

This question has been addressed by many investigators, perhaps most dramatically by Michael Merzenich of the University of California at San Francisco. Merzenich examined the representation of the hand in the sensory area of the cerebral cortex. Until recently, neuroscientists believed this representation was stable throughout life. But Merzenich and his colleagues have now demonstrated that cortical maps are subject to constant modification based on use of the sensory pathways. Since all of us are brought up in somewhat different environments, are exposed to different combinations of stimuli and are likely to exercise our sensory and motor skills in different ways, the architecture of each of our brains will be modified in slightly different ways. This

distinctive modification of brain architecture, along with a unique genetic makeup, contributes to the biological basis for the expression of individuality.

This view is best demonstrated in a study by Merzenich, in which he encouraged a monkey to touch a rotating disk with only the three middle fingers of its hand. After several thousand disk rotations, the area in the cortex devoted to the three middle fingers was expanded at the expense of that devoted to the other fingers [see illustration on preceding page]. Practice, therefore, can lead to changes in the cortical representation of the most active fingers. What mechanisms underlie the changes? Recent evidence indicates that the cortical connections in the somatosensory system are constantly being modified and updated on the basis of correlated activity, using a mechanism that appears similar to that which generates LTP.

Indeed, as we have learned from Carla J. Shatz [see "The Developing Brain," page 60], early results from cell biological studies of development suggest that the mechanisms of learning may carry with them an additional bonus. There is now reason to believe that the fine-tuning of connections during late stages of development may require an activity-dependent associative synaptic mechanism perhaps similar to LTP. If that is also true on the molecular level—if learning shares common molecular mechanisms with aspects of development and growth—the study of learning may help connect cognitive psychology to the molecular biology of the organism more generally. This broad biological unification would accelerate the demystification of mental processes and position their study squarely within the evolutionary framework of biology.

FURTHER READING

- AMNESIA FOLLOWING OPERATION ON THE TEMPORAL LOBES. Brenda Milner in *Amnesia: Clinical, Psychological and Medical Aspects*. Edited by C.W.M. Whitty and O. L. Zangwill. Butterworths, 1966.
- A CELLULAR MECHANISM OF CLASSICAL CONDITIONING IN APYLSIA: ACTIVITY-DEPENDENT AMPLIFICATION OF PRESYNAPTIC FACILITATION. R. D. Hawkins, T. W. Abrams, T. J. Carew and E. R. Kandel in *Science*, Vol. 219, pages 400-405; January 28, 1983.
- THE CURRENT EXCITEMENT IN LONG-TERM POTENTIATION. R. A. Nicoll, J. A. Kauer and R. C. Malenka in *Neuron*, Vol. 1, No. 2, pages 97-103; April 1988.
- MEMORY AND THE HIPPOCAMPUS: A SYNTHESIS FROM FINDINGS WITH RATS, MONKEYS, AND HUMANS. Larry R. Squire in *Psychological Review*, Vol. 99, No. 2, pages 195-231; April 1992.

SCIENTIFIC AMERICAN

Presents the September 1992
single topic issue

Mind and Brain

A scientific journey "inward," exploring
the relationship between mind and brain.

- ❖ Extensively illustrated with color photography, art and graphics.
- ❖ Articles by Francis H.C. Crick of the Salk Institute, Gerald D. Fischbach of Harvard Medical School, Eric R. Kandel of the College of Physicians and Surgeons, Antonio and Hanna Damasio of the University of Iowa, Doreen Kimura of the University of Western Ontario, and many others.
- ❖ Fascinating reading.
- ❖ Excellent classroom teaching aid.

As the 20th century ends, investigators are deriving deep understanding of one of the most exciting subjects in modern science: the relationship between mind and brain.

We sense the world around us. We think. We remember. We use language. What are the biological and physiological foundations of these abilities? Where and how are memories stored, and how do we relate one thought to another? Non-invasive imaging technology, molecular genetics and psychopharmacology are producing answers to these and other questions. The new knowledge stands as great fundamental science in its own right. Like all great discoveries, it promises powerful new technologies; in this instance, promising approaches to treating diseases of the mind and enhancing its development.

**Now available
in bulk for
industry and
education**

**SAVE 20%
on bulk orders
of 50 or more**

☐ Yes, I would like to receive the single
topic issue, *Mind and Brain*.

MBSA9

Name (please print) _____

Company _____

Address _____ Apt. _____

City _____ State _____ Zip _____

Photocopies of this order form are acceptable.


Please send _____ copies @ \$4.95 \$ _____

Add \$1.00 per copy
for postage and handling. \$ _____

Total enclosed \$ _____

Deduct 20% on orders of 50 or more copies. Postage and handling paid by SCIENTIFIC AMERICAN on orders of ten or more copies. Please make check or money order payable to SCIENTIFIC AMERICAN.

Send orders to: SCIENTIFIC AMERICAN, Dept. M&B,
415 Madison Avenue, New York, NY 10017, USA



Brain and Language

*A large set of neural structures serves to represent concepts; a smaller set forms words and sentences.
Between the two lies a crucial layer of mediation*

by Antonio R. Damasio and Hanna Damasio

What do neuroscientists talk about when they talk about language? We talk, it seems, about the ability to use words (or signs, if our language is one of the sign languages of the deaf) and to combine them in sentences so that concepts in our minds can be transmitted to other people. We also consider the converse: how we apprehend words spoken by others and turn them into concepts in our own minds.

Language arose and persisted because it serves as a supremely efficient means of communication, especially for abstract concepts. Try to explain the rise and fall of the communist republics without using a single word. But language also performs what Patricia S. Churchland of the University of California at San Diego aptly calls "cognitive compression." It helps to categorize the world and to reduce the complexity of conceptual structures to a manageable scale.

The word "screwdriver," for example, stands for many representations of such an instrument, including visual descriptions of its operation and purpose, specific instances of its use, the feel of the tool or the hand movement that pertains to it. Or there is the immense variety of conceptual representations denoted by a word such as "democracy." The cognitive economies of language—its facility for pulling together many concepts under one symbol—make it possible for people to establish ever more complex concepts and use them to think at levels that would otherwise be impossible.

In the beginning, however, there were no words. Language seems to have appeared in evolution only after humans and species before them had become adept at generating and categorizing actions and at creating and categorizing mental representations of objects, events and relations. Similarly, infants' brains are busy representing and evoking concepts and generating myriad actions long before they utter their first well-selected word and even longer before they form sentences and truly use language. However, the maturation of language processes may not always depend on the maturation of conceptual processes, since some children with defective conceptual systems have nonetheless acquired grammar. The neural machinery necessary for some syntactic operations seems capable of developing autonomously.

Language exists both as an artifact in the external world—

MARTIN LUTHER KING is remembered both for his vision of racial harmony and for his ability to find words that stirred his listeners to action. The central issue of the neurophysiology of language, the authors say, is to map the structures in the brain that manipulate concepts and those that turn the concepts into words.



a collection of symbols in admissible combinations—and as the embodiment in the brain of those symbols and the principles that determine their combinations. The brain uses the same machinery to represent language that it uses to represent any other entity. As neuroscientists come to understand the neural basis for the brain's representations of external objects, events and their relations, they will simultaneously gain insight into the brain's representation of language and into the mechanisms that connect the two.

We believe the brain processes language by means of three interacting sets of structures. First, a large collection of neural systems in both the right and left cerebral hemispheres represents nonlanguage interactions between the body and its environment, as mediated by varied sensory and motor systems—that is to say, anything that a person does, perceives, thinks or feels while acting in the world.


The brain not only categorizes these nonlanguage representations (along lines such as shape, color, sequence or emotional state), it also creates another level of representation for the results of its classification. In this way, people organize objects, events and relationships. Successive layers of categories and symbolic representations form the basis for abstraction and metaphor.

Second, a smaller number of neural systems, generally located in the left cerebral hemisphere, represent phonemes, phoneme combinations and syntactic rules for combining words. When stimulated from within the brain, these systems assemble word-forms and generate sentences to be spoken or written. When stimulated externally by speech or text, they perform the initial processing of auditory or visual language signals.

A third set of structures, also located largely in the left hemisphere, mediates between the first two. It can take a concept and stimulate the production of word-forms, or it can receive words and cause the brain to evoke the corresponding concepts.

Such mediation structures have also been hypothesized

ANTONIO R. DAMASIO and HANNA DAMASIO have been investigating the neural basis of language and memory for the past two decades. Antonio Damasio is professor and head of the department of neurology at the University of Iowa College of Medicine and adjunct professor at the Salk Institute for Biological Studies in San Diego. He received his M.D. and doctorate from the University of Lisbon. Hanna Damasio also holds an M.D. from the University of Lisbon. She is professor of neurology and director of the Laboratory for Neuroimaging and Human Neuroanatomy at the University of Iowa.



BRAIN SYSTEMS FOR COLOR exemplify the organization of language structures. Evidence from studies of people who have sustained brain damage indicates that the concepts for colors depend on the functioning of one system, that words for colors depend on another system and that the connections between words and concepts depend on a third.

from a purely psycholinguistic perspective. Willem J. M. Levelt of the Max Planck Institute for Psycholinguistics in Nijmegen has suggested that word-forms and sentences are generated from concepts by means of a component he calls "lemma," and Merrill F.

Garret of the University of Arizona holds a similar view.

The concepts and words for colors serve as a particularly good example of this tripartite organization. Even those afflicted by congenital color blindness know that certain ranges of hue (chro-

ma) band together and are different from other ranges, independent of their brightness and saturation. As Brent Berlin and Eleanor H. Rosch of the University of California at Berkeley have shown, these color concepts are fairly universal and develop whether or not a given culture actually has names to denote them. Naturally, the retina and the lateral geniculate nucleus perform the initial processing of color signals, but the primary visual cortex and at least two other cortical regions (known as V2 and V4) also participate in color processing; they fabricate what we know as the experience of color.

With our colleague Matthew Rizzo, we have found that damage to the occipital and subcalcarine portions of the left and right lingual gyri, the region of the brain believed to contain the V2 and V4 cortices, causes a condition called achromatopsia. Patients who previously had normal vision lose their perception of color [see "The Visual Image in Mind and Brain," by Semir Zeki, page 68]. Furthermore, they lose the ability even to imagine colors. Achromatopsics usually see the world in shades of gray; when they conjure up a typically colored image in their minds, they see the shapes, movement and texture but not the color. When they think about a field of grass, no green is available, nor will red or yellow be part of their otherwise normal evocation of blood or banana. No lesion elsewhere in the brain can cause a similar defect. In some sense, then, the concept of colors depends on this region.

Patients with lesions in the left posterior temporal and inferior parietal cortex do not lose access to their concepts, but they have a sweeping impairment of their ability to produce proper word morphology regardless of the category to which a word belongs. Even if they are properly experiencing a given color and attempting to retrieve the corresponding word-form, they produce phonemically distorted color names; they may say "buh" for "blue," for example.

Other patients, who sustain damage in the temporal segment of the left lingual gyrus, suffer from a peculiar defect called color anomia, which affects neither color concepts nor the utterance of color words. These patients continue to *experience* color normally: they can match different hues, correctly rank hues of different saturation and easily put the correct colored paint chip next to objects in a black-and-white photograph. But their ability to put names to color is dismally impaired. Given the limited set of color names available to

Components of a Sound-Based Language

PHONEMES	The individual sound units, whose concatenation, in a particular order, produces morphemes.
MORPHEMES	The smallest meaningful units of a word, whose combination creates a word. (In sign languages the equivalent of a morpheme is a visuomotor sign.)
SYNTAX	The admissible combinations of words in phrases and sentences (called grammar, in popular usage).
LEXICON	The collection of all words in a given language. Each lexical entry includes all information with morphological or syntactic ramifications but does not include conceptual knowledge.
SEMANTICS	The meanings that correspond to all lexical items and to all possible sentences.
PROSODY	The vocal intonation that can modify the literal meaning of words and sentences.
DISCOURSE	The linking of sentences such that they constitute a narrative.

those of us who are not interior decorators, it is surprising to see patients use the word "blue" or "red" when shown green or yellow and yet be capable of neatly placing a green chip next to a picture of grass or a yellow chip next to a picture of a banana. The defect goes both ways: given a color name, the patient will point to the wrong color.

At the same time, however, all the wrong color names the patient uses are beautifully formed, phonologically speaking, and the patient has no other language impairment. The color-concept system is intact, and so is the word-form implementation system. The problem seems to reside with the neural system that mediates between the two.

The same three-part organization that explains how people manage to talk about color applies to other concepts as well. But how are such concepts physically represented in the brain? We believe there are no permanently held "pictorial" representations of objects or persons as was traditionally thought. Instead the brain holds, in effect, a record of the neural activity that takes place in the sensory and motor cortices during interaction with a given object. The records are patterns of synaptic connections that can re-create the separate sets of activity that define an object or event; each record can also stimulate related ones. For example, as a person picks up a coffee cup, her visual cortices will respond to the colors of the cup and of its contents as well as to its shape and position. The somatosensory cortices will register the shape the hand assumes as it holds the cup, the movement of the hand and the arm as they bring the cup to the lips, the warmth of the coffee, and the body change some people call pleasure when they drink the stuff. The brain does not merely represent aspects of external reality; it also records how the body explores the world and reacts to it.

The neural processes that describe the interaction between the individual and the object constitute a rapid sequence of micropereceptions and micro-actions, almost simultaneous as far as consciousness is concerned. They occur in separate functional regions, and each region contains additional subdivisions: the visual aspect of perception, for example, is segregated within smaller systems specialized for color, shape and movement.

Where can the records that bind together all these fragmented activities be held? We believe they are embodied in ensembles of neurons within the


brain's many "convergence" regions. At these sites the axons of feedforward projecting neurons from one part of the brain converge and join with reciprocally diverging feedback projections from other regions. When reactivation within the convergence zones stimulates the feedback projections, many anatomically separate and widely distributed neuron ensembles fire simultaneously and reconstruct previous patterns of mental activity.

In addition to storing information about experiences with objects, the brain also categorizes the information so that related events and concepts—shapes, colors, trajectories in space and time, and pertinent body movements and reactions—can be reactivated together. Such categorizations are denoted by yet another record in another convergence zone. The essential properties of the entities and processes in any interaction are thus represented in an interwoven fashion. The collected knowledge that can be represented includes the fact that a coffee cup has dimensions and a boundary; that it is made of something and has parts; that if it is divided it no longer is a cup, unlike water, which retains its identity no matter how it is divided; that it moved along a particular path, starting at one point in space and ending at another; that arrival at its destination produced a specific outcome. These aspects of neu-


ral representation bear a strong resemblance to the primitives of conceptual structure proposed by Ray Jackendoff of Brandeis University and the cognitive semantic schemas hypothesized by George P. Lakoff of the University of California at Berkeley, both working from purely linguistic grounds.

Activity in such a network, then, can serve both understanding and expression. The activity in the network can reconstruct knowledge so that a person experiences it consciously, or it can activate a system that mediates between concept and language, causing appropriately correlated word-forms and syntactical structures to be generated. Because the brain categorizes perceptions and actions simultaneously along many different dimensions, symbolic representations such as metaphor can easily emerge from this architecture.

Damage to parts of the brain that participate in these neural patterns should produce cognitive defects that clearly delineate the categories according to which concepts are stored and retrieved (the damage that results in achromatopsia is but one example of many). Elizabeth K. Warrington of the National Hospital for Nervous Diseases in London has studied category-related recognition defects and found patients who lose cognizance of certain classes of object. Similarly, in collabor-



LANGUAGE ACTIVITY is visible in this positron emission tomographic (PET) scan of a normal individual performing a naming task. The PET image has been projected onto a three-dimensional magnetic resonance imaging (MRI) reconstruction of the same individual's brain. There are many areas of increased activity in the left hemisphere, including the motor cortex and the anterior and posterior language sectors (arrows). The image was produced by the University of Iowa's Department of Neurology, PET Facility and Image Analysis Facility.



BRAIN SYSTEMS FOR LANGUAGE in the left hemisphere include word and sentence-implementation structures and mediation structures for various lexical items and grammar. The collections of neural structures that represent the concepts themselves are distributed across both right and left hemispheres in many sensory and motor regions.

ration with our colleague Daniel Tranel, we have shown that access to concepts in a number of domains depends on particular neural systems.

For example, one of our patients, known as Boswell, no longer retrieves concepts for any unique entity (a specific person, place or event) with which he was previously familiar. He has also lost concepts for nonunique entities of particular classes. Many animals, for instance, are completely strange to him even though he retains the concept level that lets him know that they are living and animate. Faced with a picture of a raccoon, he says, "It is an animal," but he has no idea of its size, habitat or typical behavior.

Curiously, when it comes to other classes of nonunique entities, Boswell's cognition is apparently unimpaired. He can recognize and name objects, such as a wrench, that are manipulable and have a specific action attached to them. He can retrieve concepts for attributes of entities: he knows what it means for an object to be beautiful or ugly. He can grasp the idea of states or activities such as being in love, jumping or swimming. And he can understand abstract relations among entities or events such as "above," "under," "into," "from,"

"before," "after" or "during." In brief, Boswell has an impairment of concepts for many entities, all of which are denoted by nouns (common and proper). He has no problem whatsoever with concepts for attributes, states, activities and relations that are linguistically signified by adjectives, verbs, functors (prepositions, conjunctions and other verbal connective tissue) and syntactic structures. Indeed, the syntax of his sentences is impeccable.

Lesions such as Boswell's, in the anterior and middle regions of both temporal lobes, impair the brain's conceptual system. Injuries to the left hemisphere in the vicinity of the sylvian fissure, in contrast, interfere with the proper formation of words and sentences. This brain system is the most thoroughly investigated of those involved in language. More than a century and a half ago Paul Broca and Carl Wernicke determined the rough location of these basic language centers and discovered the phenomenon known as cerebral dominance—in most humans language structures lie in the left hemisphere rather than the right. This disposition holds for roughly 99 percent of right-handed people and two thirds of left-

handers. (The pace of research in this area has accelerated during the past two decades, thanks in large part to the influence of the late Norman Geschwind of Harvard Medical School and Harold Goodglass of the Boston Veterans Administration Medical Center.)

Studies of aphasic patients (those who have lost part or all of their ability to speak) from different language backgrounds highlight the constancy of these structures. Indeed, Edward Klima of the University of California at San Diego and Ursula Bellugi of the Salk Institute for Biological Studies in San Diego have discovered that damage to the brain's word-formation systems is implicated in sign-language aphasia as well. Deaf individuals who suffer focal brain damage in the left hemisphere can lose the ability to sign or to understand sign language. Because the damage in question is not to the visual cortex, the ability to see signs is not in question, just the ability to interpret them.

In contrast, deaf people whose lesions lie in the right hemisphere, far from the regions responsible for word and sentence formation, may lose conscious awareness of objects on the left side of their visual field, or they may be unable to perceive correctly spatial relations among objects, but they do not lose the ability to sign or understand sign language. Thus, regardless of the sensory channel through which linguistic information passes, the left hemisphere is the base for linguistic implementation and mediation systems.

Investigators have mapped the neural systems most directly involved in word and sentence formation by studying the location of lesions in aphasic patients. In addition, George A. Ojemann of the University of Washington and Ronald P. Lesser and Barry Gordon of Johns Hopkins University have stimulated the exposed cerebral cortex of patients undergoing surgery for epilepsy and made direct electrophysiological recordings of the response.

Damage in the posterior perisylvian sector, for example, disrupts the assembly of phonemes into words and the selection of entire word-forms. Patients with such damage may fail to speak certain words, or they may form words improperly ("loliphant" for "elephant"). They may, in addition, substitute a pronoun or a word at a more general taxonomic level for a missing one ("people" for "woman") or use a word semantically related to the concept they intend to express ("headman" for "president"). Victoria A. Fromkin of the University of California at Los Angeles has elucidated many of the linguistics

If you're looking
for a notebook
computer that's
easy to use,
there are basically
only two ways
you can go.

The hard way.



New programs, \$506 each.

Unless you buy programs specifically designed for Windows 3.1, you'll have to use conventional character-based programs and their awkward commands.


Extra memory, \$210* Most notebook computers in use today simply aren't fast enough to make Windows 3.1 practical. And many that are fast enough may require extra memory, which can add hundreds of dollars to their price tags.



Clip-on trackball, \$172. If you're going to try to use Windows 3.1 on many portable computers, you may have to plug this in and clip it onto the side of your computer. (Be careful not to catch the cord in your airplane tray.)




Added weight. Many notebook computers may claim to weigh less than 7 pounds, but by the time you add all the components you need to make Windows 3.1 work well, you're likely to find yourself with an additional 4 or 5 pounds to lug around.



Fax/data modem, \$142.

The ability to communicate on the road is what makes a notebook PC really valuable. So you'll have to pack one of these. Once you get back to the office, be prepared to shell out another \$138 to update the files on your desktop PC, and over \$500 just to access network file servers and printers.



Your time, name your price. Fiddling around with MS-DOS or Windows 3.1 on the road can cost you plenty of time. How much is yours worth?



Ordinary notebook, \$2,497. The typical notebook computer of today was designed to run DOS only, not Windows 3.1. It doesn't have enough memory, speed or screen brightness, and its screen isn't fast enough for high-quality graphical computing.

Comfort is something most notebook computer ads never talk about. Because, truth is, they're not very comfortable to use. Which may explain why they don't win awards for being well designed. (PowerBook, of course, has won more design awards than all other notebook computers combined.)



Instruction manual, 2 pounds. There's nothing like forgetting a DOS command when you're 2,000 miles from home, so you may want to carry a manual with you. And since Windows 3.1 is based on DOS, you'll definitely want instructions handy.



The coach seat factor. If you want to run Windows 3.1 on most portable computers, you'll find that you need a front-mounted keyboard and a side-mounted trackball or mouse. What you won't find is enough room on an airplane tray to hold it all.

The Apple® Macintosh® PowerBook™ is the only personal computer to make the Products of the Year lists of *Business Week*, *Fortune*® and *InfoWorld*. And the only notebook computer to be named one of the Best Designed Products of the Year by *Time*®. Because it's the only computer designed as

The easy way.

Consistent applications. Every PowerBook runs thousands of Macintosh programs (including Lotus 1-2-3 and WordPerfect) that all work in the same consistent, intuitive way. You learn one, you know the basics of all of them.

Blistering speed. Because Macintosh computers are optimized for graphical computing, PowerBook can significantly outperform comparably priced notebook computers running Windows.** Both the PowerBook 170 and the new PowerBook 145 deliver an impressive combination of speed and power at an affordable price.

Built-in Balloon Help. Point at what you want to know about, and your Macintosh will tell you what it's for. So you can leave your manual in a desk drawer at home.

Built-in networking. Plug your PowerBook into an AppleTalk® network, and conveniently access all network resources—printers, file servers, e-mail, even other networks from Novell, IBM and Digital.

Built-in fax/data modem. With this option, you can send your work directly to a fax machine or access e-mail from almost any telephone anywhere you happen to be. Just plug the phone cord into the back of the PowerBook.

Built-in peripheral support. Like every Macintosh, PowerBook lets you add up to six external hard drives, CD-ROM drives, scanners or other peripherals simply by plugging them in.

AppleTalk Remote Access. This remarkable software comes with every Macintosh PowerBook. It not only allows you to access all the files on a modem-equipped Macintosh at your office or at home, it also gives you complete access to your network—printers, files, file servers, e-mail—from anywhere there's a phone.

Built-in file sharing. On most notebook computers, sharing information with a desktop computer is virtually impossible. But every PowerBook has powerful file sharing built in. So you can easily connect your PowerBook to a desktop Mac® or another PowerBook and copy files from one computer to the other.

Palm rests. Working on the road isn't like working at your desk. So Apple designers provided a place to rest your hands, making it much easier and more comfortable to work in tight quarters—such as a middle seat on an airplane.

Integrated trackball. Everything about Macintosh computers is designed to make it easy for you to work. In the case of PowerBook, the pointing device is integrated into the computer. There's nothing to hook up or clip on. Open it up and you're ready to go. (It's comfortable whether you're left- or right-handed, too.)

It's compatible. Like every Macintosh, PowerBook is compatible with the personal computers you use at the office. It can read from and write to floppy disks from MS-DOS or OS/2 PCs, or even run DOS software using programs such as SoftPC.



a totally integrated system. Combining hardware, software and industrial design to make working on the road as comfortable as working at your desk. It's more intelligent. It's more convenient. It's power anyone can use, anywhere you happen to be. The power to be your best®.



How to connect to a network file server.

The hard way:

1. Purchase an external network adapter.
2. Install the network adapter.
3. Connect the network cable.
4. Plug in AC power supply if needed.
5. Install network software and configure parameters.
6. Start Windows.
7. Run Windows Setup to load network software.
8. Restart.
9. Start Windows.
10. Start File Manager.
11. Select the desired file server.

The easy way:

1. Connect the network cable.
2. Open Chooser.
3. Select AppleTalk "Active" button.
4. Select desired file server.

How to retrieve a deleted file.

The hard way:

1. Purchase a utility program.
2. Install it on your computer.
3. Restart the computer.
4. Start Windows.
5. Start the utility program.
6. Start the unerase feature.
7. Find the directory where your file existed.
8. Click on that directory.
9. Select the deleted file.
10. Choose unerase command.

The easy way:

1. Double-click on the trash can.
2. Drag the deleted file out of the trash can.

How to rename a directory.

The hard way:

1. Double-click on the Main group in Program Manager.
2. Start File Manager.
3. Highlight the directory you want to rename.
4. Select "Rename" from the file menu.
5. Type in a new name for the directory (must be 8 characters or less).
6. Go to Program Manager if needed.
7. For each application in the renamed directory, select "Properties" from the file menu.
8. For each application in that directory, change the command line to reflect the new path to the application program.

The easy way:

1. Highlight the directory (folder) you want to rename.
2. Type a new name (up to 31 characters).

How to add an external hard disk drive.

The hard way:

1. Purchase a parallel-to-SCSI adapter.
2. Connect it to the parallel port.
3. Plug the drive into the adapter.
4. Insert the adapter's installation disk in drive A:.
5. Select drive A:.
6. Type "INSTALL".
7. Select appropriate options for your drive.
8. Select appropriate drive for software installation.
9. Restart.
10. Run installation test program.

The easy way:

1. Plug the drive into the SCSI port.
2. Start up the PowerBook.

How to play a sound.

The hard way:

1. Purchase a sound upgrade kit.
2. Attach sound upgrade hardware.
3. Install sound upgrade's software drivers.
4. Start the sound recorder program.
5. Select Open from the file menu.
6. Select the sound file you want to play.
7. Choose OK.
8. Click on the Play button.

The easy way:

1. Double-click on the sound's icon.

How to add a pointing device.

The hard way:

1. Purchase a pointing device (such as a mouse or trackball).
2. Open notebook's back panel.
3. Attach appropriate port adapter to pointing device (if other than 9-pin).
4. Connect pointing device to proper port.
5. Choose appropriate clamp arms.
6. Attach pointing device to clamp arms and adjust tilt angle.
7. Start the computer.
8. Install driver software.
9. Restart.

The easy way:

1. The trackball is built in.

Macintosh. The more you do, the easier it gets.

For an authorized Apple reseller, call 800-538-9696, ext. 160. *This price and all the prices noted here reflect average Manufacturer's Suggested Retail Prices. **Comparisons are based on a 1991 independent research study conducted by Ingram Laboratories that tested a variety of personal computers running applications available for both the Macintosh and Microsoft Windows 3.0 environments. ©1992 Apple Computer, Inc. All rights reserved. Apple, the Apple logo, AppleTalk, Mac, Macintosh and "The power to be your best" are registered trademarks of Apple Computer, Inc. Balloon Help and PowerBook are trademarks of Apple Computer, Inc. Lotus and 1-2-3 are registered trademarks of Lotus Development Corporation. MS-DOS is a registered trademark and Windows is a trademark of Microsoft Corporation. OS/2 is a registered trademark of International Business Machines Corporation. SoftPC is a registered trademark of Insignia Solutions Inc. WordPerfect is a registered trademark of WordPerfect Corporation. L0279LL/A

tic mechanisms underlying such errors.

Damage to this region, however, does not disrupt patients' speech rhythms or the rate at which they speak. The syntactic structure of sentences is undisturbed even when there are errors in the use of functor words such as pronouns and conjunctions.

Damage to this region also impairs processing of speech sounds, and so patients have difficulty understanding spoken words and sentences. Auditory comprehension fails not because, as has been traditionally believed, the posterior perisylvian sector is a center to store "meanings" of words but rather because the normal acoustic analyses of the word-forms the patient hears are aborted at an early stage.

The systems in this sector hold auditory and kinesthetic records of phonemes and the phoneme sequences that make up words. Reciprocal projections of neurons between the areas holding these records mean that activity in one can generate corresponding activity in the other.

These regions connect to the motor and premotor cortices, both directly and by means of a subcortical path that includes the left basal ganglia and nuclei in the forward portion of the left thalamus. This dual motor route is especially important: the actual production of speech sounds can take place under the control of either a cortical or a subcortical circuit, or both. The subcortical circuit corresponds to "habit learning," whereas the cortical route implies higher-level, more conscious control and "associative learning" [see "The Biological Basis of Learning and Individuality," by Eric R. Kandel and Robert D. Hawkins, page 78].

For instance, when a child learns the word-form "yellow," activations would pass through the word-formation and motor-control systems via both the cortical and subcortical routes, and activity in these areas would be correlated with the activity of the brain regions responsible for color concepts and mediation between concept and language. In time, we suspect, the concept-mediation system develops a direct route to the basal ganglia, and so the posterior perisylvian sector does not have to be strongly activated to produce the word "yellow." Subsequent learning of the word-form for yellow in another language would again require participation of the perisylvian region to establish auditory, kinesthetic and motor correspondences of phonemes.

It is likely that both cortical "associative" and subcortical "habit" systems

operate in parallel during language processing. One system or the other predominates depending on the history of language acquisition and the nature of the item. Steven Pinker of the Massachusetts Institute of Technology has suggested, for example, that most people acquire the past tense of irregular verbs (take, took, taken) by associative learning and that of regular verbs (those whose past tense ends in *-ed*) by habit learning.

The anterior perisylvian sector, on the front side of the rolandic fissure, appears to contain structures that are responsible for speech rhythms and grammar. The left basal ganglia are part and parcel of this sector, as they are of the posterior perisylvian one. The entire sector appears to be strongly associated with the cerebellum; both the basal ganglia and the cerebellum receive projections from a wide variety of sensory regions in the cortex and return projections to motor-related areas. The role of the cerebellum in language and cognition, however, remains to be elucidated.

Patients with damage in the anterior perisylvian sector speak in flat tones, with long pauses between words, and have defective grammar. They tend in particular to leave out conjunctions and pronouns, and grammatical order is often compromised. Nouns come easier to patients with these lesions than do verbs, suggesting that other regions are responsible for their production.

Patients with damage in this sector have difficulty understanding meaning that is conveyed by syntactic structures. Edgar B. Zurif of Brandeis University, Eleanor M. Saffran of Temple University and Myrna F. Schwartz of Moss Rehabilitation Hospital in Philadelphia have shown that these patients do not always grasp reversible passive sentences such as "The boy was kissed by the girl," in which boy and girl are equally likely to be the recipient of the action. Nevertheless, they can still assign the correct meaning to a nonreversible passive sentence such as "The apple was eaten by the boy" or the active sentence "The boy kissed the girl."

The fact that damage to this sector impairs grammatical processing in both speech and understanding suggests that its neural systems supply the mechanics of component assembly at sentence level. The basal ganglia serve to assemble the components of complex motions into a smooth whole, and it seems reasonable that they might perform an analogous function in assembling word-forms into sentences. We also believe (based on experimental evidence of sim-

ilar, although less extensive structures in monkeys) that these neural structures are closely interconnected with syntactic mediation units in the frontoparietal cortices of both hemispheres [see illustration on page 92]. The delineation of those units is a topic of future research.

Between the brain's concept-processing systems and those that generate words and sentences lie the mediation systems we propose. Evidence for this neural brokerage is beginning to emerge from the study of neurological patients. Mediation systems not only select the correct words to express a particular concept, but they also direct the generation of sentence structures that express relations among concepts.

When a person speaks, these systems govern those responsible for word-formation and syntax; when a person understands speech, the word-formation systems drive the mediation systems. Thus far we have begun to map the systems that mediate proper nouns and common nouns that denote entities of a particular class (for example, visually ambiguous, nonmanipulable entities such as most animals).

Consider the patients whom we will call A.N. and L.R., who had sustained damage to the anterior and midtemporal cortices. Both can retrieve concepts normally: when shown pictures of entities or substances from virtually any conceptual category—human faces, body parts, animals and botanical specimens, vehicles and buildings, tools and utensils—A.N. and L.R. know unequivocally what they are looking at. They can define an entity's functions, habitats and value. If they are given sounds corresponding to those entities or substances (whenever a sound happens to be associated with them), A.N. and L.R. can recognize the item in question. They can perform this task even when they are blindfolded and asked to recognize an object placed in their hands.

But despite their obvious knowledge, they have difficulty in retrieving the names for many of the objects they know so well. Shown a picture of a raccoon, A.N. will say: "Oh! I know what it is—it is a nasty animal. It will come and rummage in your backyard and get into the garbage. The eyes and the rings in the tail give it away. I know it, but I cannot say the name." On the average they come up with less than half of the names they ought to retrieve. Their conceptual systems work well, but A.N. and L.R. cannot reliably ac-

cess the word-forms that denote the objects they know.

The deficit in word-form retrieval depends on the conceptual category of the item that the patients are attempting to name. A.N. and L.R. make fewer errors for nouns that denote tools and utensils than for those naming animals, fruits and vegetables. (This phenomenon has been reported in similar form by Warrington and her colleague Rosaleen A. McCarthy of the National Hospital for Nervous Diseases and by Alfonso Caramazza and his colleagues at Johns Hopkins University.) The patients' ability to find names, however, does not split neatly at the boundary of natural and man-made entities. A.N. and L.R. can produce the words for such natural stimuli as body parts perfectly, whereas they cannot do the same for musical instruments, which are as artificial and as manipulable as garden tools.

In brief, A.N. and L.R. have a problem with the retrieval of common nouns denoting certain entities regardless of their membership in particular conceptual categories. There are many reasons why some entities might be more or less vulnerable to lesions than others. Of necessity, the brain uses different neural systems to represent entities that differ in structure or behavior or entities that a person relates to in different ways.

A.N. and L.R. also have trouble with proper nouns. With few exceptions, they cannot name friends, relatives, celebrities or places. Shown a picture of Marilyn Monroe, A.N. said, "Don't know her name but I know who she is; I saw her movies; she had an affair with the president; she committed suicide; or maybe somebody killed her; the police, maybe?" These patients do not have what is known as face agnosia or prosopagnosia—they can recognize a face without hesitation—but they simply cannot retrieve the word-form that goes with the person they recognize.

Curiously, these patients have no difficulty producing verbs. In experiments we conducted in collaboration with Tranel, these patients perform just as well as matched control subjects on tasks requiring them to generate a verb in response to more than 200 stimuli depicting diverse states and actions. They are also adept at the production of prepositions, conjunctions and pronouns, and their sentences are well formed and grammatical. As they speak or write, they produce a narrative in which, instead of the missing noun, they will substitute words like "thing" or "stuff" or pronouns such as "it" or "she" or

"they." But the verbs that animate the arguments of those sentences are properly selected and produced and properly marked with respect to tense and person. Their pronunciation and prosody (the intonation of the individual words and the entire sentence) are similarly unexceptionable.

The evidence that lexical mediation systems are confined to specific regions is convincing. Indeed, the neural structures that mediate between concepts and word-forms appear to be graded from back to front along the occipitotemporal axis of the brain. Mediation for many general concepts seems to occur at the rear, in the more posterior left temporal regions; mediation for the most specific concepts takes place at the front, near the left temporal pole. We have now seen many patients who have lost their proper nouns but retain all or most of their

common nouns. Their lesions are restricted to the left temporal pole and medial temporal surface of the brain, sparing the lateral and inferior temporal lobes. The last two, in contrast, are always damaged in the patients with common noun retrieval defects.

Patients such as A.N. and L.R., whose damage extends to the anterior and midtemporal cortices, miss many common nouns but still name colors quickly and correctly. These correlations between lesions and linguistic defects indicate that the temporal segment of the left lingual gyrus supports mediation between color concepts and color names, whereas mediation between concepts for unique persons and their corresponding names requires neural structures at the opposite end of the network, in the left anterior temporal lobe. Finally, one of our more recent patients, G.J., has extensive damage that encompasses all of these left occipito-

Components of a Concept

Concepts are stored in the brain in the form of "dormant" records. When these records are reactivated, they can re-create the varied sensations and actions associated with a particular entity or a category of entities. A coffee cup, for example, can evoke visual and tactile representations of its shape, color, texture and warmth, along with the smell and taste of the coffee or the path that the hand and the arm take to bring the cup from the table to the lips. All these representations are re-created in separate brain regions, but their reconstruction occurs fairly simultaneously.



temporal regions from front to back. He has lost access to a sweeping universe of noun word-forms and is equally unable to name colors or unique persons. And yet his concepts are preserved. The results in these patients support Ojemann's finding of impaired language processing after electrical stimulation of cortices outside the classic language areas.

It appears that we have begun to understand fairly well where nouns are mediated, but where are the verbs? Clearly, if patients such as A.N. and L.R. can retrieve verbs and functor words normally, the regions required for those parts of speech cannot be in the left temporal region. Preliminary evidence points to frontal and parietal sites. Aphasia studies performed by our group and by Caramazza and Gabriele Miceli of Catholic University of the Sacred Heart, Milan, and Rita Berndt of the University of Maryland show that

patients with left frontal damage have far more trouble with verb retrieval than with noun retrieval.

Additional indirect evidence comes from positron emission tomography (PET) studies conducted by Steven E. Petersen, Michael I. Posner and Marcus E. Raichle of Washington University. They asked research subjects to generate a verb corresponding to the picture of an object—for example, a picture of an apple might generate "eat." These subjects activated a region of the lateral and inferior dorsal frontal cortex that corresponds roughly to the areas delineated in our studies. Damage to these regions not only compromises access to verbs and functors, it also disturbs the grammatical structure of the sentences that patients produce.

Although this phenomenon may seem surprising at first, verbs and functors constitute the core of syntactic structure, and so it makes sense that the

mediation systems for syntax would overlap with them. Further investigations, either of aphasic patients or of normal subjects, whose brain activity can be mapped by PET scans, may clarify the precise arrangement of these systems and yield maps like those that we have produced to show the differing locations of common and proper nouns.

During the past two decades, progress in understanding the brain structures responsible for language has accelerated significantly. Tools such as magnetic resonance imaging have made it possible to locate brain lesions accurately in patients suffering from aphasia and to correlate specific language deficits with damage to particular regions of the brain. And PET scans offer the opportunity to study the brain activity of normal subjects engaged in linguistic tasks.

Considering the profound complexity of language phenomena, some may wonder whether the neural machinery that allows it all to happen will ever be understood. Many questions remain to be answered about how the brain stores concepts. Mediation systems for parts of speech other than nouns, verbs and functors, have been only partially explored. Even the structures that form words and sentences, which have been under study since the middle of the 19th century, are only sketchily understood.

Nevertheless, given the recent strides that have been made, we believe these structures will eventually be mapped and understood. The question is not if but when.



FURTHER READING

THE SIGNS OF LANGUAGE. Edward S. Klima and Ursula Bellugi. Harvard University Press, 1979.

KNOWLEDGE OF LANGUAGE ITS NATURE, ORIGIN, AND USE. Noam Chomsky. Greenwood Press, 1986.

LESION ANALYSIS IN NEUROPSYCHOLOGY. Hanna Damasio and Antonio R. Damasio. Oxford University Press, 1989.


NEURAL REGIONALIZATION OF KNOWLEDGE ACCESS: PRELIMINARY EVIDENCE. A. R. Damasio, H. Damasio, D. Tranel and J. P. Brandt in *Cold Spring Harbour Symposia on Quantitative Biology*, Vol. LV: *The Brain*. Cold Spring Harbor Laboratory Press, 1990.

APHASIA. A. R. Damasio in *New England Journal of Medicine*, Vol. 326, No. 8, pages 531-539; February 20, 1992.

AN INTRODUCTION TO LANGUAGE. Victoria Fromkin and Robert Rodman. Harcourt Brace Jovanovich College Publications, 1992.

SCIENTIFIC
AMERICAN

DECADE OF THE BRAIN



Brain disorders include well over 650 disorders—ranging from stroke, head and spinal cord injury, to Alzheimer's Disease, schizophrenia, alcoholism and drug addiction—which devastate the lives of an estimated 50 million Americans. Some disorders, like brain tumors and multiple sclerosis, are killers. Others, like learning disabilities and sleep disorders diminish much of the meaning of life. As a category, brain disorders are a leading cause of death in the United States today and they are the most common and severe cause of social, economic, and psychological disability in this country.

The National Foundation for Brain Research (NFBR) is dedicated to the proposition that brain research is the best investment the American people can make to prevent or ameliorate some of the most debilitating disorders of the human condition. To accomplish this goal, the non-profit NFBR has galvanized professional societies, lay organizations, and governmental and corporate bodies to work toward a common objective: the enhancement of brain research.

As the only national organization promoting all aspects of brain research, the National Foundation for Brain Research's ongoing mission is to:

- Improve public awareness and support for scientific research on the brain and its disorders;
- Serve as a clearinghouse for national and international activities concerning the "Decade of the Brain"; and
- Bring together government, medical, and scientific societies, voluntary health agencies, and industry in joint programs of public and professional education.

Ongoing NFBR projects include the development of television series, dissemination of government reports on brain research, and an annual scientific symposium in Washington. One of NFBR's most exciting projects is a science museum exhibit on the brain. The exhibit opened in June at the Franklin Institute Science Museum in Philadelphia and will tour eight cities, with the last stop in October 1994 scheduled for the California Museum of Science & Industry in Los Angeles.


NFBR is widely recognized for its leadership on the Decade of the Brain initiative. In recognition of the importance of neurological and mental research, President George Bush signed into law a Congressional resolution

THE NEW SCIENCE

Two achievements of the human intellect crown the last decade of the 20th Century: Quantum Cosmology, the new science of the Universe, and Neuroscience, the new science of the Brain. The former probes our beginnings, the latter our destiny.

What makes neuroscience "new" and what it can do for humankind are not trivial questions. Neuroscience is not a traditional discipline but a new way of thinking about the brain. Questions such as: How does the brain learn, store memories, recognize one's grandmother, or plan for the future must tap a variety of specialties including cognitive psychology, computational science, neuroimaging, neurophysiology, neurogenetics, molecular biology, and many other disciplines.

What makes neuroscience new is its emergence from a set of principles that are unlikely to be falsified. For example, we now know for certain that nerve impulses result from the flow of different ions through various membrane channels, that communication between neurons occurs via chemical messengers, that neurons and their




Neuroscience is a spectrum of specialties spanning many disciplines.

target cells secrete trophic factors, that brain development is a consequence of genetic and epigenetic interactions, and that neuronal systems can be modified by experience, or lack thereof. As reductionist biology defines the molecular and genetic foundations of these principles, cognitive and systems neuroscience is disclosing constructionist "rules" of brain function.

What neuroscience can do for humankind follows from the foregoing principles and their implications. First, neuroscience can tell scientists how the brain is put together and how it works in health and disease. It also can tell researchers what therapeutic interventions and strategies for prevention can interdict brain dysfunction and degeneration. Finally, neuroscience can help tell us who we are and whether intelligence has survival value for *H. sapiens*.

—Dominick P. Purpura, M.D.,
Chairman, National Foundation for Brain Research



SCHERING-PLOUGH CORP.

Schering-Plough Corp. has recently built on its strengths in chemical synthesis and molecular biology to expand into promising areas of the neurosciences. The Madison, N.J., company has established advanced programs in central nervous system research to develop treatments for Alzheimer's disease and schizophrenia.

"While investing in our biotechnology capabilities in areas such as immunology," says Dr. Frank Bullock, senior vice president for the Schering-Plough Research Institute in Bloomfield, N.J., "we built the infrastructure necessary to expand into other areas. As the neurosciences emerged, we evaluated how emerging areas of science, like cell growth and differentiation and recombinant DNA, could be applied to CNS therapeutics."

One area of investigation involves studies of selective antagonists of dopamine, a brain neurotransmitter. There are now at least five types of dopamine receptors—D1 through D5. Current schizophrenia drugs act by blocking the action of dopamine on D2 receptors. While effectively alleviating the behavioral symptoms of schizophrenia, these drugs cause serious motor abnormalities. Schering-Plough hopes to develop compounds that selectively block D1 receptors in an effort to eliminate the side effects. Using conventional techniques and cloned dopamine receptor sites for screening, researchers are now finishing animal studies and initiating clinical trials.

Schering-Plough has also invested heavily in Alzheimer's therapeutics. One program, which assumes the disease could be due to a deficit in neurotrophic growth factors, seeks to increase the production of naturally occurring growth factor proteins in the brain using small neurotrophic factor-enhancing molecules. A second approach, assuming the disease is caused by inappropriate processing and deposition of beta-amyloid precursor proteins in neurons, seeks to prevent its formation. Both programs employ biotechniques resulting from an agreement between Schering-Plough and Cephalon Inc., West Chester, Pa. A third approach assumes that a deficit in cholinergic neurotransmission characterizes the disease.

Schering-Plough has also entered into a licensing agreement with Carter-Wallace Inc., New York, for overseas marketing rights to felbamate, a promising compound for treating epilepsy.

When the Research Institute moves in December to the Drug Discovery Facility, a 1-million sq.-ft., state-of-the-art facility in Kenilworth, N.J., additional CNS product areas will be considered. Says Dr. Bullock: "The neurosciences and CNS will present some very attractive research opportunities."



A new Drug Discovery Facility, nearing completion in Kenilworth, N.J., will unite most of Schering-Plough's researchers at one site.

designating the 1990s as the "Decade of the Brain." The Decade of the Brain initiative has become an important unifying symbol in the effort to reduce the human suffering as well as the economic impacts of brain disorders. Its passage also has inspired a new age of commitment to neuroscience research by both government and the private sector.

THE COST OF BRAIN DISORDERS

It is impossible to measure the toll that brain disorders take in terms of the suffering of victims and their families. But, in the most comprehensive study of the economic cost of these conditions to date, NFBF recently estimated the economic impact of brain disorders to be as much as \$401.1 billion. Based on a population of 238 million, the U.S. spends roughly \$1,638 per person to combat brain disorders.

This price tag encompasses "direct" and "indirect" costs. The direct costs of medical treatment and services are approximately \$104 billion, or more than \$1 out of every \$7 spent on health care in 1991. The indirect costs include lost wages, opportunity costs associated with family caregiving, or costs of crimes associated with disorders of addiction.

Brain disorders may be divided in four general categories. Psychiatric diseases, which range from schizophrenia to cognitive impairment, account for costs of \$136.1 billion. Neurologic disorders—dementia, mental retardation, multiple sclerosis, head and spinal cord injuries, stroke, cerebral palsy, and epilepsy—cost \$103.7 billion. The cost of alcohol abuse is estimated at \$90.1 billion, and drug abuse at \$71.2 billion. Indirect costs account for more than half to three-quarters of the total cost for each category. (Copies of "The Cost of Disorders of the Brain" are available for \$10 by writing to NFBF, Suite 300, 1250 24th Street, NW, Washington, DC 20037.)

SEIZING THE OPPORTUNITIES

Advances in the neurosciences have significantly raised expectations about our ability to understand the healthy brain and to develop more effective diagnostic techniques and treatments for brain disorders. Our capacity to capitalize on the opportunities that have been created by these advances depends on a strong and continued commitment by both the public and private sectors throughout the 1990s and beyond.

Research operates on a continuum. Basic brain research focuses on understanding the fundamental structure and functions of the brain. In particular, scientists are intent on discovering the fundamental ways in which neurons—the building blocks of the brain—work. At the other end of the spectrum, applied research builds upon the information derived from basic research to develop treatments and diagnostic devices for diseases of the brain.

The federal government is the primary sponsor of basic brain research. Pharmaceutical companies and medical products manufacturers conduct applied research

Neuroscience: Turning expertise into advantage.

Schering-Plough is building on proven strengths in chemical synthesis, biochemistry, pharmacology and biotechnology to pursue new treatments for serious brain disorders.

Building on vast scientific knowledge and advanced technologies, our research scientists are developing compounds that seek to alleviate the symptoms of schizophrenia without the side effects of today's treatments. Pioneering work is also under way to discover novel therapeutics for Alzheimer's disease and age-associated memory impairment.

At Schering-Plough, we are determined to nurture an environment that welcomes new approaches to scientific discovery. Our gifted researchers do far more than discover new products: they continually *re-invent* the organization, allowing for a perpetually innovating company.



Schering-Plough

Schering-Plough Research Institute
2000 Galloping Hill Road
Kenilworth, N.J. 07033

HOECHST-ROUSSEL PHARMACEUTICALS

Hoechst-Roussel Pharmaceuticals Inc. is the center for neurosciences R&D for the Hoechst worldwide organization. The Somerville, N.J., center is dedicated to discovering new compounds for the treatment of Alzheimer's disease, schizophrenia, and obsessive compulsive disorders.

Since 1981, Alzheimer's disease has been a major focus of CNS research. Hoechst-Roussel's three-pronged strategy involves development of drugs to treat the disease's early symptoms (gradual loss of memory and intellectual ability); drugs to treat the advanced symptoms (personality, behavior, and mood changes); and agents to arrest or reverse the disease process.

All three approaches rely on understanding the formation and pathological significance of beta-amyloid plaques, as well as the role of the neurotransmitters acetylcholine and norepinephrine, which are reduced in the brains of Alzheimer's patients. Treatment of symptoms focuses on the role of neurotransmitters, while understanding the role of plaques will more likely lead to the arrest of the disease.

The company has several Alzheimer's compounds in development in the laboratory and in clinical trials. One compound, Mentane™, is in development to treat patients with mild to moderate symptoms. The



Friederike Wirtz-Brugger, Research Pharmacologist for Hoechst-Roussel Pharmaceuticals Inc., maps brain waves for research on Alzheimer's Disease.

drug works by inhibiting the breakdown of acetylcholine, critical to cognitive functioning in the brain.

In the search for better schizophrenia treatments, Hoechst-Roussel is looking at drugs that can block serotonin-2 receptors. The company currently has a compound in clinical trials. Research into treatments for obsessive-compulsive disorders also involve some dysregulation or deficiency in serotonin receptors. "While we believe that serotonin dysregulation is critical, it is not the only player," says Dr. Richard C. Allen, vice president and general manager of the Hoechst Neuroscience Strategic Business Unit. "We are looking for drugs with a balanced profile in terms of their effects on a variety of receptors."

Hoechst-Roussel recently broke ground on its new molecular neurobiology laboratory. The new 46,000 sq.-ft. facility will allow multiple approaches to evaluating treatments for CNS diseases, as well as more long-term approaches, such as the control of gene expression and the establishment of assays on cloned human receptors. "When we can find the genes that are responsible for these disorders," says Dr. Allen, "we will be able to develop drugs to control gene transcription and expression, and halt the cause."

to find effective drugs and technologies. A number of federal agencies support research on the brain for a combined total of roughly \$2 billion a year. Much of this activity is funded by the National Institutes of Health, and the Alcohol, Drug Abuse, and Mental Health Administration. These agencies conduct their own research in intramural laboratories, but the majority of the research is supported through extramural grants to scientists in academic settings and medical centers.

The commitment to brain research also is seen at the highest policy levels of government. The White House Office of Science and Technology Policy (OSTP) was named by Congress as the coordinating agency for federal Decade of the Brain activities. In April 1991, OSTP issued the report "Maximizing Human Potential: Decade of the Brain 1990-2000," a comprehensive overview of brain research activities throughout the federal government. This report is an historic milestone, representing the highest level of attention ever given to basic brain and behavioral research in the federal science and technology policy establishment. In addition, the report represents the first time that all the federal agencies which support research on the brain have collaborated on an undertaking of this magnitude. (Copies of the OSTP report are available from NFBR.)

A DYNAMIC NEW INDUSTRY

Advances in brain research also have spawned the emergence of a dynamic neuroscience industry. The medical devices and pharmaceutical industries, through sustained investment in research and development, have made major contributions in bringing promising new diagnostic devices and therapeutics to clinicians and consumers, often at great cost. According to one study, it takes an average \$231 million, spread over 12 years, from synthesis to regulatory clearance, to bring a prescription drug to market. This does not include the costs of the underlying basic research. Pharmaceutical companies spend relatively more money on R&D than any other industry; 83% of total R&D funds are spent on new product research.

The dramatic revolution in biomedical technology, rising from advances in molecular biology and genetics, also gave birth to the biotechnology industry. Known for their innovative research, several new biotechnology companies have been formed, dedicated specifically to the development of diagnostics and therapeutics for treatment of brain disorders.

NEW TREATMENTS, BETTER UNDERSTANDING

The combination of basic scientific advances and modern technology has revolutionized the treatment of brain-related disorders. Brain operations previously thought impossible are now routine; natural compounds are being re-engineered to correct imbalances in the brain; and new drugs are under development which have the potential to alleviate a variety of brain disorders.

For many of these disorders, scientists often do not know exactly what area of the brain is involved. Similarly,

Hoechst-Roussel
Pharmaceuticals
Inc.



Pioneering Research in the Cause, Treatment and Cure of Alzheimer's Disease

Hoechst-Roussel Pharmaceuticals Inc.
Somerville, New Jersey 08876-1258

The name and logo HOECHST are registered trademarks of Hoechst AG.

Hoechst 


Q67003-1291

BIO-IMAGING TECHNOLOGIES INC.

Despite the analytic potential of imaging technologies that display the brain, "biomedicine has been almost exclusively observationally oriented," points out James J. Conklin, M.D., president and CEO of Bio-Imaging Technologies Inc. Further, "The use of proprietary hardware-software systems has prevented an exchange of image data between MRI devices, PET and SPECT scans, and CAT technologies."

Bio-Imaging Technologies' task is to bring quantitative disciplines into imaging. In doing so, the two-and-a-half year-old company from West Trenton, N.J., hopes to speed the process of clinical evaluation and regulatory review for its clients: biotechnology and pharmaceutical companies seeking FDA approval of drugs and medical devices. Because more precise data gleaned from images means the reliability of statistical evidence improves, Bio-Imaging theoretically can decrease the number of patients needed to provide clinical trial results. That, in turn, could reduce the average \$231 million cost and slash the typical 12-year timetable to bring a new drug to market.

Bio-Imaging's proprietary technology includes tools to insure that each imaging technology—whether it's a CAT or a PET scan—studies the same exact site. The company's image processing technology, based on satellite imaging techniques, digitizes and converts images to a standard format. Data from various systems



"Fused" MRI and PET images allow Bio-Imaging Technologies to pinpoint blood perfusion defects and evaluate recovery in stroke patients.

are fused and translated into three-dimensional image displays and quantitatively analyzed. By following particular patients over time, Bio-Imaging can build far more statistically valid analyses than more subjective measures allow.

For example, MRI brain studies of a stroke patient map the individual's anatomy; a PET scan clearly shows areas of the brain where blood does not flow and therefore are likely damaged. By combining those two tools in progressive time-studies, Bio-Imaging's technology can help pharmaceutical researchers to methodically chart improvement.

Bio-Imaging's technology applies to all diseases and conditions amenable to imaging—from gastrointestinal ulcers and heart disease to skin lesions. However, "The roots of our technology are in the brain," says Dr. Conklin. The technology was first used in CNS studies at Maryland Psychiatric Research Center and Johns Hopkins.

the chemical reactions associated with brain disorders often are not understood. Consequently, effectiveness of treatments vary, and side-effects are difficult to eliminate.

RECENT ADVANCES IN BRAIN RESEARCH

More has been learned about the brain's structure and functions in recent decades than in prior recorded history. During the 1960s and 1970s, enormous advances occurred in the methods used to study the chemistry of the brain, resulting in increased knowledge about the roles of natural chemicals and pharmaceuticals in mental disorders. In the 1980s, advances in molecular biology and imaging techniques opened new doors in the study of the roles of genetics in brain anatomy and activity. In the 1990s, the Decade of the Brain, science is poised to combine the experience of the past 20 years to refine the study of brain activity in living subjects. Science holds great promise for breakthroughs in a number of areas, including neurogenetics, neurobehavioral sciences, and neural injury.

Even a cursory review of where brain research has brought us in the past 20 years is staggering. Unprecedented advances in molecular genetics and functional and structural brain imaging have provided new ways to conceptualize the healthy brain and the injured or diseased brain. Currently, much attention is focused on alterations in neurotransmitter systems. Scientists are investigating the possible association of increased or decreased levels of dopamine with symptoms of schizophrenia, and the role of monoamines in mood disorders.


Molecular biology, combined with the techniques and knowledge of neuroscience, is bringing the development of drugs for diseases of the brain into a new phase through what is known as rational drug design (RDD). The principle behind RDD is that if the structure of a specific drug receptor is known, then it is possible to "tailor design" drugs which selectively interact with, and modulate, its functions. RDD focuses on understanding the physiological basis of disease, concentrating on the activity of hormones and hormone receptors, enzymes, cell replication, and protein synthesis.

BRAIN IMAGING

Until recently, examination of the human brain was possible only through surgery or autopsy. In the past 20 years, the potential to understand and treat brain disorders has been enormously advanced through the development of ways of "imaging" or seeing the brain operating intact, allowing for unprecedented access. Imaging techniques, already useful in the diagnosis of some mental disorders, are of fundamental importance in basic research aimed at mapping the regions of the brain and correlating those regions with function.

One technique, positron emission tomography (PET), allows measurement of energy levels—or activity—in the brain. By giving moment-to-moment pictures of the working brain, PET scans provide an inside look at

(continued on page BR11)



WE CAN MEASURE WHAT THE EYE CAN'T SEE

BIO-IMAGING TECHNOLOGIES, INC. utilizes advanced, computer-assisted medical image processing and image analysis to assist pharmaceutical and biotechnology companies in the clinical evaluation of drugs and medical devices.

BIO-IMAGING TECHNOLOGIES generates biostatistical measures from medical images captured by a variety of imaging and sensor devices, from MRI, CAT, PET and nuclear scanners to fiberscopes and infrared sensors. These biostatistical measures are used to quantitatively analyze the therapeutic effect of drugs or the diagnostic utility of medical devices.

BIO-IMAGING TECHNOLOGIES applies its medical image processing, image base and data management expertise to prepare fully interactive, computerized regulatory submissions.

To learn how we can help you to facilitate clinical development and regulatory review and **reduce time to market**, call William L. Robbins, Director, Business Development at 609-883-2000.



BIO-IMAGING

Technologies, Inc.

THE NEW MEASURE FOR DRUG AND DEVICE DEVELOPMENT

BIO-IMAGING TECHNOLOGIES, INC.
830 Bear Tavern Road, West Trenton, NJ 08628-1020
TEL: 609-883-2000 FAX: 609-883-7719

CIBA-GEIGY CORP.

Ciba-Geigy, the wholly owned U.S. subsidiary of a \$15 billion conglomerate headquartered in Basel, Switzerland, generates about one-third of Ciba-Geigy's worldwide pharmaceutical business from its Summit, N.J., division. Summit's research concentrates on arthritis, pulmonary disease, cardio-vascular, cancer, and central nervous system disorders.

"We have a corporate commitment to researching and developing drugs for CNS disorders," says Carlos Staglis, product director for the CNS area. "There is a tremendous emotional stigma and burden with CNS conditions and we're committed to

doing something about that."

Ciba-Geigy's long list of drugs for CNS conditions include Lioresal, for the relief of spasms in multiple sclerosis conditions; Ritalin, for attention deficit disorders; and Tofranil, for depression.

One of the most widely prescribed medications worldwide for the treatment of epilepsy is Ciba-Geigy's Tegretol. First introduced in Europe during the early 1960s, Tegretol became available in the U.S. in 1968 for the treatment of trigeminal neuralgia and in 1974 for epilepsy. Earlier this year, Ciba-Geigy submitted a new drug application for Tegretol



One of the most widely prescribed medications worldwide for the treatment of epilepsy is Ciba-Geigy's Tegretol.

OROS, a new slow-release formulation of Tegretol.

The CNS therapeutic area can claim its share of major successes lately. For example, Habitrol, a transdermal nicotine patch to aid in smoking cessation, was launched in December with the intent of helping many of the nation's estimated 50 million smokers quit by reducing cigarette craving. Another of the division's claims to fame is Anafranil, the first and only drug approved in the United States for treating Obsessive-Compulsive Disorder (OCD).

As many as one in 40 people, or 5 million Americans, suffers from OCD. These individuals experience uncontrollable recurring thoughts (obsessions) and feel forced to act out meaningless behaviors frequently and repeatedly (compulsions). The most common obsessions involve preoccupation with dirt and fear of acting on violent thoughts. The most common compulsions entail excessive cleaning, washing, counting, arranging rituals, and hair pulling.

Cleared by the FDA for marketing in 1989, Anafranil blocks the reuptake of serotonin and norepinephrine—neurochemicals that carry messages between nerve cells in the brain that have been linked to OCD behavior. Patients who responded to the drug averaged a 35% to 42% reduction in symptoms, with 85% saying they experienced some reduction in symptoms.

Books on the Brain

Stress, the Aging Brain, and the Mechanisms of Neuron Death

Robert M. Sapolsky

"A provocative hypothesis concerning the war and tear on the brain associated with stress and aging." — Bruce S. McEwen, Rockefeller University

A Bradford Book 448 pp., 79 illus. \$55.00 (October)

Behavioral Endocrinology

edited by Jill B. Becker, S. Marc Breedlove, and David Crews

"This book provides a superb introduction to the field... It will stimulate students and attract many to the next generation of behavioral endocrinologists."

— Arthur P. Arnold, UCLA

A Bradford Book 574 pp., 133 illus.
\$65.00 cloth, \$34.95 softcover

Culturing Nerve Cells

edited by Gary Banker and

Kimberly Goslin

foreword by Charles F. Stevens

"An excellent book that fills a substantial need in biomedical literature."

— Dennis W. Choi, Washington University School of Medicine

Cellular and Molecular Neuroscience Series

A Bradford Book 453 pp., 89 illus. \$55.00

The Computational Brain

Patricia S. Churchland and

Terrence J. Sejnowski

"...Essential reading for anyone interested in neuroscience." — James H. Schwartz, Columbia University, C.P.S.

Computational Neuroscience Series


A Bradford Book 576 pp., 366 illus. \$39.95

To order books call toll-free 1.800.356.0343 or 617.628.8569. MasterCard & VISA accepted.

55 Hayward Street **The MIT Press** Cambridge, MA 02142

Quality is never an accident.
It is always the result
of intelligent effort.
There must be the will
to produce a superior thing.

John Ruskin



At CIBA-GEIGY,
people are
the active ingredient
to pharmaceutical
innovations of tomorrow...

Pharmaceuticals Division
CIBA-GEIGY Corporation
Summit, New Jersey 07901

DU PONT MERCK

The Du Pont Merck Pharmaceutical Company was founded in 1991 with a vision of its becoming a multinational, research-based drug company which would bring novel scientific solutions to meeting major unmet medical needs.

The new company today is a fully integrated, free-standing entity owned equally by Du Pont and Merck & Co. Inc. Its headquarters are in Wilmington, Delaware.

"Our long-term research commitments are arthritis, dementia, thrombosis, atherosclerosis, and cancer," says David W. Martin, executive vice president for research and development. "We are committed to these specific diseases, and if a particular therapeutic hypothesis fails, we won't abandon our commitment; we will continue with another target for the same disease."

This commitment may come to fruition soon, as two promising new compounds, one for hypertension and one for dementia of the Alzheimer's type, are now in the final phases of clinical testing. Other near-term entities are potential drugs for breast cancer, psychosis, arthritis, and organ transplantation. Altogether, the company spends about \$250 million annually on R&D.

In this "Decade of the Brain," it is Du Pont Merck's "Aviva" (linopirdine) that is generating considerable anticipation as it reaches completion of Phase III trials. It is designed to enhance normal neurotransmission in the brain by targeting the enhanced release of certain neurotransmitters that play key roles in information processing, memory, and cognition.

While other therapies are being developed that affect one neurotransmitter—acetylcholine—Errol DeSouza, Du Pont Merck's director of central nervous system research, explains that: "'Aviva' seems to enhance the release of three neurotransmitters—acetylcholine, serotonin, and dopamine. It acts in multiple transmitter systems."

The goal of this "multiple transmitter" therapeutic approach is to help alleviate some of the conditions that often occur in the early-to-moderate stages of Alzheimer's disease. These most often include a deficit in information processing, which impairs memory and a decline in activities of daily living. This represents one of the most important immediate research goals of Du Pont Merck, an emerging leader in pharmaceuticals.



Du Pont Merck locates its research facilities in Wilmington, Delaware.

(continued from page BR7)

how and where brain activity occurs; they provide maps of emotions, learning, vision, and memory.

While PET scans tell us about brain activity, magnetic resonance imaging (MRI) shows the structures of the brain and is useful for detecting tumors and following the progress of diseases, as well as helping us to understand normal changes as the brain ages. Magnetic resonance spectroscopy (MRS) is similar to MRI technology, but examines brain chemistry rather than anatomy. The newer imaging techniques supplement and complement the more familiar techniques of computerized axial tomography (CAT) scans and the electroencephalograph (EEG).

The information gathered from imaging studies has affected our understanding of the mind/brain relationship by breaching the biological limitations imposed by the inaccessibility of the human brain.

INFORMATION TECHNOLOGY: NEURAL NETWORKS

Brain research has contributed to, as well as benefitted from advances in computers. The brain and nervous system are marvelously effective in processing sensory information, relying on complex connections between neurons that give the brain the crucial ability to recognize patterns and learn from them.

Scientists try to copy parts of the brain's design in computer systems called artificial neural networks. There are two kinds of neural networks. One is a type of computer software that allows a normal computer to simulate information in "brain-like" patterns or "chunks," rather than processing one piece of data at a time. The other type of neural network is a special kind of computer with thousands of simple devices connected to one another in a brain-like way.

A neural network attempts to solve a problem with the information that it gets from repeated exposure to patterns, such as speech or handwriting. The network is able to "learn from supervision" and adapt to new information.

In addition to its contributions in the field of neuroscience, neural network research also has significant implications for the nation's economic competitiveness. One current emphasis in industry is to develop newer neural-network chips that will speed information processing. ■

THE NATIONAL FOUNDATION FOR BRAIN RESEARCH JOINS
SCIENTIFIC AMERICAN IN THANKING THE INDUSTRY
SPONSORS WHO MADE THIS REPORT POSSIBLE:

BIO-IMAGING TECHNOLOGIES INC.

CIBA-GEIGY CORP.

DU PONT MERCK PHARMACEUTICAL COMPANY

HOECHST-ROUSSEL PHARMACEUTICALS

MIT PRESS

SCHERING-PLOUGH CORP.


Cover photo courtesy
of Bio-Imaging
Technologies Inc.



A Moment to Remember

Why have we committed so many resources, and so many people,
in hope of finding answers to the devastating effects of dementia and Alzheimer's disease?

The answer is really quite simple.



© 1992 Du Pont Merck
Wilmington, DE
© 1992 SCIENTIFIC AMERICAN, INC

DECADE OF THE BRAIN

WHERE TO FIND MORE INFORMATION



Many organizations exist to help patients and families living with various disorders of the brain. These organizations typically provide information about the treatments available for the particular disorder and useful information to families, and many support research.

ALZHEIMER'S DISEASE

Alzheimer's Association
70 East Lake Street
Suite 600
Chicago, IL 60601
(312)853-3060; (800)621-0379

French Foundation for Alzheimer Research
11620 Wilshire Boulevard
Suite 820
Los Angeles, CA 90025
(213)445-4650; (800)477-2243

AMYOTROPIC LATERAL SCLEROSIS

The ALS Association
21021 Ventura Boulevard
Suite 321
Woodland Hills, CA 91364
(818)340-7500; (800)782-4747

ANXIETY DISORDERS

Anxiety Disorders Association of America
6000 Executive Boulevard, #513
Rockville, MD 20852
(900)737-3400

ATAXIA

National Ataxia Foundation
15500 Wayzata Boulevard, #750
Wayzata, MN 55391
(612)473-7666

ARTERIOVENOUS MALFORMATION

The AVM Support Group
107 Bella Vista Way
San Francisco, CA 94127
(415)421-7330

AUTISM

Autism Society of America
8601 Georgia Avenue
Suite 503
Silver Spring, MD 20910
(301)565-0433

The New Jersey Center for Outreach and Services for the Autism Community Inc.
123 Franklin Corner Road
Suite 215
Lawrenceville, NJ 08648
(609)895-0190

BATTEN'S DISEASE

Batten's Disease Support and Research Association
2600 Parsons Avenue
Columbus, OH 43207
(800)448-4570

BIRTH DEFECTS

March of Dimes Birth Defects Foundation
1275 Mamaroneck Avenue
White Plains, NY 10605
(914)428-7100

BLEPHAROSPASM

Benign Essential Blepharospasm Foundation Inc.
2929 Calder Avenue, Suite 304
Beaumont, TX 77704
(409)832-0788

BRAIN TUMOR

American Brain Tumor Assn.
3725 N. Talman Avenue
Chicago, IL 60618-4712
(312)286-5571; (800)886-2282

Brain Research Fund Foundation of London
111 Waterloo Street, Suite 600
London, Ontario N6B 2M4
Canada
(519)624-7755

Brain Tumor Foundation for Children Inc.
2231 Perimeter Park Dr.
Suite 9
Atlanta, GA 30341
(404)458-5564

Brain Tumor Society
258 Harvard Street, Suite 308
Brookline, MA 02146
(617)433-7033

The Children's Brain Tumor Foundation
35 Alpine Lane
Chappaqua, NY 10514
(914)238-4917

National Brain Tumor Foundation
323 Geary Street, Suite 510
San Francisco, CA 94102
(800)934-CURE

CAREGIVING

Family Survival Project for Brain-Impaired Adults
425 Bush Street
Suite 500
San Francisco, CA 94108
(415)434-3388

CHARCOT-MARIE-TOOTH DISEASE

Charcot-Marie-Tooth Association
601 Upland Avenue
Upland, PA 19015
(215)499-7486

Charcot-Marie-Tooth Intl.
1 Springbank Drive
St. Catherine, Ontario L2S 2K1
Canada
(416)687-3630

DEPRESSION

Depression After Delivery
P.O. Box 1282
Morrisville, PA 19067
(215)295-3994; (800)944-4773

National Foundation for Depressive Illnesses
P.O. Box 2257
New York, NY 10116
(800)248-4344

DIZZINESS

Vestibular Disorders Association
P.O. Box 4467
Portland, OR 97208-4467
(503)229-7705

DYSAUTONOMIA

Dysautonomia Foundation Inc.
20 East 46th Street
Suite 302
New York, NY 10017
(212)949-6644

DYSLEXIA

The Orton Dyslexia Society
Chester Building
8600 LaSalle Road
Suite 382
Baltimore, MD 21204-6020
(410)296-0232

DYSTONIA

Dystonia Medical Research Foundation
8383 Wilshire Blvd., Suite 800
Beverly Hills, CA 90211
(213)852-1630

EPILEPSY

Epilepsy Foundation of America
4351 Garden City Drive
Landover, MD 20785
(301)459-3700

GENERAL DISABILITIES

National Easter Seal Society Inc.
70 East Lake Street
Chicago, IL 60601
(312)726-6200 (Voice); -4258

GENETIC DISORDERS

Alliance of Genetic Support Groups
35 Wisconsin Circle, Suite 440
Chevy Chase, MD 20815
(301)652-5553; (800)336-GENE

National Foundation for Jewish Genetic Diseases
250 Park Avenue, Suite 1000
New York, NY 10017
(212)554-9712

GUILLAIN-BARRE SYNDROME

Guillain-Barre Syndrome Foundation International
P.O. Box 262
Wynnewood, PA 19096
(215)667-0131

HEADACHE

American Association for the Study of Headache
875 Kings Highway
Suite 200
West Deptford, NJ 08096
(609)845-0322; (800)255-ACHE

American Council for Headache Education
875 Kings Highway
Suite 200
West Deptford, NJ 08096
(609)845-0322; (800)255-ACHE

National Headache Foundation
5252 N. Western Avenue
Chicago, IL 60625
(800)843-2256

HEAD INJURY

National Head Injury Foundation
1140 Connecticut Avenue, NW
Suite 812
Washington, DC 20036
(202)296-6443; (800)444-6443

HUNTINGTON'S DISEASE

Hereditary Disease Foundation
1427 7th Street, Suite 2
Santa Monica, CA 90401-9990
(213)458-4183

Huntington's Disease Society of America

140 W. 22nd Street, 6th Floor
New York, NY 10011-2420
(212)242-1968; (800)345-HDSA

HYDROCEPHALUS

National Hydrocephalus Foundation
400 N. Michigan Ave., Suite 1102
Chicago, IL 60611
(312)645-0701; (800)785-6161

LEARNING DISABILITIES

Learning Disabilities Association of America
4156 Library Road
Pittsburgh, PA 15234
(412)341-1515; (412)341-8077

National Center for Learning Disabilities
99 Park Avenue
6th Floor
New York, NY 10016
(212)687-7211

LEUKODYSTROPHY
United Leukodystrophy Foundation
2304 Highland Drive
Sycamore, IL 60178
(800)728-5483

MENTAL ILLNESS
Natl. Alliance for the Mentally Ill
2101 Wilson Blvd.
Suite 302
Arlington, VA 22201
(703)524-7600; (800)950-NAMI

National Mental Health Assn.
1021 Prince Street
Alexandria, VA 22314
(800)969-NMHA

MENTAL RETARDATION
The Arc
500 E. Border Street
Suite 300
Arlington, TX 76010
(817)261-6003

MULTIPLE SCLEROSIS
Multiple Sclerosis Association of America
601-605 White Horse Pike
Oaklyn, NJ 08107
(800)833-4MSA

Multiple Sclerosis Foundation
6350 N. Andrews Avenue
Fort Lauderdale, FL 33309
(305)776-6805; (800)441-7055

National Multiple Sclerosis Society
733 3rd Avenue
6th Floor
New York, NY 10017
(212)986-3240; (800)624-8236

MUSCULAR DYSTROPHY
Muscular Dystrophy Association
3561 East Sunrise Drive
Tucson, AZ 85718
(800)572-1717

MYASTHENIA GRAVIS
Myasthenia Gravis Foundation
53 West Jackson Boulevard,
Suite 600
Chicago, IL 60604
(312)427-6252; (800)541-5454

MYOCLONUS
Myoclonus Research Foundation
200 Old Palisade Road
Suite 17D
Fort Lee, NJ 07024
(201)585-0770

NARCOLEPSY
American Narcolepsy Association
P.O. Box 26230
San Francisco, CA 94126-6230
(415)788-4793

Narcolepsy Network
P.O. Box 1365
FDR Station
New York, NY 10150
(415)591-7884; (914)834-2855

National Sleep Foundation
122 S. Robertson Boulevard
Suite 201
Los Angeles, CA 90048
(310)288-0466

NEUROFIBROMATOSIS
Neurofibromatosis Inc.
3401 Woodridge Court
Mitchellville, MD 20721-2817
(301)577-8984

OBSESSIVE-COMPULSIVE DISORDER
Obsessive Compulsive Foundation
P.O. Box 9573
New Haven, CT 06535
(203)772-0565

PAIN
American Chronic Pain Assn.
P.O. Box 850
Rocklin, CA 95677
(916)632-0922

National Chronic Pain Outreach Association
7979 Old Georgetown Road
Suite 100
Bethesda, MD 20814-2429
(301)652-4948

PARAPLEGIA
American Paraplegia Society
75-20 Astoria Boulevard
Jackson Heights, NY 11370
(718)803-3782

PARKINSON'S DISEASE
American Parkinson Disease Association
60 Bay Street
Staten Island, NY 10301
(718)981-8001; (800)223-2732;
(800)732-9550 (toll free in NY)

California Parkinson's Foundation
2444 Moorpark Avenue
Suite 316
San Jose, CA 95128
(408)998-2722

National Parkinson Foundation
1501 NW 9th Ave.
Bob Hope Road
Miami, FL 33136
(305)547-6666; (800)327-4545;
(800)433-7022 (toll free in FL)

Parkinson Support Groups of America
11376 Cherry Hill Road
#204
Beltsville, MD 20705
(301)937-1545

Parkinson's Disease Foundation
Columbia-Presbyterian Medical Center
650 West 168th Street
New York, NY 10032
(212)923-4700; (800)457-6676

Parkinson's Educational Program - USA
3900 Birch Street
#105
Newport Beach, CA 92660
(714)250-2975; (800)344-7872

United Parkinson Foundation
360 W. Superior Street
Chicago, IL 60610
(312)664-2344

POSTPOLIO SYNDROME
International Polio Network
5100 Oakland Avenue
#206
St. Louis, MO 63110
(314)534-0475

Polio Survivors Association
12720 La Reina Avenue
Downey, CA 90242
(310)862-4508

PROGRESSIVE SUPRANUCLEAR PALSY
Society for Progressive Supranuclear Palsy Inc.
2904 Marnat Road
Baltimore, MD 21209-2420
(410)484-8771

RETT SYNDROME
International Rett Syndrome
8511 Rose Marie Drive
Fort Washington, MD 20744
(301)248-7031

REYE'S SYNDROME
National Reye's Syndrome Foundation
426 North Lewis
Bryan, OH 43508
(419)636-2679; (800)233-7893;
(800)231-7393 (Toll free in OH)

SPINAL CORD INJURY
American Paralysis Association
500 Morris Avenue
Springfield, NJ 07081
(201)379-2690; (800)225-0292

National Spinal Cord Injury Association
600 W. Cummings Park
Suite 2000
Woburn, MA 01801
(617)935-2722; (800)962-9629

Paralyzed Veterans of America
Spinal Cord Research Foundation
801 18th Street, NW
Washington, DC 20006
(202)416-7659

Spinal Cord Society
Wendell Road
Fergus Falls, MN 56537
(218)739-5252

SPINAL MUSCULAR ATROPHY
Families of S.M.A.
P.O. Box 1465
Highland Park, IL 60035
(708)432-5551

STROKE
American Heart Association
7272 Greenville Avenue
Dallas, TX 75231
(214)373-6300
(800)242-8721

National Stroke Association
300 E. Hampden Avenue
Suite 240
Englewood, CO 80123-2654
(303)762-9922
(800)STR-OKES

STURGE-WEBER SYNDROME
The Sturge-Weber Foundation
P.O. Box 460931
Aurora, CO 80046
(303)360-7290; (800)627-5482

SYRINGOMYELIA
American Syringomyelia Alliance Project
P.O. Box 1586
Longview, TX 75606-1586
(903)236-7079

TAY-SACHS
National Tay-Sachs and Allied Diseases Association Inc.
2001 Beacon Street
Brookline, MA 02146
(617)277-4463

TOURETTE SYNDROME
Tourette Syndrome Association
42-40 Bell Boulevard
Bayside, NY 11361
(718)224-2999
(800)237-0717

TREMOR
International Tremor Foundation
360 W. Superior Street
Chicago, IL 60610
(312)664-2344

TUBEROUS SCLEROSIS
National Tuberous Sclerosis Association
8000 Corporate Drive
Suite 120
Landover, MD 20785
(301)459-9888; (800)225-6872



Working Memory and the Mind

Anatomic and physiological studies of monkeys are locating the neural machinery involved in forming and updating internal representations of the outside world. Such representations form a cornerstone of the rational mind


by Patricia S. Goldman-Rakic

The seeming simplicity of everyday life belies the enormously complex ongoing operations of the mind. Even routine tasks such as carrying on a conversation or driving to work draw on a mixture of current sensory data and stored knowledge that has suddenly become relevant. The combination of moment-to-moment awareness and instant retrieval of archived information constitutes what is called the working memory, perhaps the most significant achievement of human mental evolution. It enables humans to plan for the future and to string together thoughts and ideas, which has prompted Marcel Just and Patricia Carpenter of Carnegie Mellon University to refer to working memory as "the blackboard of the mind."

Until recently, the fundamental processes involved in such higher mental functions defied description in the mechanistic terms of science. Indeed, for the greater part of this century, neurobiologists often denied that such functions were accessible to scientific analysis or declared that they belonged strictly to the domain of psychology and philosophy. Within the past two decades, however, neuroscientists have made great advances in understanding the relation between cognitive processes and the anatomic organization of the brain. As a consequence, even global mental attributes such as thought and intentionality can now be meaningfully studied in the laboratory.

The ultimate goal of that work is extraordinarily ambitious. Eventually researchers such as myself hope to be able to analyze higher mental functions in terms of the coordinated activation of neurons in various structures in the brain. It should also be possible to identify the cells that mediate the activity of those structures. Such research will help explain the origin of mind. It may also lead to more complete descriptions of baffling mental disorders such as schizophrenia.

For many years, insight into the operation of the brain was stymied by the misconception that memory is a single entity that could be traced to a single structure or location. Since the 1950s, neuroscientists have increasingly come to appreciate that memory consists of multiple components constructed around a distributed network of neurons. According to present thinking, a form of memory known as associa-



tive memory acquires facts and figures and holds them in long-term storage. That knowledge is of no use, however, unless it can be accessed and brought to mind in order to influence current behavior.

Working memory complements associative memory by providing for the short-term activation and storage of symbolic information, as well as by permitting the manipulation of that information. A simple activity involving working memory is the carry-over operation in mental arithmetic, which requires tempo-

rarily storing a string of numbers and holding the sum of one addition in mind while calculating the next. More complex examples include planning a chess move or constructing a sentence. Working memory in humans is considered fundamental to language comprehension, to learning and to reason.

Numerous lines of evidence indicate that the operations of working memory are carried out in a part of the brain known as the prefrontal lobes of the cerebral cortex. (Cortex derives from the Latin word meaning bark; the cerebral cortex consists of an outer rind of so-called gray matter neurons surrounding the cerebrum.) Much of the evidence identifying this structure as the center for working memory comes from observations of the effects of injuries to the prefrontal part of the hemispheres. For example, patients having frontal lobe damage exhibit gross deficiencies in how they use knowledge to guide their behavior in everyday situations. Nevertheless, they often retain a full store of information and may continue to score well on conventional tests of intelligence.

Although most fully developed in humans, some elements of working memory exist in other animals, especially in other primates; if their prefrontal cortices are damaged, those animals develop symptoms much like the ones seen in humans. Neuroscientists have therefore turned to monkeys in their efforts to explore the nature of working memory.


WORKING MEMORY enables a human to retrieve stored symbolic information, such as the bowings and fingerings of a memorized piece of music, and to translate that information into a controlled set of motor activities. Studies of similar but simpler information processing performed by primates is revealing the structure of working memory.

PATRICIA S. GOLDMAN-RAKIC has devoted her academic career to studying the neurobiology of memory and cognition. She received a Ph.D. from the University of California, Los Angeles, in 1963. Two years later she joined the Intramural Research Program of the National Institute of Mental Health. In 1979 she moved to the Yale University School of Medicine, where she is a professor of neuroscience. Goldman-Rakic sits on several national advisory boards and is a member of the National Academy of Sciences. She recently served as president of the Society for Neuroscience. Her current research focuses on identifying the neural mechanisms that carry out higher cortical functions in primates.

WORKING MEMORY TASK



CUE




DELAY




RESPONSE

ASSOCIATIVE MEMORY TASK



CUE AND
RESPONSE



DELAY



CUE AND
RESPONSE

RIGHT WRONG

MEMORY TASKS help to assess the workings of the mind. In the classical working memory task (*left*), a monkey briefly views a target stimulus—in this case, a morsel of food. Only after a delay is the animal allowed to retrieve the food. The experimenter randomly varies the location of the food between trials, so that each response tests only the animal's short-term retention of visual and spatial information. An associative memory task (*right*), in contrast, follows a consistent pattern throughout. Here a plus sign always indicates the correct response. The task therefore measures the animal's ability to retain long-term rules.

Such exploration has been aided by the design of repeatable tests of working memory functions.

Working memory is being assessed in monkeys by means of tasks known as delayed-response tests, which evaluate an organism's ability to react to situations on the basis of stored or internalized representations rather than

on information immediately present in the environment. In the prototypical delayed-response test, an animal receives a brief visual or auditory stimulus that is then hidden or taken away. After a delay of several seconds, the animal is given a signal that tells it to respond to the location where the stimulus had appeared. If the response is correct, the

animal receives a reward, usually food or juice.

Delayed-response tests tap working memory processes because the animal must retain the memory of the location of the stimulus during the period of the delay. The proper response at the end of the delay is indicated not by external stimuli but by the memory of what the subject saw on the previous trial. Furthermore, the correct response may differ from one trial to the next, depending on new information presented to the subject in each trial. Correct responses in working memory tasks, as in human affairs, are guided by memory rather than by immediate sensory information, and they depend on constant updating of the relevant information.

Delayed-response tests resemble very closely the object-permanence task, developed in the early part of this century by the French child psychologist Jean Piaget, that is widely used to chart the cognitive development of young children. For Piaget's task, a child is shown two boxes, one of which contains a toy. The boxes are then closed. After a brief wait, during which the child is purposely distracted, the child is asked to pick which box contains the toy. Once the child gives several consecutive correct responses, the toy is switched into the other box while the child watches. The experimenter then continues the test to find out whether the child will change his or her response in accord with the updated information.

A series of studies has demonstrated that performance on the object-permanence task, like the ability to conduct delayed-response activities, depends on the degree of maturity of the subject's prefrontal cortex. Human infants less than about eight months old (whose cortices have not yet acquired adult circuitry) perform poorly on these tasks, as do monkeys whose prefrontal regions have been surgically ablated. In both cases, the subjects' responses are guided by habit and by reflex rather than by representational principles. Infants and brain-injured monkeys tend to repeat the response that previously was reinforced (for example, choosing the box on the right even after they have seen that the toy was transferred to the box on the left) rather than change their response to agree with newly presented information. Both humans and monkeys act as if "out of sight" is "out of mind."

Such behavior implies that the mechanism for guiding behavior by representational knowledge is destroyed in monkeys having prefrontal lesions and not yet developed in human infants. In

support of that notion, I, along with Jean-Pierre Bourgeois and Pasko Rakic, also at Yale University, have examined the rate at which neural connections form in the prefrontal cortices of juvenile monkeys.

The time of most rapid synapse formation in the animals' prefrontal region occurs when the animals are roughly two to four months old, the same age at which the monkeys acquire the capacity to perform delayed-response tasks. The concept that an object exists continuously in space and time even when out of view and, more generally, the ability to form abstract concepts may depend on a fundamental capacity to store representations of the outside world and to respond to those representations even when the real objects are not present.


The studies described above raised the enticing possibility of identifying more precisely the brain structures associated with delayed-response activities and representational memory. Much of the progress toward that goal has derived from experiments that monitor electrical activity in single neurons in monkeys' prefrontal cortices while the animals perform tasks that depend on specific delayed-response skills.

Joaquin M. Fuster of the University of California at Los Angeles, along with Kiso Kubota and Hiroaki Niki of the Kyoto Primate Center in Japan, performed the first experiments of how individual neurons behave in the prefrontal cortex. The researchers introduced fine electrodes into the prefrontal cortices of monkeys trained to perform simple delayed-response tasks and then recorded the animals' neuronal activity in relation to the events in the task. Those studies revealed a range of responses among the neurons in the prefrontal cortex. Some cells showed heightened electrical activity when information was presented, whereas others became active during the delay period, when the animals were remembering the information. A third set of neurons responded most strongly when the animals began their motor response.

At Yale, Shintaro Funahashi, Charles J. Bruce and I have used the single-neuron technique in conjunction with a delayed-response experiment that tests spatial memory. For our experiment, a monkey is trained to fix its gaze on a small spot in the center of a television screen. A visual stimulus, typically a small square, appears briefly in one of eight locations on the screen and then vanishes. At the

end of a delay of three to six seconds, the central light, or fixation spot, switches off, instructing the animal to move its eyes to the location where the stimulus was seen before the delay. If the response is correct, the animal is rewarded with a sip of grape juice. Because the animal's gaze is locked onto the fixation spot, each stimulus activates a specific set of retinal cells. Those cells, in turn, trigger only a certain subset of the visual pathways in the brain.

Using the eye-movement experiment, we have demonstrated that certain neurons in the prefrontal cortex possess what we call "memory fields": when a particular target disappears from view, an individual prefrontal neuron switches into an active state, producing electrical signals at more than twice the baseline rate. The neuron remains activated until the end of the delay period, when the animal delivers its response. A given neuron appears always to code the same visual location. For example, some neurons fire only if the stimulus appears at the nine o'clock position on the television screen; the cell does not respond to visual stimuli that appear elsewhere in the monkey's visual field. Other neurons code for other target locations in working memory.



DELAYED-RESPONSE TASK has been used to study working memory in monkeys. While a monkey fixes its gaze on a central spot, a target flashes on the screen (*left*), then vanishes. During a delay of several seconds, the monkey keeps a memory of the spot "in mind" (*center*). When the central spot turns

off, the animal moves its eyes to look where the target appeared (*right*). Measurements of electrical activity show that certain neurons in the prefrontal cortex react to the appearance of the target, others hold the memory of it in mind and still others fire in preparation for a motor response.



NEURONAL CIRCUITRY connects the prefrontal cortex to the sensory, limbic and motor systems in a monkey brain (*top*). Anatomic studies show that neural projections from the parietal lobe to the prefrontal cortex exhibit a modular pattern, as seen in this frontal cross section (*middle*). Radioactive tracers reveal the metabolic activity in a frontal cross section of the brain of a monkey performing a delayed-response task (*bottom*). The distribution of activity closely conforms to the anatomic links.

The neurons capable of retaining the visual and spatial coordinates of a stimulus (in other words, of keeping its location "in mind" after it vanishes) appear to be organized together within a specific area of the prefrontal cortex. These neurons collectively form the core of the spatial working memory system. If the activity of one or more of these neurons falters during the delay period—if the animal is distracted, for example—the animal will probably make an error.

The activation of prefrontal neurons during the delay period of a delayed-response task depends neither on the presence of an external stimulus nor on the execution of a response. Rather the neural activity corresponds to a mental event interposed between the stimulus and the response. Monkeys whose prefrontal cortices have been damaged have no difficulty in moving their eyes to a visible target or in reaching for a desired object, but they cannot direct those motor responses by remembering targets and objects that are no longer in evidence.

Because the prefrontal cortex functions as an intermediary between memory and action, one can imagine that damage to the prefrontal cortex could spare knowledge about the outside world yet destroy the organism's ability to bring that stored knowledge to mind and to utilize it. Indeed, monkeys whose prefrontal cortices have been damaged, as well as many humans with similar injuries, exhibit no difficulty learning sensory-discrimination tasks. All forms of associative, or long-term, learning are preserved as long as the subject can still find the familiar environmental stimuli associated with certain consequences and expectations [see "The Biological Basis of Learning and Individuality," by Eric R. Kandel and Robert D. Hawkins, page 78].

Over the past decade, improved techniques for investigating the anatomy of the brain have provided for the first time an accurate and detailed picture of how the prefrontal cortex connects



with major sensory and motor control centers. Various researchers have found that the part of the cortex near the principal sulcus, a large groove in the prefrontal cortex, is critical for the visual and spatial working memory functions. I have focused my research on this particular region in the belief that an in-depth neurobiological analysis of one major subdivision of the prefrontal cortex could serve as a starting point for analysis of the other subdivisions of the brain and help lead the way to development of a unified theory of the function of the entire prefrontal cortex.

Studies of direct and indirect neuronal linkages in the brain reveal that the

prefrontal cortex is part of an elaborate network of reciprocal connections between the principal sulcus and the major sensory, limbic and premotor areas of the cerebral cortex. That particular network seems to be dedicated to spatial information processing. The network's structure probably follows the same basic plan as do other similarly organized networks that draw on multiple parts of the brain and are dedicated to other cognitive functions—object recognition, language production and comprehension, and mathematical reasoning, for example.

As previously noted, delay-response experiments demonstrate that neurons in the principal sulcus are sensitive to the specific location of visual stimuli. Those neurons must therefore have access to visual and spatial information originating elsewhere in the brain. The principal sulcus does in fact receive signals from the posterior parietal cortex, where the brain processes spatial vision. Clinical studies have documented that damage to the parietal cortex in humans causes spatial neglect, a loss of awareness of the body and its relation to objects in the outside world.

Given that working memory depends on accessing and bringing to mind information that is stored in long-term memory, one might presume that the principal sulcus also interacts with the hippocampus, the neuronal structure that controls associative, or learned, memory. Researchers have used radioactive amino acids to trace direct connections between the principal sulcus and the hippocampus.

My colleague Harriet Friedman, also at Yale, and I have used a remarkable technique known as autoradiography to measure brain metabolism. Our work shows that the hippocampus and the principal sulcal areas of the cortex are often simultaneously active during delayed-response tests. My co-workers and I think that the primary role of the hippocampus is to consolidate new associations, whereas the prefrontal cortex is necessary for retrieving the products of such associative learning (facts, events, rules) from long-term storage elsewhere in the brain for use in the task at hand.

A particularly useful version of autoradiography, called the 2-deoxyglucose method, has made it possible to observe directly which parts of the brain are activated during specific tasks. In this technique, developed by Louis Sokoloff of the National Institute of Mental Health, animals are injected with the compound 2-deoxyglucose, a molecule that appears chemically identical to glucose, the sugar that cells consume to

provide energy. The more active a cell is, the more 2-deoxyglucose it takes in. Unlike normal glucose, however, 2-deoxyglucose cannot be broken down by metabolic activity, so it accumulates in the cell. Sokoloff uses a radioactive version of the compound. The concentration of radioactivity in each part of the brain is therefore directly proportional to how active the cells there have been.

For our studies, a monkey trained to perform the delayed-response tasks receives an intravenous injection of radioactive 2-deoxyglucose. Immediately after completing the task, the animal is sacrificed and its brain is dissected into thin slices that are placed on photographic film. Radioactivity darkens the film, so each exposure serves as a snapshot of the activity of the cells in one particular slice of the brain.

My colleagues and I have found that the prefrontal cortex, as well as many of the areas with which it is connected (for example, the hippocampus, the bottom portion of the parietal cortex and the thalamus), exhibits a high level of metabolic activity during delayed-response performance. The same areas are notably less active when the monkey performs associative memory tasks that do not depend on short-term, rapid updating of information.

These results confirm anatomic studies of the connections between the prefrontal cortex and other parts of the brain. More significantly, they also reveal the degree to which various parts of the brain are engaged in certain discrete memory tasks. The studies also hint at how the prefrontal cortex organizes the many different kinds of information that must flood through it. In fact, patterns of brain activity appear distinctly different depending on whether the task calls up memories of location or of attributes of objects.

I think the prefrontal cortex is divided into multiple memory domains, each specialized for encoding a different kind of information, such as the location of objects, the features of objects (color, size and shape) and additionally, in humans, semantic and mathematical knowledge. Recently Fraser Wilson and James Skelly in my laboratory at Yale have begun to define an area below the principal sulcus in monkeys where neurons respond preferentially to complex attributes of objects rather than to their locations. They have found neurons there that increase their rate of firing when a monkey is remembering a red circle but not when calling up a memory of a green square, for example.

Noninvasive imaging techniques are

increasingly being used to monitor activation patterns in the human brain and to identify which neurons are engaged during specific mental tasks. One, known as positron emission tomography (PET), resembles autoradiography in that the subject takes in a radioactive compound that exposes changes in blood flow to a given region of the brain, indirectly displaying that region's degree of metabolic activity. Another way to record human brain activity is to measure the changing electrical potentials on the scalp in response to controlled sensory stimulation, a procedure called electroencephalography (EEG). Neither PET scans nor EEGs can provide anything close to the resolution possible in 2-deoxyglucose studies in animals, but they are invaluable tools for monitoring the human brain during mental activity.

A series of PET studies at Hamner-Smith Hospital in London and at Washington University examined subjects performing tasks that required them to keep a mental record of recently presented lists of words. Another PET experiment by the Washington University group required subjects to generate an appropriate verb to accompany a noun flashed in front of them on a card. The participants in all three tests displayed heightened neuronal activity in the prefrontal cortex while performing their tasks, all of which engaged working memory.


In a complementary study, Robert T. Knight of the University of California at Davis looked at EEGs of patients whose frontal lobes were injured. He

asked them to perform tasks that depended on comparing current auditory stimuli with recently presented ones in order to detect whether they are the same or different. Frontal lobe patients displayed patterns of electrical activity quite unlike those of healthy subjects performing the same task, suggesting that the patients do not store recent information in memory the same way as do normal people.


In one study, subjects were exposed to steady patterns of low and high tones and occasional, unexpected auditory stimuli. Healthy people developed positive electrical potentials on their cortices within one third of a second of hearing the anomalous sound. Patients who had lesions in their prefrontal cortices showed no such response, although they reacted normally to the familiar background tones. These data are consistent with the notion that the prefrontal cortex temporarily stores information against which current stimuli are judged.

The ultimate function of the neurons in the prefrontal cortex is to excite or inhibit activity in other parts of the brain. In this way, information processed in the principal sulcus can direct neurons in the motor centers that in turn carry out movements of the eyes, mouth, hands and other parts of the body. Whole-brain studies tell only part of the story; to understand the details of how signals pass to and from the prefrontal cortex, one must scrutinize the brain on a cellular scale.

When viewed through a conventional microscope, the cerebral cortex appears



PYRAMIDAL NEURON (left) in the prefrontal cortex is thought to modulate signals to and from the prefrontal cortex. Each such neuron is covered with thousands of spines, bulblike projections where synaptic connections occur. Synapses have different morphologies depending on whether they are excitatory or inhibitory (right). The dopamine-containing connections in the cortex are of the inhibitory type.



ELABORATE FLOW OF NEURAL SIGNALS is involved in producing a memory-guided eye movement. A neuron in the fifth layer of the prefrontal cerebral cortex transmits signals along a chain of neurons in the striatum, the substantia nigra and the superior colliculus, where they trigger motor response in the eyes. Impulses from the substantia nigra travel to the mediodorsal thalamus and back to the cortex, indicating the completion of the motor response and signaling the prefrontal neuron to return to a baseline level of activity. The graphs show the electrical activity of the neurons; inverted triangles indicate the nearly instantaneous travel of the signals.

to be divided into six layers of varying cellular composition and density. Cells in each layer form their own set of connections within the brain. One class of cell, which resides in the fifth layer of the cortex, projects to areas beyond the cortex, including the caudate nucleus and putamen (which regulate a variety of motor activities) and the superior colliculus (which specifically processes visual motor functions). Neurons in the sixth layer of the cerebral cortex project into the thalamus, through which sensory inputs from the brain's periphery travel to reach the cortex.

The prefrontal cortex probably cannot independently trigger motor responses. Nevertheless, it may regulate motor behavior by initiating, programming, facilitating and canceling commands to brain structures that are more immediately involved in directing muscular movement. Such commands are transmitted via an elaborate set of chemical pathways in the brain. Neuroscientists and biochemists around the world have been racing to learn more about these chemicals and how they regulate the operation of the brain.

A number of researchers studying rodent brains, including Anne Marie Thierry and Jacques Glowinski of the College of France in Paris, Brigitte Berger of Pitié Salpêtrière Hospital, also in Paris, and Tomas Hökfelt of the Karolinska Institute in Sweden, along with many colleagues, find that the prefrontal cortex abounds in catecholamines, a family of compounds that prepare the body for a stressful situation. Those compounds also act as neurotransmitters, substances that transmit neuronal impulses in the brain. My co-workers and I have discovered a similar abundance of catecholamines in the prefrontal cortices of nonhuman primates. One of the most familiar catecholamines, dopamine, regulates how neurons react to stimuli and seems to play a central role in schizophrenia.

A growing body of evidence suggests that dopamine is one of the most important of the chemicals that regulate cell activity associated with working memory. An imbalance in the abundance of dopamine in the prefrontal cortex can induce deficits in the working memory similar to those resulting from lesions in the principal sulcus region of the prefrontal cortex. For example, aged monkeys whose prefrontal cortices are deficient in dopamine and norepinephrine (a chemical relative of adrenaline) perform poorly in delayed-response tests. Injecting the aged animals with the deficient neurotransmitters restored their memory function

so that they tested roughly as well as younger, healthy monkeys.

Many of my colleagues and I are striving to learn which cells respond to dopamine and how they affect working memory. Within the past several years, we have collected evidence showing that neurons in certain layers of the cerebral cortex contain a great abundance of D₁ receptors, one of the chemical sites where dopamine binds to a cell. Interestingly, the neurons that are rich in D₁ receptors are those that project to the thalamus, the brain structure that relays information to the cortex.

Csaba Leranthy, John Smiley and F. Mark Williams of Yale are examining the cellular structures that enable dopamine to modulate responses to sensory inputs in the cerebral cortex. The researchers use an antibody developed by Michel Geffard of the Institute of Cellular Biochemistry and of Neurochemistry of the National Center of Scientific Research in Bordeaux, France, to label the neurons and their axonal projections that contain dopamine. They then scrutinize those cells under an electron microscope. The team looked in particular at the points of contact between dopamine-releasing cells and the neuronal spines, small protuberances where the cells receive incoming signals. Spines are discrete sites where calcium ions can enter and activate cellular mechanisms involving information processing and modulation of neuronal responses.

In most cases, the dopamine-releasing cells make symmetric contact with the spines—that is, the cell projections on either side of the synaptic cleft show roughly the same density. Such symmetric contacts are thought to have an inhibitory effect: when the postsynaptic site is activated, the cell's normal, spontaneous electrical activity is dampened. A large proportion of the spines of pyramidal cells—the major class of neuron that projects out of the cortex—receive asymmetric contacts from the axons of another cell whose point of origin has not yet been identified but which is thought to carry signals from other cortical areas. Those asymmetric contacts probably have an opposite, excitatory effect.

Pyramidal cells receive the major sensory or informational signals arriving at the cerebral cortex. The network of excitatory and inhibitory synapses, or connections, noted by the Yale group provides a mechanism by which dopamine could alter the way that various classes of pyramidal neurons respond to integrate such signals across thousands of spines in their dendrites. In this way, dopamine may regulate the overall output of the cortex. Further

analysis of the physical and chemical interactions between pyramidal cells and other neurons in the cerebral cortex should clarify how dopamine and other neurotransmitters influence cognition by stimulating or repressing the cellular responses of cortical neurons.

Investigations of the workings of the prefrontal cortex are revealing not only how the mind operates but also what goes wrong when it malfunctions. Medical researchers have implicated dysfunction of the prefrontal cortex as the cause of many neurological and psychiatric disorders, including Parkinson's disease and especially schizophrenia. The abnormal mental attributes associated with schizophrenia strongly resemble those caused by physical damage to the prefrontal cortex: thought disorders, reduced attention span, inappropriate or flattened emotional responses and lack of initiative, plans and goals. Schizophrenic patients, like frontal lobe patients and monkeys afflicted with prefrontal lobe lesions, retain a normal ability to perform routine procedures or habits but exhibit fragmented, disorganized behavior when attempting to perform tasks involving symbolic or verbal information [see "Major Disorders of Mind and Brain," by Elliot S. Gershon and Ronald O. Rieder, page 126].

Schizophrenic patients taking tests such as the Wisconsin Card Sort test tend to repeat a previous response even when it is clear that it is no longer the correct one; normal subjects, in contrast, shift hypotheses much sooner after making an error. Schizophrenic individuals are also severely impaired both on spatial delayed-response tasks and on a variety of tests of problem solving, abstraction and planning.

Studies of cerebral blood flow by David H. Ingvar of University Hospital in Lund, Sweden, and by Daniel R. Weinberger, Karen F. Berman and others at the National Institute of Mental Health, as well as measurements of local cerebral metabolism made by Monte S. Buchsbaum of the University of California at Irvine, show that schizophrenic patients have below-average blood flow into their prefrontal cortices, indicative of a depressed level of activity in that part of the brain. Schizophrenic subjects often suffer from impaired ability to move their eyes to track and project the forward trajectories of moving targets, further evidence that the disorder involves malfunctions in a posterior part of the prefrontal cortex, where the eye-movement centers involved in predictive tracking are located.

Sohee Park and Philip S. Holzman of


Harvard University have shown that schizophrenic subjects exhibit impaired performance on working memory tasks much like those my colleagues and I have used to study working memory in rhesus monkeys. Conversely, Martha MacAvoy and Bruce of Yale have demonstrated that monkeys with lesions in the relevant portions of the prefrontal cortex exhibit the same type of predictive tracking disorder that has long been considered a marker of schizophrenia in humans.

Perhaps researchers should begin to think of schizophrenia as a breakdown in the processes by which representational knowledge governs behavior. In my view, neural pathways in the prefrontal cortex update inner models of reality to reflect changing environmental demands and incoming information. Those pathways guide short-term memory and moment-to-moment behavior. If they fail, the brain views the world as a series of disconnected events, like a slide show, rather than as a continuous sequence, like a movie. The result is schizophrenic behavior, excessively dominated by immediate stimulation rather than by a balance of current, internal and past information.

At present, theories describing the fundamental causes of schizophrenia are inadequate, much as knowledge of the functioning of the working memory system remains frustratingly sketchy. Fortunately, neurobiological research has been advancing at a breathless pace in the past few years. Such research should lead to a greater understanding not only of schizophrenia but of the prefrontal cortex and how it shapes short-term memory and the broader working of the rational mind.

FURTHER READING

- WORKING MEMORY. Alan Baddeley. Oxford University Press, 1986.
- CIRCUITRY OF PRIMATE PREFRONTAL CORTEX AND REGULATION OF BEHAVIOR BY REPRESENTATIONAL MEMORY. P. S. Goldman-Rakic in *Handbook of Physiology*, Section 1, Vol. 5: *Higher Functions of the Brain*, Part 1. Edited by Fred Plum. Bethesda, Md., American Physiological Society, 1987.
- MNEMONIC CODING OF VISUAL SPACE IN THE MONKEY'S DORSOLATERAL PREFRONTAL CORTEX. Shintaro Funahashi, Charles J. Bruce and Patricia S. Goldman-Rakic in *Journal of Neurophysiology*, Vol. 61, No. 2, February 1989.
- PREFRONTAL CORTICAL DYSFUNCTION IN SCHIZOPHRENIA: THE RELEVANCE OF WORKING MEMORY. Patricia S. Goldman-Rakic in *Psychopathology and the Brain*. Edited by Bernard J. Carroll and James E. Barrett. Raven Press, 1991.



Sex Differences in the Brain

Cognitive variations between the sexes reflect differing hormonal influences on brain development. Understanding these differences and their causes can yield insights into brain organization

by Doreen Kimura

Women and men differ not only in physical attributes and reproductive function but also in the way in which they solve intellectual problems. It has been fashionable to insist that these differences are minimal, the consequence of variations in experience during development. The bulk of the evidence suggests, however, that the effects of sex hormones on brain organization occur so early in life that from the start the environment is acting on differently wired brains in girls and boys. Such differences make it almost impossible to evaluate the effects of experience independent of physiological predisposition.

Behavioral, neurological and endocrinologic studies have elucidated the processes giving rise to sex differences in the brain. As a result, aspects of the physiological basis for these variations have in recent years become clearer. In addition, studies of the effects of hormones on brain function throughout life suggest that the evolutionary pressures directing differences nevertheless allow for a degree of flexibility in cognitive ability between the sexes.

Major sex differences in intellectual function seem to lie in patterns of ability rather than in overall level of intelligence (IQ). We are all aware that people have different intellectual strengths. Some are especially good with words, others at using objects—for instance, at constructing or fixing things. In the same fashion, two individuals may have the same overall intelligence but have varying patterns of ability.

Men, on average, perform better than women on certain spatial tasks. In particular, men have an advantage in tests that require the subject to imagine rotating an object or manipulating it in some other way. They outperform women in mathematical reasoning tests and in navigating their way through a route. Further, men are more accurate in tests of target-directed motor skills—that is, in guiding or intercepting projectiles.

Women tend to be better than men at rapidly identifying matching items, a skill called perceptual speed. They have



greater verbal fluency, including the ability to find words that begin with a specific letter or fulfill some other constraint. Women also outperform men in arithmetic calculation and in recalling landmarks from a route. Moreover, women are faster at certain precision manual tasks, such as placing pegs in designated holes on a board.

Although some investigators have reported that sex differences in problem solving do not appear until after puberty, Diane Lunn, working in my laboratory at the University of Western Ontario, and I have found three-year-old boys to be better at targeting than girls of the same age. Moreover, Neil V. Watson, when in my laboratory, showed that the extent of experience playing sports does not account for the sex difference in targeting found in young adults. Kimberly A. Kerns, working with Sheri A. Berenbaum of the University of Chicago, has found that sex differences in spatial rotation performance are present before puberty.

Differences in route learning have been systematically studied in adults in laboratory situations. For instance, Liisa Galea in my department studied undergraduates who followed a route on a tabletop map. Men learned the route in fewer trials and made fewer errors than did women. But once learning was complete, women remembered more of the landmarks than did men. These results, and those of other researchers, raise the possibility that women tend to use landmarks as a strategy to orient themselves in everyday life. The prevailing strategies used by males have not yet been clearly established, although they must relate to spatial ability.

Marion Eals and Irwin Silverman of York University studied another function that may be related to landmark memory. The researchers tested the ability of individuals to recall objects and their locations within a confined space—such as in a room or on a tabletop. Women were better able to remember whether an item had been displaced or not. In addition, in my laboratory, we measured the accuracy of object location: subjects were shown an array of objects and were later asked to replace them in their exact positions. Women did so more accurately than did men.

ROUTES in a landscape such as the one in this painting, *The Old Oaken Bucket*, by Grandma Moses (1860–1961), may be learned differently by women and men. In laboratory experiments, researchers have found that women tend to remember landmarks—such as the well at the lower right or a tree at an intersection. Men appear to learn routes faster, but they cannot recall landmarks as readily: they may rely preferentially on spatial cues such as distance and direction.

DOREEN KIMURA studies the neural and hormonal basis of human intellectual function. She is professor of psychology and honorary lecturer in the department of clinical neurological sciences at the University of Western Ontario in London. Kimura, a fellow of the Royal Society of Canada, received the 1992 John Dewan Award for outstanding research from the Ontario Mental Health Foundation. She recently finished a book on neuromotor mechanisms in communication.

Problem-Solving Tasks Favoring Women

Women tend to perform better than men on tests of perceptual speed, in which subjects must rapidly identify matching items—for example, pairing the house on the far left with its twin:



In addition, women remember whether an object, or a series of objects, has been displaced:



On some tests of ideational fluency, for example, those in which subjects must list objects that are the same color, and on tests of verbal fluency, in which participants must list words that begin with the same letter, women also outperform men:

L _ _ _	Limp, Livery, Love, Laser, Liquid, Low, Like, Lag, Live Lug, Light, Lift, Liver, Lime, Leg, Load, Lap, Lucid ...
---------	---

Women do better on precision manual tasks—that is, those involving fine-motor coordination—such as placing the pegs in holes on a board:



And women do better than men on mathematical calculation tests:

77	$14 \times 3 - 17 + 52$
43	$2(15 + 3) + 12 - \frac{15}{3}$

It is important to place the differences described above in context: some are slight, some are quite large. Because men and women overlap enormously on many cognitive tests that show average sex differences, researchers use variations within each group as a tool to gauge the differences between groups.

Imagine, for instance, that on one test the average score is 105 for women and 100 for men. If the scores for women ranged from 100 to 110 and for men from 95 to 105, the difference would be more impressive than if the women's scores ranged from 50 to 150 and the men's from 45 to 145. In the latter case, the overlap in scores would be much greater.

One measure of the variation of scores within a group is the standard deviation. To compare the magnitude of a sex difference across several distinct tasks, the difference between groups is divided by the standard deviation. The resulting number is called the effect size. Effect sizes below 0.5 are generally considered small. Based on my data, for instance, there are typically no differences between the sexes on tests of vocabulary (effect size 0.02), nonverbal reasoning (0.03) and verbal reasoning (0.17).

On tests in which subjects match pictures, find words that begin with similar letters or show ideational fluency—such as naming objects that are white or red—the effect sizes are somewhat larger: 0.25, 0.22 and 0.38, respectively. As discussed above, women tend to outperform men on these tasks. Researchers have reported the largest effect sizes for certain tests measuring spatial rotation (effect size 0.7) and targeting accuracy (0.75). The large effect size in these tests means there are many more men at the high end of the score distribution.

Since, with the exception of the sex chromosomes, men and women share genetic material, how do such differences come about? Differing patterns of ability between men and women most probably reflect different hormonal influences on their developing brains. Early in life the action of estrogens and androgens (male hormones chief of which is testosterone) establishes sexual differentiation. In mammals, including humans, the organism has the potential to be male or female. If a Y chromosome is present, testes or male gonads form. This development is the critical first step toward becoming a male. If the gonads do not produce male hormones or if for some reason the hormones cannot act on the tissue, the default form of the organism is female.

Once testes are formed, they produce two substances that bring about the development of a male. Testosterone causes masculinization by promoting the male, or Wolffian, set of ducts and, indirectly through conversion to dihydrotestosterone, the external appearance of scrotum and penis. The Mülle-

rian regression factor causes the female, or Müllerian, set of ducts to regress. If anything goes wrong at any stage of the process, the individual may be incompletely masculinized.

Not only do sex hormones achieve the transformation of the genitals into male organs, but they also organize corresponding male behaviors early in life. Since we cannot manipulate the hormonal environment in humans, we owe much of what we know about the details of behavioral determination to studies in other animals. Again, the intrinsic tendency, according to studies by Robert W. Goy of the University of Wisconsin, is to develop the female pattern that occurs in the absence of masculinizing hormonal influence.

If a rodent with functional male genitals is deprived of androgens immediately after birth (either by castration or by the administration of a compound that blocks androgens), male sexual behavior, such as mounting, will be reduced. Instead female sexual behavior, such as lordosis (arching of the back), will be enhanced in adulthood. Similarly, if androgens are administered to a female directly after birth, she displays more male sexual behavior and less female behavior in adulthood.

Bruce S. McEwen and his co-workers at the Rockefeller University have shown that, in the rat, the two processes of defeminization and masculinization require somewhat different biochemical changes. These events also occur at somewhat different times. Testosterone can be converted to either estrogen (usually considered a female hormone) or dihydrotestosterone. Defeminization takes place primarily after birth in rats and is mediated by estrogen, whereas masculinization involves both dihydrotestosterone and estrogen and occurs for the most part before birth rather than after, according to studies by McEwen. A substance called alpha-fetoprotein may protect female brains from the masculinizing effects of their estrogen.

The area in the brain that organizes female and male reproductive behavior is the hypothalamus. This tiny structure at the base of the brain connects to the pituitary, the master endocrine gland. Roger A. Gorski and his colleagues at the University of California at Los Angeles have shown that a region of the pre-optic area of the hypothalamus is visibly larger in male rats than in females. The size increment in males is promoted by the presence of androgens in the immediate postnatal, and to some extent prenatal, period. Laura S. Allen in Gorski's laboratory has found a similar sex difference in the human brain.

Other preliminary but intriguing studies suggest that sexual behavior may reflect further anatomic differences. In 1991 Simon LeVay of the Salk Institute for Biological Studies in San Diego reported that one of the brain regions that is usually larger in human males than in females—an interstitial nucleus of the anterior hypothalamus—is smaller in homosexual than in heterosexual men. LeVay points out that this finding supports suggestions that sexual preference has a biological substrate.

Homosexual and heterosexual men may also perform differently on cognitive tests. Brian A. Gladue of North Dakota State University and Geoff D. Sanders of City of London Polytechnic report that homosexual men perform less well on several spatial tasks than do heterosexual men. In a recent study in my laboratory, Jeff Hall found that homosexual men had lower scores on targeting tasks than did heterosexual men; however, they were superior in ideational fluency—listing things that were a particular color.

This exciting field of research is just starting, and it is crucial that investigators consider the degree to which differences in life-style contribute to group differences. One should also keep in mind that results concerning group differences constitute a general statistical statement; they establish a mean from which any individual may differ. Such studies are potentially a rich source of information on the physiological basis for cognitive patterns.

The lifelong effects of early exposure to sex hormones are characterized as organizational, because they appear to alter brain function permanently during a critical period. Administering the same hormones at later stages has no such effect. The hormonal effects are not limited to sexual or reproductive behaviors: they appear to extend to all known behaviors in which males and females differ. They seem to govern problem solving, aggression and the tendency to engage in rough-and-tumble play—the boisterous body contact that young males of some mammalian species display. For example, Michael J. Meaney of McGill University finds that dihydrotestosterone, working through a structure called the amygdala rather than through the hypothalamus, gives rise to the play-fighting behavior of juvenile male rodents.

Male and female rats have also been found to solve problems differently. Christina L. Williams of Barnard College has shown that female rats have a greater tendency to use landmarks

in spatial learning tasks—as it appears women do. In Williams's experiment, female rats used landmark cues, such as pictures on the wall, in preference to geometric cues, such as angles and the shape of the room. If no landmarks were available, however, females used geometric cues. In contrast, males did not use landmarks at all, preferring geometric cues almost exclusively.

Interestingly, hormonal manipulation during the critical period can alter these behaviors. Depriving newborn males of testosterone by castrating them or administering estrogen to newborn females results in a complete reversal of sex-typed behaviors in the adult animals. (As mentioned above, estrogen can have a masculinizing effect during brain development.) Treated females behave like males, and treated males behave like females.

Natural selection for reproductive advantage could account for the evolution of such navigational differences. Steven J. C. Gaulin and Randall W. FitzGerald of the University of Pittsburgh have suggested that in species of voles in which a male mates with several females rather than with just one, the range he must traverse is greater. Therefore, navigational ability seems critical to reproductive success. Indeed, Gaulin and FitzGerald found sex differences in laboratory maze learning only in voles that were polygynous, such as the meadow vole, not in monogamous species, such as the prairie vole.


Again, behavioral differences may parallel structural ones. Lucia F. Jacobs in Gaulin's laboratory has discovered that the hippocampus—a region thought to be involved in spatial learning in both birds and mammals—is larger in male polygynous voles than in females. At present, there are no data on possible sex differences in hippocampal size in human subjects.

Evidence of the influence of sex hormones on adult behavior is less direct in humans than in other animals. Researchers are instead guided by what may be parallels in other species and by spontaneously occurring exceptions to the norm in humans.


One of the most compelling areas of evidence comes from studies of girls exposed to excess androgens in the prenatal or neonatal stage. The production of abnormally large quantities of adrenal androgens can occur because of a genetic defect called congenital adrenal hyperplasia (CAH). Before the 1970s, a similar condition also unexpectedly appeared when pregnant women took various synthetic steroids. Although the consequent masculinization of the geni-

Problem-Solving Tasks Favoring Men


Men tend to perform better than women on certain spatial tasks. They do well on tests that involve mentally rotating an object or manipulating it in some fashion, such as imagining turning this three-dimensional object



or determining where the holes punched in a folded piece of paper will fall when the paper is unfolded:




Men also are more accurate than women in target-directed motor skills, such as guiding or intercepting projectiles:



They do better on disembedding tests, in which they have to find a simple shape, such as the one on the left, once it is hidden within a more complex figure:




And men tend to do better than women on tests of mathematical reasoning:




tals can be corrected early in life and drug therapy can stop the overproduction of androgens, effects of prenatal exposure on the brain cannot be reversed.

Studies by researchers such as Anke A. Ehrhardt of Columbia University and June M. Reinisch of the Kinsey Institute have found that girls with excess expo-


SPATIAL



MATHEMATICAL REASONING



PERCEPTUAL SPEED



TESTOSTERONE LEVELS can affect performance on some tests (see boxes on pages 120 and 121 for examples of tests). Women with high levels of testosterone perform better on a spatial task (*top*) than do women with low levels; men with low levels outperform men with high levels. On a mathematical reasoning test (*middle*), low testosterone corresponds to better performance in men; in women there is no such relation. On a test in which women usually excel (*bottom*), no relation is found between testosterone and performance.

sure to androgens grow up to be more tomboyish and aggressive than their unaffected sisters. This conclusion was based sometimes on interviews with subjects and mothers, on teachers' ratings and on questionnaires administered to the girls themselves. When ratings are used in such studies, it can be difficult to rule out the influence of expectation either on the part of an adult who knows the girls' history or on the part of the girls themselves.

Therefore, the objective observations of Berenbaum are important and convincing. She and Melissa Hines of the University of California at Los Angeles observed the play behavior of CAH-affected girls and compared it with that of their male and female siblings. Given a

choice of transportation and construction toys, dolls and kitchen supplies or books and board games, the CAH girls preferred the more typically masculine toys—for example, they played with cars for the same amount of time that normal boys did. Both the CAH girls and the boys differed from unaffected girls in their patterns of choice. Because there is every reason to think that parents would be at least as likely to encourage feminine preferences in their CAH daughters as in their unaffected daughters, these findings suggest that the toy preferences were actually altered in some way by the early hormonal environment.

Spatial abilities that are typically better in males are also enhanced in CAH girls. Susan M. Resnick, now at the Na-

tional Institute on Aging, and Berenbaum and their colleagues reported that affected girls were superior to their unaffected sisters in a spatial manipulation test, two spatial rotation tests and a disembedding test—that is, the discovery of a simple figure hidden within a more complex one. All these tasks are usually done better by males. No differences existed between the two groups on other perceptual or verbal tasks or on a reasoning task.

Studies such as these suggest that the higher the androgen levels, the better the spatial performance. But this does not seem to be the case. In 1983 Valerie J. Shute, when at the University of California at Santa Barbara, suggested that the relation between levels of androgens and some spatial capabilities might be nonlinear. In other words, spatial ability might not increase as the amount of androgen increases. Shute measured androgens in blood taken from male and female students and divided each into high- and low-androgen groups. All fell within the normal range for each sex (androgens are present in females but in very low levels). She found that in women, the high-androgen subjects were better at the spatial tests. In men the reverse was true: low-androgen men performed better.

Catherine Gouchie and I recently conducted a study along similar lines by measuring testosterone in saliva. We added tests for two other kinds of abilities: mathematical reasoning and perceptual speed. Our results on the spatial tests were very similar to Shute's: low-testosterone men were superior to high-testosterone men, but high-testosterone women surpassed low-testosterone women. Such findings suggest some optimum level of androgen for maximal spatial ability. This level may fall in the low male range.

No correlation was found between testosterone levels and performance on perceptual speed tests. On mathematical reasoning, however, the results were similar to those of spatial ability tests for men: low-androgen men tested higher, but there was no obvious relation in women.

Such findings are consistent with the suggestion by Camilla P. Benbow of Iowa State University that high mathematical ability has a significant biological determinant. Benbow and her colleagues have reported consistent sex differences in mathematical reasoning ability favoring males. These differences are especially sharp at the upper end of the distribution, where males outnumber females 13 to one. Benbow argues

that these differences are not readily explained by socialization.

It is important to keep in mind that the relation between natural hormonal levels and problem solving is based on correlational data. Some form of connection between the two measures exists, but how this association is determined or what its causal basis may be is unknown. Little is currently understood about the relation between adult levels of hormones and those in early life, when abilities appear to be organized in the nervous system. We have a lot to learn about the precise mechanisms underlying cognitive patterns in people.

Another approach to probing differences between male and female brains is to examine and compare the functions of particular brain systems. One noninvasive way to accomplish this goal is to study people who have experienced damage to a specific brain region. Such studies indicate that the left half of the brain in most people is critical for speech, the right for certain perceptual and spatial functions.


It is widely assumed by many researchers studying sex differences that the two hemispheres are more asymmetrically organized for speech and spatial functions in men than in women. This idea comes from several sources. Parts of the corpus callosum, a major neural system connecting the two hemispheres, may be more extensive in women; perceptual techniques that probe brain asymmetry in normal-functioning people sometimes show smaller asymmetries in women than in men, and damage to one brain hemisphere sometimes has a lesser effect in women than the comparable injury has in men.

In 1982 Marie-Christine de Lacoste, now at the Yale University School of Medicine, and Ralph L. Holloway of Columbia University reported that the back part of the corpus callosum, an area called the splenium, was larger in women than in men. This finding has subsequently been both refuted and confirmed. Variations in the shape of the corpus callosum that may occur as an individual ages as well as different meth-

ods of measurement may produce some of the disagreements. Most recently, Allen and Gorski found the same sex-related size difference in the splenium.

The interest in the corpus callosum arises from the assumption that its size may indicate the number of fibers connecting the two hemispheres. If more connecting fibers existed in one sex, the implication would be that in that sex the hemispheres communicate more fully. Although sex hormones can alter callosal size in rats, as Victor H. Denenberg and his associates at the University of Connecticut have demonstrated, it is unclear whether the actual number of fibers differs between the sexes. Moreover, sex differences in cognitive function have yet to be related to a difference in callosal size. New ways of imaging the brain in living humans will undoubtedly increase knowledge in this respect.

The view that a male brain is functionally more asymmetric than a female brain is long-standing. Albert M. Galaburda of Beth Israel Hospital in Boston and the late Norman Geschwind of Har-



RIGHT HEMISPHERIC DAMAGE affects spatial ability to the same degree in both sexes (*graphs at bottom*), suggesting that women and men rely equally on that hemisphere for certain spatial tasks. In one test of spatial rotation performance

(*left*), a series of drawings of a gloved right or left hand must be matched to a right- or left-handed glove. In a second test (*right*), photographs of a three-dimensional object must be matched to one of two mirror images of the same object.

vard Medical School proposed that androgens increased the functional potency of the right hemisphere. In 1981 Marian C. Diamond of the University of California at Berkeley found that the right cortex is thicker than the left in male rats but not in females. Jane Stewart of Concordia University in Montreal, working with Bryan E. Kolb of the University of Lethbridge in Alberta, recently pinpointed early hormonal influences on this asymmetry: androgens appear to suppress left cortex growth.

Last year de Lacoste and her colleagues reported a similar pattern in human fetuses. They found the right cortex was thicker than the left in males. Thus, there appear to be some anatomic reasons for believing that the two hemispheres might not be equally asymmetric in men and women.


Despite this expectation, the evidence in favor of it is meager and conflicting, which suggests that the most striking sex differences in brain organization may not be related to asymmetry. For example, if overall differences between men and women in spatial ability were related to differing right hemispheric dependence for such functions, then damage to the right hemisphere would perhaps have a more devastating effect on spatial performance in men.

My laboratory has recently studied the ability of patients with damage to one hemisphere of the brain to rotate certain objects mentally. In one test, a series of line drawings of either a left or a right gloved hand is presented in various orientations. The patient indicates the hand being depicted by simply pointing to one of two stuffed gloves that are constantly present.

The second test uses two three-dimensional blocklike figures that are mirror images of one another. Both figures are present throughout the test. The patient is given a series of photographs of these objects in various orientations, and he or she must place each picture in front of the object it depicts. (These nonverbal procedures are employed so that patients with speech disorders can be tested.)

As expected, damage to the right hemisphere resulted in lower scores for both sexes on these tests than did damage to the left hemisphere. Also as anticipated, women did less well than men on the block spatial rotation test. Surprisingly, however, damage to the right hemisphere had no greater effect in men than in women. Women were at least as affected as men by damage to the right hemisphere. This result suggests that the normal differences between men and women on such rotational tests are

INCIDENCE OF APHASIA



not the result of differential dependence on the right hemisphere. Some other brain systems must be mediating the higher performance by men.

Parallel suggestions of greater asymmetry in men regarding speech have rested on the fact that the incidence of aphasias, or speech disorders, are higher in men than in women after damage to the left hemisphere. Therefore, some researchers have found it reasonable to conclude that speech must be more bilaterally organized in women. There is, however, a problem with this conclusion. During my 20 years of experience with patients, aphasia has not been disproportionately present in women with right hemispheric damage.

In searching for an explanation, I discovered another striking difference between men and women in brain organization for speech and related motor function. Women are more likely than men to suffer aphasia when the front part of the brain is damaged. Because restricted damage within a hemisphere more frequently affects the posterior than the anterior area in both men and women, this differential dependence may explain why women incur aphasia less often than do men. Speech functions are thus less likely to be affected in women not because speech is more bilaterally organized in women but because the critical area is less often affected.

A similar pattern emerges in studies of the control of hand movements, which are programmed by the left hemisphere. Apraxia, or difficulty in selecting appropriate hand movements, is very


APHASIAS, or speech disorders, occur most often in women when damage is to the front of the brain. In men, they occur more frequently when damage is in the posterior region. The data presented above derive from one set of patients.

common after left hemispheric damage. It is also strongly associated with difficulty in organizing speech. In fact, the critical functions that depend on the left hemisphere may relate not to language per se but to organization of the complex oral and manual movements on which human communication systems depend. Studies of patients with left hemispheric damage have revealed that such motor selection relies on anterior systems in women but on posterior systems in men.

The synaptic proximity of women's anterior motor selection system (or "praxis system") to the motor cortex directly behind it may enhance fine-motor skills. In contrast, men's motor skills appear to emphasize targeting or directing movements toward external space—some distance away from the self. There may be advantages to such motor skills when they are closely meshed with visual input to the brain, which lies in the posterior region.

Women's dependence on the anterior region is detectable even when tests involve using visual guidance—for instance, when subjects must build patterns with blocks by following a visual model. In studying such a complex task, it is possible to compare the effects of damage to the anterior and posterior regions of both hemispheres because per-

INCIDENCE OF APRAXIA



APRAXIA, or difficulty in selecting hand movements, is associated with frontal damage to the left hemisphere in women and with posterior damage in men. It is also associated with difficulties in organizing speech.

formance is affected by damage to either hemisphere. Again, women prove more affected by damage to the anterior region of the right hemisphere than by posterior damage. Men tend to display the reverse pattern.

Although I have not found evidence of sex differences in functional brain asymmetry with regard to basic speech, motor selection or spatial rotation ability, I have found slight differences in more abstract verbal tasks. Scores on a vocabulary test, for instance, were affected by damage to either hemisphere in women, but such scores were affected only by left-sided injury in men. This finding suggests that in reviewing the meanings of words, women use the hemispheres more equally than do men.

In contrast, the incidence of non-right-handedness, which is presumably related to lesser left hemispheric dependence, is higher in men than in women. Even among right-handers, Marion Annett, now at the University of Leicester in the U.K., has reported that women are more right-handed than men—that is, they favor their right hand even more than do right-handed men. It may well be, then, that sex differences in asymmetry vary with the particular function being studied and that it is not always the same sex that is more asymmetric.

Taken altogether, the evidence sug-

gests that men's and women's brains are organized along different lines from very early in life. During development, sex hormones direct such differentiation. Similar mechanisms probably operate to produce variation within sexes, since there is a relation between levels of certain hormones and cognitive makeup in adulthood.

One of the most intriguing findings is that cognitive patterns may remain sensitive to hormonal fluctuations throughout life. Elizabeth Hampson of the University of Western Ontario showed that the performance of women on certain tasks changed throughout the menstrual cycle as levels of estrogen went up or down. High levels of the hormone were associated not only with relatively depressed spatial ability but also with enhanced articulatory and motor capability.

In addition, I have observed seasonal fluctuations in spatial ability in men. Their performance is improved in the spring when testosterone levels are lower. Whether these intellectual fluctuations are of any adaptive significance or merely represent ripples on a stable baseline remains to be determined.

To understand human intellectual functions, including how groups may differ in such functions, we need to look beyond the demands of modern life. We did not undergo natural selection for reading or for operating computers. It seems clear that the sex differences in cognitive patterns arose because they proved evolutionarily advantageous. And their adaptive significance probably rests in the distant past. The orga-

nization of the human brain was determined over many generations by natural selection. As studies of fossil skulls have shown, our brains are essentially like those of our ancestors of 50,000 or more years ago.

For the thousands of years during which our brain characteristics evolved, humans lived in relatively small groups of hunter-gatherers. The division of labor between the sexes in such a society probably was quite marked, as it is in existing hunter-gatherer societies. Men were responsible for hunting large game, which often required long-distance travel. They were also responsible for defending the group against predators and enemies and for the shaping and use of weapons. Women most probably gathered food near the camp, tended the home, prepared food and clothing and cared for children.

Such specializations would put different selection pressures on men and women. Men would require long-distance route-finding ability so they could recognize a geographic array from varying orientations. They would also need targeting skills. Women would require short-range navigation, perhaps using landmarks, fine-motor capabilities carried on within a circumscribed space, and perceptual discrimination sensitive to small changes in the environment or in children's appearance or behavior.


The finding of consistent and, in some cases, quite substantial sex differences suggests that men and women may have different occupational interests and capabilities, independent of societal influences. I would not expect, for example, that men and women would necessarily be equally represented in activities or professions that emphasize spatial or math skills, such as engineering or physics. But I might expect more women in medical diagnostic fields where perceptual skills are important. So that even though any one individual might have the capacity to be in a "nontypical" field, the sex proportions as a whole may vary.

FURTHER READING

SEX DIFFERENCES IN THE BRAIN: THE RELATION BETWEEN STRUCTURE AND FUNCTION. Edited by G. J. DeVries, J.P.C. DeBruin, H.B.M. Uylings and M. A. Corner in *Progress in Brain Research*, Vol. 61. Elsevier, 1984.

MASCULINITY/FEMININITY. Edited by J. M. Reinisch, L. A. Rosenblum and S. A. Sanders. Oxford University Press, 1987.

BEHAVIORAL ENDOCRINOLOGY. Edited by Jill B. Becker, S. Marc Breedlove and David Crews. The MIT Press/Bradford Books, 1992.



Major Disorders of Mind and Brain

Schizophrenia and manic-depressive illness are shaped by heredity and marked by structural and biochemical changes in the brain. The predisposing genes remain unknown


by Elliot S. Gershon and Ronald O. Rieder

Madness was understood for centuries by religion and poetry as an affliction of the spirit and by medicine as a disorder of various humors and organs of the body. In the past century, physicians have recognized the most common forms of psychosis (our current word for madness) as two chronic disorders—schizophrenia and mania—and have begun to understand the abnormalities in brain structure and function that accompany them. Each affects about 1 percent of the population. Both flare episodically, although schizophrenia follows a deteriorating course, whereas patients with bipolar manic-depressive illness, who have episodes of mania and depression, are usually mentally normal between episodes.

The anatomic, biochemical and hereditary bases of these disorders are now emerging. Some research has already shaped the development of new treatments. These subjects form the focus of our article. First, however, it is useful to consider what these disorders are like for the people who have them.

When Mrs. T. was 16 years old, she began to experience her first symptom of schizophrenia: a profound feeling that people were staring at her. These bouts of self-consciousness soon forced her to end her public piano performances. Her self-consciousness led to withdrawal, then to fearful delusions that others were speaking of her and finally to suspicions that they were plotting to harm her. At first Mrs. T.'s illness was intermittent, and the return of her intelligence, warmth and ambition between episodes allowed her to complete several years of college, to marry and to rear three children. She had to enter a hospital for the first time at 28, after the birth of her third child, when she began to hallucinate.

Now, at 45, Mrs. T. is never entirely well. She has seen dinosaurs on the street and live animals in her refrigerator. While hallucinating, she speaks and writes in an incoherent, but almost poetic, way. At other times, she is more lucid, but even then her voices sometimes lead her to do dangerous things, such as driving very fast down the highway in the middle of the night, dressed only in a nightgown. As an episode winds down, Mrs. T. usually becomes deeply depressed and hopeless about her condition. Often she sits in her car



with the engine running and contemplates committing suicide.

Over the past five years she has taken antipsychotic medications, such as haloperidol, that suppress the hallucinations and help her stay out of the hospital. Stress, however, can bring the hallucinations and delusions back for days or weeks, as happened after her recent separation from her husband and the subsequent sale of her home. At such times, her voices shout terrible criticisms. After her daughter left for college, they shouted, "You'll


never see her again, you have been a bad mother, she'll die." At other times and without any apparent stimulus, Mrs. T. has bizarre visual hallucinations. For example, she saw cherubs in the grocery store. These experiences leave her preoccupied, confused and frightened, unable to perform such everyday tasks as cooking or playing the piano. When feeling well, however, she does volunteer work at church.

The mood disorders, which are distinct from schizophrenia, are called unipolar when the patient has episodes of depression alone and bipolar when there are episodes of both mania and depression. (The term "manic-depressive illness" encompasses both the unipolar and the bipolar form; the term "bipolar" is also used for the rare cases in which mania occurs without depression.) The depressions are quite severe, and suicide is an all too frequent outcome. Mania, a state of excitement usually characterized by impulsive behavior, can, when untreated, ultimately ruin marriages, careers and fortunes.

Mania can develop suddenly and shockingly, as illustrated in a case cited by a group led by Robert L. Spitzer of Columbia University. Daryl, a 25-year-old dancer, was cast in a

HALLUCINATORY SELF-PORTRAIT exemplifies the visual creativity that schizophrenia sometimes induces; it also conveys the pain and perceptual distortions of this illness.

ELLIOT S. GERSHON and RONALD O. RIEDER began collaborating on the study of mental illness at the Intramural Research Program of the National Institute of Mental Health. Gershon heads the program's Clinical Neurogenetics Branch and specializes in the population genetics and molecular genetics of normal and abnormal behavior. He graduated from Harvard Medical School in 1965, trained in psychiatry at the Massachusetts Mental Health Center and moved to the NIMH in 1969. Rieder graduated from Harvard Medical School in 1968 and trained in psychiatry at Albert Einstein College of Medicine. At the NIMH he conducted research on schizophrenia. Now he directs research and psychiatric residency training at Columbia University.



STRUCTURAL CHANGES appear in the shrunken hippocampus (yellow) and enlarged, fluid-filled ventricles (gray) of the brain of a schizophrenic patient (top), as contrasted with that of a normal volunteer (bottom). These three-dimensional MRI reconstructions were made by Nancy C. Andreasen of the University of Iowa.

part for a Broadway show. He began to come home at the end of rehearsal making disparaging remarks about the sessions and the director. A week later a fellow performer called Daryl's wife to complain that her husband had been trying to take over the rehearsals, giving unsolicited advice to the director and the other performers. At this point, his wife realized that Daryl's usually

easygoing demeanor had turned tense and irritable. He began making nasty comments about his wife's figure and their recent sex life. Three days later he began shouting obscenities at other performers and was ejected from the theater. At home, he talked "a mile a minute" and paced incessantly, dressed only in his underwear. He felt no need to eat or sleep. The next day he skipped

work to make a number of extravagant purchases.

At this time, two weeks after he had displayed his first symptoms, Daryl accepted hospitalization. He received one dose of a tranquilizer yet spent most of the night disrupting the ward. Then he signed out, against medical advice, in the morning. Eventually he responded well to lithium carbonate. Daryl's father has had a similar but more prolonged history, losing many jobs over 20 years after episodes of excited confrontations with his bosses. Over the past five years, however, he too has responded well to lithium.

Although schizophrenia and manic-depression can devastate patients' lives, the disorders do not preclude the performance of highly creative work. Schizophrenic patients confined in institutions have occasionally produced extraordinary works of graphic art [see *illustration on page 126*]. Manic-depressive illness often occurs in conjunction with extraordinary talent, even genius, in politics and military leadership, as well as in literature and music and the other performing arts. Among those thought to have had the disorder are William Blake, Lord Byron, Virginia Woolf, Robert Schumann, Oliver Cromwell and Winston Churchill. Many observers have suggested that extremes of mood and changes in outlook may spur creativity; they also speculate that the energy and facility of thought that typify the milder stages of mania can be a source of creativity.

Even though schizophrenia and severe mood disorders manifest themselves as intangible mental experiences, they are biologically determined to a major degree. (Only a few of the biological discoveries can be discussed in this brief article.) The first evidence of determinants came early in this century, when genetic studies showed that both schizophrenia and manic-depressive illness ran in families. Most workers discounted these correlations, however, on the grounds that families share environment as well as genes. To consider the two factors in isolation, researchers turned to adoptees, who, once adopted, have environmental families that are different from their genetic families.

In the best-known study, begun in the 1960s, Seymour S. Kety and his colleagues at the National Institute of Mental Health and at a psychological institute in Scandinavia identified schizophrenics adopted in infancy and traced their biological relatives through the adoption register. The study indicated that biological relatives had an

increased risk of developing the illness—es but adoptive relatives did not. The control group—biological relatives of nonpsychotic adoptees—faced no excess risk of schizophrenia or of any mental illness resembling it.

Twin studies are also revealing because different types of twins vary widely in their genetic relatedness. When schizophrenia or bipolar disease develops in one twin, the chance that it will develop in the other is much greater in identical twins, who share all their genes, than in fraternal twins, who share only about half. Moreover, although about half of the identical twins of schizophrenics never develop the illness, the children of even the well twins are at increased risk. These correlations imply two things. The risk of illness rises with increasing genetic similarity, but even a perfect identity of genes does not produce a perfect correspondence. Some environmental factor, or interaction of genes with the environment, must therefore push susceptible people over the threshold of illness. Studies have already implicated one possible factor: prenatal exposure to the influenza virus.

Mood disorders also stem from the interaction of genes with some aspect of the environment. Rates of major depression in every age group have steadily increased in several of the developed countries since the 1940s. This trend was first spotted some 10 years ago in an epidemiological study in Sweden. A similar increase in suicide over the same four decades occurred in Alberta, Canada. These findings have been firmly established as birth-cohort effects: suicide rates among 15- to 19-year-olds, for instance, were 10 times higher for those born in the late 1950s than for those born in the early 1930s. Similar birth-cohort increases appeared, over these decades, in suicide and unipolar disorder in the U.S., in bipolar disorder in the U.S. and Switzerland, and in alcoholism in males in the U.S. [see illustration on page 131].

Rates of depression, mania and suicide continue to rise as each new birth cohort ages, a pattern that harbors ominous public health consequences. Such birth-cohort effects are even more pronounced in the relatives of patients than in the general population—in other words, at comparable ages, the children of patients are far more susceptible to these disorders than are their ill parents' siblings. This relation clearly implies an interaction between genes and some environmental factor, which must have been changing continuously over the past few decades. The factor remains a mystery.

The biological abnormalities that genes and environment somehow put into motion were quite mysterious until the 1970s, when new imaging technologies allowed physicians to visualize the living brain in great detail.

One imaging technique, computerized tomography (CT scanning), was first applied to the brains of schizophrenic patients in 1978 by Eve C. Johnstone and her colleagues at the Clinical Research Centre in Middlesex, England. They observed that the lateral cerebral ventricles were much larger than in normal subjects. If the ventricles or the spaces between convolutions are enlarged, one can conclude there has been a failure of development or a loss of brain tissue. Other x-ray evidence confirmed this conclusion by showing less tissue and more fluid-filled spaces around the convolutions in the cerebral cortex.


Another technique, magnetic resonance imaging (MRI), confirmed the ventricular enlargement. Daniel R. Weinberger's group at the National Institute of Mental Health used MRI to compare identical twins in which one twin had schizophrenia and the other did not. In 12 of 15 sets of such twins, the schizophrenic one had the larger cerebral ventricles. Relative diminution of specific brain structures has been demonstrated, too, in autopsies and by MRI scans of schizophrenic patients. The most striking examples of such diminution appear in the hippocampal region, part of the limbic system in the temporal lobe of the cerebrum, which modulates emotional response, memory and other functions [see illustration on opposite page].

The new imaging devices also showed functional abnormalities for the first time. In 1974 David H. Ingvar of University Hospital in Lund, Sweden, found reduced blood flow in the frontal cerebrum of schizophrenic patients, implying decreased neuronal activity there. This finding has since been corroborated many times [see illustration at right].

Weinberger's group presents evidence linking both structural and functional abnormalities in the brain to a schizophrenic cognitive trait. They found that normal subjects show increased blood flow in the prefrontal cerebral cortex while taking the Wisconsin Card Sort, a test of working memory and abstract thinking, whereas schizophrenic subjects show less of an increase in flow and do worse on the test. Moreover, those schizophrenic patients whose hippocampal structures are the smallest show the greatest deficit in prefrontal blood flow. The hippocampus connects to the prefrontal cortex,

which manages working memory in primates [see "Working Memory and the Mind," by Patricia S. Goldman-Rakic, page 110].

Postmortem studies of schizophrenic patients have also uncovered abnormalities in the number of brain cells and in their organization, particularly in the temporal lobe. Yet the tissue shows none of the scarring one would expect from an infection, nor do the abnormalities progress over time. Some researchers therefore speculate that the abnormalities stem from a developmental disorder, perhaps a failure of the growth of neurons and the development of their connections, or from a disturbance in the "pruning" of neurons that normally




FUNCTIONAL DEFICIENCY is evident in positron emission tomographic scans made while the subjects were taking a test of vigilance. The test enhances prefrontal cortex metabolism in the normal volunteer (*top*) but not in the schizophrenic patient (*bottom*). The data were provided by Monte S. Buchsbaum of the University of California at Irvine.

Medicines for Mental Disorders

Drugs may act on several points in the synapse. Antidepressants that affect the presynaptic cell include those that block the cell's reuptake of monoamines (a). These drugs include tricyclic antidepressants, such as imipramine, which block reuptake of several monoamines, and more specific blockers, such as fluoxetine, for serotonin, and bupropion, for dopamine. Other antidepressants known as monoamine oxidase inhibitors (b) prevent the presynaptic cell from metabolizing monoamines. Drugs that affect the postsynaptic cell include agents that either block monoamine receptors or stimu-

late their ability to respond (c). Haloperidol, an antipsychotic, is a dopamine receptor blocker. Finally, some drugs affect the second messenger (d) that is normally produced after a receptor has been activated. For example, lithium carbonate, an antidepressant and antimanic agent, works by inhibiting the synthesis of phosphatidyl inositol. Here a postsynaptic receptor is shown coupled to a stimulatory G protein; this is its activated state, which causes more second-messenger chemicals to be synthesized, triggering molecular cascades that determine how the postsynaptic cell will respond.



occurs between the ages of three and 15 [see "The Developing Brain," by Carla J. Shatz, page 60].

How might such abnormalities cause the symptoms of schizophrenia? While conducting surgery on nonschizophrenic patients, Wilder G. Penfield of the Montreal Neurological Institute discovered that certain structures in the brain are related to hallucinations. (Such patients are often kept awake so that they can help ascertain the functions of brain tissues near the field in which the surgeon is operating.) Penfield found that when he touched his diagnostic electrode to the temporal lobe he often elicited in his patients sights and sounds resembling hallucinations.

Further research showed that the frontal parts of the temporal cortex receive highly processed and filtered sensory information from other parts of the cortex. That information eventually reaches the limbic system and other structures that mediate emotional response, or affect. Perhaps, then, some overactivation of the temporal cortex or abnormalities in the filtering process produce the common experiences of schizophrenic patients: auditory hallucinations and a sense of being overwhelmed by all the senses.

The first effective medications for schizophrenia and depression were discovered serendipitously, without any knowledge of their effects on brain

chemistry. Chlorpromazine was developed in the 1950s as a surgical anesthetic but turned out to alleviate the symptoms of both schizophrenia and mania. It thus became the first widely used antipsychotic drug. Scientists then used it as a model for the synthesis of imipramine, which they expected would also serve as an antipsychotic agent. Instead it turned out to be very effective in the treatment of depression. Lithium was introduced into the treatment of manic-depressive illness after John Cade, an Australian psychiatrist, noted in 1949 that lithium salts sedated rodents in his laboratory.

Insight into the way antidepressant agents act began with the study of reser-

pine, a drug derived from the plant *Rauwolfia serpentina*, used in traditional medicine in India. Reserpine was one of the first effective medications for high blood pressure. Physicians noted, however, that the drug sometimes brought on severe depression in patients—a few even committed suicide.

Biochemists discovered that reserpine depletes certain neurotransmitters classed as monoamines, among them norepinephrine, dopamine and serotonin. All the antidepressant drugs known in the mid-1960s effectively concentrated these monoamines in the synapse, either by inhibiting their metabolic breakdown or by preventing cells from reabsorbing them from the synaptic space (a process known as reuptake).

This pattern led Joseph J. Schildkraut, then at the National Institute of Mental Health, to propose in 1965 that depression was associated with a reduction in synaptic availability of catecholamines (norepinephrine and dopamine), particularly dopamine, and mania with an increase of catecholamines. Nevertheless, there are antidepressant drugs, such as iprindole, that are associated with no observable change in norepinephrine reuptake or metabolism.

Biochemical pharmacologists therefore looked beyond the neurotransmitters to the synaptic receptor molecules that bind with them. The workers knew that norepinephrine has several pharmacologically distinct receptors, called adrenoceptors, but their experiments in binding various antidepressant drugs to the receptors produced no consistent change.

Then, in 1975, Fridolin Sulser's laboratory at Vanderbilt University found an answer by looking not at binding itself but at the intracellular response that one type of binding elicits. They studied how norepinephrine stimulates beta-receptors, a class of adrenoceptors that mediates the release of cyclic adenosine monophosphate inside the nerve cell. This molecule then serves as a second chemical messenger. But after long-term administration of certain antidepressants, including iprindole, this secondary response consistently decreases. Virtually all antidepressants, including those discovered after the finding was published, produce this result. So does electroconvulsive therapy, a very effective treatment for depression in which shocks are administered to the cerebrum to induce artificial seizures.

A number of receptors for each of the monoamines exist, and new receptors continue to be discovered. With each discovery, workers learn more about how antidepressants alter these receptors and their second-messenger sys-

tems. It turns out that many, but not all, antidepressants produce dysregulations in other receptors, including postsynaptic and presynaptic adrenoceptors and certain subclasses of the dopamine and serotonin receptors.

These multiple actions of therapeutic drugs suggest that many kinds of biological defects may play a part in manic-depressive illness. Possible defects in neurotransmission include abnormalities in receptors and related molecules; various components of the second-messenger pathways; various proteins that modulate ion transport and indirectly increase or decrease activity in second-messenger systems; and various G proteins, which couple to receptors and stimulate or inhibit intracellular second messengers [see "G Proteins," by Maurice E. Linder and Alfred G. Gilman; *SCIENTIFIC AMERICAN*, July]. So far, however, no one has found direct evidence of any such molecular abnormality in patients.

Episodes of illness gradually become more frequent and more severe in many manic-depressive patients. This pattern of deterioration suggested to Robert M. Post of the National Institute of Mental Health an analogy with an experimental process known as kindling. Scientists kindle convulsions in rodents by stimulating the animals' brains with electricity to induce seizures. Each repetition of the experiment lowers the electrical threshold for the next seizure, leading finally to spontaneous seizures.


Post proposed that manic-depressive illness progresses in a similar fashion, each episode facilitating the next one. This mechanism would account for both the progression of the illness and the

deleterious effects of interrupting treatment with lithium or anticonvulsant medication. After such an interruption, patients may fail to respond to a resumption of the medication, even if it had been effective earlier.

One can also infer aspects of the biology of schizophrenia from the biochemical action of the therapeutic neuroleptic drugs, which include chlorpromazine. Arvid Carlsson of the University of Göteborg sought to explain why these drugs cause animals to produce increased quantities of breakdown products of dopamine. He suggested the effect was a compensatory response of the presynaptic neuron to a postsynaptic blockade [see illustration on opposite page].

As the different molecular and pharmacologic forms of dopamine receptors became known, the D_2 dopamine receptor emerged as the principal site of action of antipsychotic medications. Some of the drugs seem to work by means of interactions with other neurotransmitter systems: among these interactions are the balance between the neural pathways containing the D_1 and the D_2 dopamine receptors and the balance between pathways containing certain serotonin receptors ($5HT_2$) and the D_2 dopamine receptors.

Maria and Arvid Carlsson recently proposed that schizophrenia is characterized by the disruption of a balance between dopamine neurons originating in the midbrain and glutamate neurons originating in the cerebral cortex. The imbalance could be an excess of dopamine or a deficit of glutamate, or both. A reduction in glutamate neurons would be consistent with the apparent cortical atrophy seen in schizophrenia. This theory fits with the effects of known psychosis-producing drugs of abuse: PCP,



ALARMING GENERATIONAL TREND suggests that some environmental factor is increasing the incidence of mood disorders in people who are genetically susceptible to it. These graphs compare two groups of relatives of bipolar patients—one born before 1940, the other born later. At each age at which a comparison is possible, the later cohort is at far higher risk of developing bipolar illness or related psychosis.

a hallucinogen, blocks glutamate receptors, whereas amphetamine, which can produce psychosis with chronic use, stimulates dopamine release.

Studies of the biochemical action of these drugs and related clinical research have brought us to an era of rationally based design of medications. Once neuropharmacologists understood that chlorpromazine works by blocking dopamine receptors, they were able to synthesize haloperidol, which strongly blocks dopamine receptors but has little effect on other receptors. Similarly, after scientists found that the antidepressant imipramine blocks reuptake of the neurotransmitter serotonin, they were able to design fluoxetine, which specifically blocks serotonin reuptake but has very little effect on the reuptake of other monoamines.

Pharmacology and neurobiology continue to feed off each other. Over the past few years, clinical trials have shown


that clozapine helps about 30 percent of those schizophrenic patients who do not respond to other antipsychotic medications or who develop intolerable side effects. This drug's unusual properties, such as its specific interaction with certain dopamine and serotonin receptors (D_4 and $5HT_2$), may help in the effort to understand schizophrenia itself.

Depression turns out to involve hormonal systems of a much wider scope than had been realized. Cortisol, a hormone secreted by the adrenal glands, constitutes the main circulating steroid associated with stress in humans. Many severely depressed patients show persistently elevated blood cortisol, implying a malfunction in the system that normally governs it.

George P. Chrousos of the National Institute of Child Health and Devel-

opment and Philip W. Gold of the National Institute of Mental Health interpret this failure as the result of a prolonged activation of the brain's stress system. This system—a complex of neuronal, hormonal and immunologic responses—comes into play when some stress provokes the brain, causing its hypothalamic centers to secrete corticotropin-releasing hormone (CRH). This factor in turn stimulates the pituitary gland—just under the brain—to produce the hormone adrenocorticotropin, which circulates to the adrenal glands and stimulates their release of cortisol. This process normally turns itself off when the excess cortisol reaches its receptors (glucocorticoid receptors) in the brain and suppresses CRH production there. In depressed patients, however, Gold found that production of CRH is excessive and that this suppression fails.

The CRH-producing neurons of the



The difference between worldwide and worldwise.



What difference does it make that Lufthansa flies to more countries worldwide than any other airline? Or that every 60 seconds a Lufthansa plane is taking off for yet another one of our 182 destinations around the world? The point is that, over the years, every experience with every traveler from every corner of the world has helped us to understand you. We've grown accustomed to differences in custom. All the better to help you feel at home everywhere. And that makes all the difference in the world.

A passion for perfection.®



Lufthansa

For more information and reservations call 1-800-645-3880 or see your Travel Agent. Lufthansa is a participant in the mileage programs of United, Delta and USAir.

A rose, twelve stories high

Fish that eat fruit.

And the air you breathe.

This is the rain forest.

Prepared as a public service by Ogilvy & Mather Photo by Duncan Sim © 1990 World Wildlife Fund



Every second another acre of rain forest is destroyed forever. World Wildlife Fund needs your help. 1-800-CALL-WWF.

World Wildlife Fund  Rain Forest Rescue Campaign

© 1992 SCIENTIFIC AMERICAN, INC

hypothalamus are principally regulated by neurons containing norepinephrine, which originate in the hindbrain. These CRH and norepinephrine neurons serve the stress system as central stations. Each set of neurons stimulates the other. In addition, each responds similarly to many neurotransmitters and peptide modulators of neurotransmission. Because many antidepressant drugs affect these neurotransmitters, they must also influence the regulation of the stress system.

The stress system of the brain sets the level of arousal and the emotional tone, alters the ease with which various kinds of information can be retrieved and analyzed, and aids in the initiation of specific actions. All these functions are disordered in depressed patients, and as a result they become sad, have trouble concentrating and become incapable of making decisions.

The anatomy of the stress system starts with the locus coeruleus in the hindbrain (the major source of norepinephrine-producing neurons) and the paraventricular nucleus of the hypothalamus (the brain's major CRH-producing region). From there the connections reach into the cerebrum; these connections include dopamine-producing neurons that project into the mesolimbic dopamine tract, which helps to control motivation, reward and reinforcement. A connection of CRH neurons to the amygdala and hippocampus is important for memory retrieval and emotional analysis of information pertinent to the environmental events that induced the stress.

The general concepts of stress system dysregulation apply to many psychiatric and other diseases, and it will require a considerable amount of basic and clinical research to determine whether a causal relation exists between mood disorders and this kind of stress response dysregulation.

Molecular genetics can test hypotheses on the biology of these diseases because the predisposition to them is almost certainly inherited. The task is

BRAIN STRESS SYSTEM extends from the hindbrain and hypothalamus to cerebral destinations inside and outside the limbic system (*blue*). It includes neurons containing norepinephrine (*dark blue*), CRH (*red*) and dopamine (*green*). When activated, the system affects mood, thought and, indirectly, the secretion of cortisol by the adrenal glands. Deactivation normally begins when cortisol binds to hypothalamic receptors. But in depression this shutdown fails, producing chronic activation.

made difficult, however, by the complexity of inheritance in schizophrenia and manic-depression. Neither illness is inherited through one dominant or recessive gene. Analysts must take into account that illness might result from coordinated actions of several genes at several different locations (loci) or, alternatively, from genetic heterogeneity (in which the same illness can be caused by a mutation at any one of several loci).

There are two major strategies for finding pathogenic genes. One can systematically search each chromosome, or one can investigate a candidate gene—such as that for a given receptor—which is known to code for proteins related to the disease.

DNA markers now exist for nearly every segment of every chromosome. In any family, each parent contributes to the child a single segment from either one or the other chromosome in a given pair of chromosomes. The illness is linked to a marker location on the genetic map if, and only if, one ancestral chromosome segment is consistently inherited with illness throughout a pedigree. Whenever there is a linkage to the marker, one can be sure that a gene for the illness resides somewhere on that chromosome segment.

Such mapping has consistently shown that a proportion of the families of Alzheimer's patients have illness linked to markers on the long arm of chromosome 21. No such linkages have been demonstrated conclusively for manic-depressive illness or schizophrenia. One widely publicized study of a large Amish pedigree linked manic-depressive illness to markers on chromosome 11; another study of a series of pedigrees in Iceland and England linked schizophrenia to markers on chromosome 5. But later analyses led the investigators to withdraw their conclusions, and no other researchers have confirmed the findings.

Linkage to manic-depressive illness has been reported more than once at the tip of the long arm of the X chromosome, but the linkage remains controversial. One can expect more definitive results from several large-scale international efforts, now under way, to scan the entire gene map of families touched by either schizophrenia or manic-depressive illness.

Many genes encoding the molecules involved in neurotransmission are candidates for the defects underlying manic-depressive illness or schizophrenia. A study of several such candidates for manic-depressive illness was performed on a series of 20 pedigrees at the National Institute of Mental Health by

Margret R. Hoehe, Sevilla D. Detera-Wadleigh, Wade H. Berrettini, Pablo V. Gejman and one of us (Gershon). The group tested structural genes for many of the receptors we have described, including norepinephrine (three alpha-receptor genes and two beta-receptor genes), dopamine (D_2 and D_4) and corticosteroid receptors, and the gene of a G protein subunit ($G_{s\alpha}$). Other investigators have studied the D_2 dopamine receptor in schizophrenia pedigrees. Linkage was strongly excluded for each of these genes.

When linkage of a candidate gene to illness can be ruled out, one can conclude that no mutation in the candidate gene determines inheritance of the susceptibility to the illness. The only qualification to this general rule is the statistical and technical limits on our power to detect or rule out linkage. Methods other than linkage can also scan for mutations in candidate genes, and many genes remain to be examined.

We expect our understanding of the biology of schizophrenia and mood disorders to expand dramatically, fueled by the impressive advances in neurobiology, cognitive neuroscience and genetics. Precise diagnostic tests for persons at risk for illness, treatments based on knowledge of molecular alterations that lead to illness, understanding of how environmental events interact with the brain to produce illness and, eventually, the development of gene therapy are all goals that may be achieved.

FURTHER READING


INTERACTIONS BETWEEN GLUTAMATERGIC AND MONOAMINERGIC SYSTEMS WITHIN THE BASAL GANGLIA—IMPLICATIONS FOR SCHIZOPHRENIA AND PARKINSON'S DISEASE. M. Carlsson and A. Carlsson in *Trends in Neurosciences*, Vol. 13, No. 7, pages 272-276; July 1990.

GENETIC MAPPING OF COMMON DISEASES: THE CHALLENGES OF MANIC-DEPRESSIVE ILLNESS AND SCHIZOPHRENIA. E. S. Gershon, M. Martinez, L. R. Goldin and P. V. Gejman in *Trends in Genetics*, Vol. 6, No. 9, pages 282-287; September 1990.

MANIC-DEPRESSIVE ILLNESS. Frederick K. Goodwin and Kay Redfield Jamison. Oxford University Press, 1990.

THE BIOCHEMICAL BASIS OF NEUROPHARMACOLOGY. Jack R. Cooper, Floyd E. Bloom and Robert H. Roth. Oxford University Press, 1991.

THE CONCEPTS OF STRESS AND STRESS SYSTEM DISORDERS: OVERVIEW OF PHYSICAL AND BEHAVIORAL HOMEOSTASIS. George P. Chrousos and Philip W. Gold in *Journal of the American Medical Association*, Vol. 267, No. 9, pages 1244-1252; March 4, 1992.



Aging Brain, Aging Mind

Late in life the human brain suffers attrition of certain neurons and undergoes chemical alterations. Yet for many people, these changes do not add up to a noticeable decline in intelligence

by Dennis J. Selkoe

Contemplate senescence, as Shakespeare did. In *As You Like It*, his memorable character Lord Jaques enumerates seven ages of man, concluding with this sad description:

*Last scene of all, that ends this strange
eventful history,
Is second childishness and mere oblivion.*


For many of us, as for the melancholy Jaques, the prospect of aging continues to conjure images of inexorable, devastating decline, a slow march toward mindlessness and mortality. But is severe deterioration of the brain—and thus the mind—inevitable?

The answer is no. Admittedly, research indicates that as youth fades, certain molecules and cells in the brain become increasingly impaired or disappear. Some of the changes can undoubtedly disrupt cognition if they accumulate past critical thresholds. Yet studies of human behavior suggest that a mind-eroding buildup of damage is by no means an automatic accompaniment of longevity.

Older adults who truly lose their minds probably do so because a specific disease markedly accelerates or is superimposed on the aging process. In developed nations, the leading cause of senile dementia—loss of memory and reason in the elderly—is Alzheimer's disease. Other causes include the occurrence of multiple strokes or Parkinson's disease.

Physicians cannot always distinguish between older people who suffer from minor, relatively stable forgetfulness and those who are in the early stages of Alzheimer's disease or another progressive, dementing disorder. Ongoing research into normal aging and into disorders of the mind will both enable doctors to make those distinctions and give rise to palliative and preventive therapies. For most students of the aging brain, the ultimate goal is to enhance the quality of its function in old age, not necessarily to prolong life, although the latter could certainly result from the former.

Scientists who study the structural and chemical changes that typify the aging brain in the absence of disease find that the changes are heterogeneous, like the brain itself. The brain consists not only of diverse neurons (the signal-conveying cells) but also of varied glial cells (which help to support and repair neurons) and of blood vessels. Certain subsets of cells and areas of the brain are more prone to age-related damage than others. And the time of onset and the mix and extent of



physical alterations, as well as the effects on intellect, can differ dramatically from person to person. In general, however, it seems safe to say that most of the structural and chemical modifications I will discuss become apparent in late middle life, that is, in the fifties and sixties. Some of them become pronounced after age 70. Because there probably is no unifying mechanism that underlies all senescence (age-related dysfunction of cells and molecules) in the brain, it seems unlikely that investigators will find a singular elixir

that will retard or reverse every decline.

Age-associated changes have been most studied in neurons, which in general do not multiply after birth. As individuals grow older, their overall number of brain neurons decreases, but the pattern is by no means uniform. For example, very few neurons disappear from areas of the hypothalamus that regulate the secretion of certain hormones by the pituitary gland.


In contrast, many more nerve cells tend to vanish from the substantia nigra and locus coeruleus, which are specialized populations of cells in the brain stem. Parkinson's disease can decimate some 70 percent or more of the neurons in those areas, seriously disrupting motor function. Aging alone usually eliminates many fewer cells, although older individuals who exhibit mild symptoms reminiscent of Parkinson's disease—decreased flexibility, slowness of movement and a stooped, shuffling gait—may have lost up to 30 or 40 percent of the original complement.

Parts of the limbic system, including the hippocampus, undergo variable amounts of cell death as well. (The limbic system is central to learning, memory and emotion.) Researchers have estimated that about 5 percent of neurons in the hippocampus disappear with each decade in the second half of life. This calculation suggests that about 20 percent of neurons are lost in that period. The attrition is patchy, though; some hippocampal areas show no significant decline.

Even when neurons themselves survive, their cell bodies and their complex extensions known as axons and dendrites (or, collectively, as neurites) may atrophy. Neurons bear a single axon that relays signals to other neurons, often a dis-

GEORGE BERNARD SHAW, who died in 1950 when he was 94, wrote several plays in his nineties. A major goal of research into the aging brain is to increase the number of people who retain mental vigor throughout life.

DENNIS J. SELKOE, who in 1988 received the Leadership and Excellence in Alzheimer's Disease Award from the National Institute on Aging, is co-director of the Center for Neurologic Diseases at Brigham and Women's Hospital in Boston. He is also professor of neurology and neuroscience at Harvard Medical School. Selkoe received his medical degree from the University of Virginia. This is his second article for *Scientific American*.



BRAIN STRUCTURES indicated in boldface are involved in learning, memory and reasoning. They normally undergo a number of anatomic changes late in life that can potentially impair cognition. The alterations listed are just a sampling, and they generally occur only in selected parts of the affected structures. The weight of the brain declines with age as well (*graph*). Robert D. Terry and his colleagues at the University of California at San Diego, who collected the data plotted on the graph, conclude that the decrease stems in part from shrinkage of large neurons.

tance away. Dendrites, which are more abundant and are found in large branching arbors, generally receive signals from other neurons.

Cell-body and neuritic atrophy commonly occur with aging in a number of brain areas important to learning, memory, planning and other complex intellectual functions. Large neurons, in particular, shrink in parts of the hippocampus and the cerebral cortex. And cell bodies and axons can degenerate in certain acetylcholine-secreting neurons that project from the basal forebrain to the hippocampus and diverse areas of the cortex. Acetylcholine is one of various neurotransmitters by which neurons convey signals to one another.

Yet not all neuronal changes are necessarily destructive. Some may represent attempts by surviving neurons to compensate for loss or shrinkage of other neurons and their projections. Indeed, Paul D. Coleman, Dorothy G. Flood and Stephen J. Buell of the University of Rochester Medical Center have observed a net growth of

dendrites in some regions of the hippocampus and cortex between middle age (the forties and fifties) and early old age (the early seventies), followed by a regression of dendrites in late old age (the eighties and nineties). These investigators postulate that the initial dendritic growth reflects an effort by viable neurons to cope with the age-associated loss of their neighbors. Apparently, this ability to compensate fails in very old neurons. Studies of adult rats show a similar capacity for growth; longer and more complex dendrites appear in the visual cortex after the animals are exposed to visually stimulating environments.

Such findings are encouraging. They suggest both that the brain is capable of dynamic remodeling of its neuronal connections, even in the later years, and that therapy of some kind might augment this plasticity. On the other hand, the functionality of the dendrites that appear in old age has yet to be determined.

In addition to changes in their number and in the structure of their cell

bodies and neurites, neurons can undergo alteration of their internal architecture. For example, the cytoplasm of certain cells of the hippocampus and other brain areas vital to memory and learning can begin to fill with bundles of helically wound protein filaments known as neurofibrillary tangles. An abundance of such tangles in these and other brain areas is believed to contribute to the dementia of Alzheimer's disease, but the significance of small amounts in the undiseased brain is less clear. The development of tangles during aging seems to indicate that certain proteins, particularly those of the cytoskeleton, or the internal scaffolding of the cell, are being chemically modified in ways that could contribute to inefficient signaling by these neurons.

In another internal alteration, neuronal cytoplasm in many parts of the brain becomes increasingly dotted with innumerable granules containing lipofuscin, a fluorescent pigment. This pigment is thought to derive from lipid-rich internal membranes that have been incompletely digested. Again, investiga-

tors disagree over whether lipofuscin granules harm cells or are mere markers of longevity.

As is true of neurons, glial cells, which have a supporting role in brain function, also become altered. Robert D. Terry of the University of California at San Diego and other investigators have established that the type known as fibrous astrocytes increases steadily in size and number after age 60. Proliferation of these cells, which are capable of releasing diverse factors that promote neuronal and neuritic growth, is of unknown consequence. Perhaps it represents another attempt by the brain to compensate for gradual decrements in the number and structure of neurons.

Meanwhile the areas between neurons are also undergoing change. In humans, monkeys, dogs and certain other animals, the extracellular spaces of the hippocampus, cerebral cortex and other brain regions commonly accumulate moderate numbers of spherical deposits called senile plaques. These plaques, which develop very slowly, are primarily aggregates of a small molecule known as the beta-amyloid protein. Amyloid protein also accumulates in scattered blood vessels in these regions and in the meninges, the connective tissue that envelops the brain.

Researchers have not fully determined which cells give rise to these protein deposits and what effect the accumulations have on nearby neurons in healthy elders. The answers should emerge soon, however, because the dramatically increased deposition of amyloid protein in patients with Alzheimer's disease has pushed such issues to the forefront of research.

The diverse structural alterations that occur in the aging brain result from deleterious changes in the activity or abundance of molecules that are important to the integrity and functioning of cells. One of the most venerable theories of aging holds that cells throughout the body senesce because defects slowly accrue in their DNA, the material from which genes are constructed. Genes carry the chemical instructions that inform cells precisely how to synthesize proteins. At some point, the scenario goes, damage to DNA lowers the quality or quantity of critical proteins (such as certain enzymes) in cells. Or the damage may increase the activity or amount of undesirable proteins (such as those that promote the development of cancers).

Until recently, genetic research focused almost entirely on the chromosomal DNA in the nucleus, the long strands of helical DNA that collectively

store the genes for virtually all the proteins made in cells. Such work indicated that the enzymatic machinery designed to excise and repair faulty nuclear DNA becomes less efficient late in life and perhaps in the presence of certain brain diseases. Scientists have also found evidence that cellular controls governing genetic activity may be relaxed during aging. One mechanism may involve subtle elimination of methyl groups (CH_3) from certain parts of DNA molecules [see "A Different Kind of Inheritance," by Robin Holliday; *SCIENTIFIC AMERICAN*, June 1989].



In recent years, investigators have begun to suspect that DNA in a special cellular location—the mitochondria—may also contribute to senescence of the brain. Mitochondria are intracellular "power plants" that provide cells with critically needed energy. They contain their own snippet of DNA, which bears instructions for the manufacture of 13 proteins needed for energy generation. If mitochondrial DNA slowly became partially defective, the defects could result in production of damaged mitochondrial proteins or in the elimination of such proteins.

The evidence favoring a causal role for mitochondrial DNA in some of the

changes associated with aging includes the finding that such DNA seems to be more susceptible to damage than is nuclear DNA. One reason may be that the DNA-repair machinery in the organelles is less effective than that in the nucleus. Mitochondrial DNA is also probably subject to more attack by highly reactive oxygenated compounds known as free radicals. Such compounds are a continual by-product of the reactions through which mitochondria produce energy. Free radicals are made during other cellular processes as well and in response to ionizing radiation. They oxidize, or add oxygen atoms to, molecules, thereby altering them.

Furthermore, investigators have discovered that a crucial enzyme encoded by mitochondrial DNA—cytochrome oxidase—declines with age in rat brains. Several researchers have also identified specific deletions in stretches of mitochondrial DNA in aged brains and in patients with certain age-linked brain disorders, such as Parkinson's disease.




Even if most nuclear and mitochondrial genes remained unaltered and gave rise to the proper amounts of normal proteins, subsequent modifications of the proteins



HIPPOCAMPAL NEURONS (photographs) have been stained in brain specimens from healthy humans in their fifties, seventies and nineties and from a patient with Alzheimer's disease (left to right). Dorothy G. Flood and Paul D. Coleman of the University of Rochester Medical Center find that the average length of dendritic trees (branched shapes) on such neurons increases between middle age and old age in healthy adults, regressing only in late old age (graph). The normal growth may reflect an attempt by the brain to compensate for destructive age-linked changes. Dendrites in Alzheimer's patients do not exhibit age-related growth.

Dementia in the U.S.

In 1992 the Framingham Study, which repeatedly assesses the health of a large group of subjects as they age, estimated the prevalence of dementia (a), including Alzheimer's disease (b). The Alzheimer's figures differ from those of a survey conducted in East Boston (by a team led by Denis A. Evans of Harvard Medical School), probably because the Framingham group applied a narrower definition of the disease. The Framingham Study, consistent with others, also found Alzheimer's disease to be the leading cause of persistent dementia in late life (c). Some currently treatable causes are listed (d).



*Numbers do not add up to 100 percent because of rounding off.

d SOME TREATABLE CAUSES OF DEMENTIA

Medications	Certain tumors or infections of the brain
Emotional depression	Blood clots pressing on the brain
Vitamin B ₁₂ deficiency	Metabolic imbalances (including thyroid, kidney or liver disorders)
Chronic alcoholism	

could lead to molecular mischief in late life. Proteins can undergo a number of distinct chemical modifications, including oxidation of certain of their component amino acids, glycosylation (addition of carbohydrate side chains) or cross-linking (formation of strong chemical bridges between proteins). Such modifications occur normally and enable proteins to carry out their functions. On the other hand, there is abundant evidence that as people age, many proteins accumulate unhelpful modifications. For example, the levels of oxidized proteins in skin cells of humans and brain cells of rats increase progressively with age. In very old rats, such proteins can account for 30 to 50 percent of the total protein content of the cell. Cells from young adults with progeria, a rare and remarkable syndrome marked by premature aging of many body tissues, contain levels of oxidized proteins that approach those found in healthy 80-year-olds.

Because enzymes are the proteins that catalyze many of the most important chemical reactions in cells, they have received the greatest scrutiny. Several enzymes that synthesize neurotransmitters or their receptors become less active as an individual ages. For some of these enzymes, postsynthetic modification may be partly at fault.

It is a cruel irony that proteases, the very enzymes responsible for degrading oxidized and other proteins, themselves undergo oxidation and loss of activity. The problem of damaged proteases, potentially bad enough on its own, may be exaggerated by a parallel decline in the enzymes superoxide dismutase and catalase. These proteins normally inactivate free radicals and defend against oxidative damage to diverse kinds of molecules. At least in rats, they become increasingly scarce in old age.

John M. Carney of the University of Kentucky and Robert A. Floyd of the Oklahoma Medical Research Foundation and their co-workers have recently provided some of the first evidence that such oxidation may lead to a loss of mental function. Comparing aged and young gerbils, they showed that significantly more proteins in the older animals were oxidized. This increase was accompanied by a decrease in activity of certain critical enzymes. Moreover, the older animals had more trouble than did the younger ones in navigating a radial maze.

When the scientists injected the aged gerbils with a compound (*N*-tert-butyl- α -phenylnitron) known to inactivate oxygen free radicals and thus decrease oxidation, the levels of oxidized proteins declined, and the activity of the en-

zymes increased to the levels characteristic of young animals. What is more, the biochemical improvements were accompanied by the restoration of maze-running skill to youthful levels. When the therapy was stopped, the amount of oxidized protein and enzymatic activity reverted to that typical of old animals.

Many important nonprotein molecules in the brain also change significantly in structure or amount during aging. There is evidence that the long chains of carbon atoms making up the lipids in membranes that envelop cells and internal organelles undergo chemical modifications. Among these is destructive oxidation by free radicals. As a result, the precise composition of the membranes can shift, subtly altering their behavior.

For example, investigators have documented an age-related decrease in the fluidity of the membranes making up synaptosomes, tiny neuronal vesicles involved in the storage and release of neurotransmitters. And age-related changes occur in the lipid composition of the myelin that sheathes and insulates axons. Alteration of myelin can have a measurable effect on the speed and efficiency with which nerve fibers propagate electrical impulses over long distances.


The molecular changes I have discussed are but a small sampling of those that have been found in aged brains of humans and other mammals. In trying to make sense of such alterations, scientists immediately confront the problem of determining whether a phenomenon they have documented is the proverbial "cart" or "horse." For instance, there is little doubt that DNA accrues damage over the years. But does the damage result, for example, in increased oxidation of enzymes, or does oxidation occur first and lead to accumulation of DNA deficits? The likelihood is that both sequences can happen. Once many processes get started, they undoubtedly exacerbate others, triggering a complex cascade of events.

Equally important is the question of what bearing all these diverse anatomic and physiological age-linked changes have on the mind. In many people, the answer may be "very little." Until scientists can record the mental functioning of large numbers of healthy people close to the time they die and then correlate such data with structural and chemical changes in their brains, any links between specific physical alter-

ations and disruptions of the intellect will remain murky.

We do know that in people who are free of Alzheimer's disease and other specific brain-threatening disorders, the extent of anatomic and physiological alterations tends to be modest. In many studies reporting an age-related neurochemical deficit—such as reduction in the activity of a particular enzyme or in the levels of selected proteins or RNA molecules—the measures reported for elderly adults have ranged from 5 to 30 percent below those in young adults. The degree of neuronal loss in various regions of the brain falls into roughly the same range.

Although a 30 percent loss might seem quite high, such gradual declines



TISSUE FROM THE BRAIN of a 69-year-old man is riddled with the classic lesions of Alzheimer's disease—senile plaques and neurofibrillary tangles. The plaque visible in this fragment (*large dark-gold sphere*) consists of beta-amyloid protein and, at the periphery, damaged axons and dendrites (*dark squiggles*). The tangles, which are twisted fibers that fill the cytoplasm, make a number of cells appear blackened (*small dark blobs*). Plaques and tangles appear in the brains of healthy aged adults, but usually to a much lesser extent and in restricted regions.

often appear to have little practical effect on thinking. Indeed, positron emission tomographic (PET) imaging indicates that the brains of healthy people in their eighties are almost as active as those of people in their twenties. As is true of other organs, the brain appears to have considerable physiological reserve and to tolerate small losses of neuronal function.

Epidemiological and psychological studies paint a similar picture. Estimates of the prevalence of dementia vary, but the most dire of them—derived from a door-to-door study conducted by Denis A. Evans of Harvard Medical School and his colleagues—indicates that as a group, almost 90 percent of all people older than 65 years are free of dementia. Evans and his co-workers reported in 1991 that fewer than 5 percent of subjects aged 65 to 75 exhibited symp-

toms of dementia, a figure that rose to about 20 percent in the subjects aged 75 to 84. Then the number jumped to about 50 percent in people older than 85 years (which is twice as high as certain other estimates). As disturbing as the data are for people older than 75 years, the figures still indicate that a good many people escape major disturbances in cognition in the later years.

Analyses of performance in healthy old adults lead to a similar conclusion. For instance, Arthur L. Benton, Daniel Tranel and Antonio R. Damasio of the University of Iowa College of Medicine found that when people in their seventies and eighties remain in good health, they show only a subtle decline in performance on tests of memory, perception and language.

One decrement on which many studies agree is a reduction in the speed of some aspects of cognitive processing. Hence, septuagenarians may be unable to retrieve certain details of a particular past event quickly—say, the precise date or place—but they are often able to recall the information minutes or hours later. Given enough time and an environment that keeps anxiety at bay, most healthy elderly people score about as well as young or middle-aged adults on tests of mental performance. The more complex a task is (for example, a multistep mathematical problem), the more likely it is that an otherwise healthy elder will perform less well than does a young adult. A message of guarded optimism emerges from many investigations of normal aging: one may not learn or remember quite as rapidly during healthy late life, but one may learn and remember nearly as well.

Taken together, then, the physical, epidemiological and psychological findings suggest that mild to moderate decreases in memory or speed of intellectual processing may be related to a gradual accumulation of standard anatomic and physiological brain changes that accompany aging. In contrast, dementia apparently arises from more specific and excessive changes in subsets of neurons and in neural circuits. In other words, diseases having distinct causes and mechanisms underlie senile dementia. Of course, one might well wonder why people become more prone to various debilitating brain disorders, including Alzheimer's disease, as they age. In many instances, the answers are uncertain.

Because Alzheimer's disease is by far the most common cause of severe in-

tellectual decline in the elderly, let me briefly review the latest research into why it develops, why it becomes increasingly prevalent late in life and how it might eventually be treated. Fortunately, progress in this area has been truly remarkable of late [see "Amyloid Protein and Alzheimer's Disease," by Dennis J. Selkoe; SCIENTIFIC AMERICAN, November 1991].


Until very recently, the question of what causes Alzheimer's disease had to be answered, "We don't know." But rapid advances from many laboratories analyzing the chemistry and molecular biology of the beta-amyloid deposits has now led to identification of the first specific molecular cause of this complex and devastating disorder. In 1991 studies by Alison M. Goate and John A. Hardy and their colleagues at St. Mary's Hospital Medical School in London, and subsequently other research teams, established that particular genetic mutations are at fault in at least some instances.

These DNA mutations occur within the gene that gives rise to the beta-amyloid precursor protein, or beta-APP. This precursor includes within it the beta-amyloid protein that constitutes both senile plaques and vascular amyloid deposits. The normal functions of beta-APP have not yet been revealed, but those of us who study Alzheimer's disease have found that the precursor is made by most cells of the body. We know, too, that the mutated version somehow leads to accelerated extracellular and vascular deposition of the beta-amyloid segment. Some mutations may lead to more or faster amyloid accumulation than others. Hastened depositions, in turn, could partly explain why some people show symptoms earlier than others.

Research on Down's syndrome has contributed importantly to the new understanding. Individuals with that syndrome are born with three copies of chromosome 21 (where the beta-APP gene is located) rather than the normal two copies. They also invariably develop myriad senile plaques and neurofibrillary tangles in the fourth and fifth decades of life. Neuropathological examination of Down's patients who have died early in life has revealed that a few amorphous deposits of amyloid protein can begin to appear in the teens, decades before full-blown senile plaques and neurofibrillary tangles and clinical signs of dementia develop. That critical finding, together with the discovery of

beta-APP mutations in inherited Alzheimer's disease, now makes it clear that amyloid protein deposition can serve as a seminal event in some if not all cases of Alzheimer's disease.

No one is sure how the initially inert protein leads over a long time to the extensive structural and biochemical changes in axons, dendrites, neuronal cell bodies and glial cells that ravish the minds of Alzheimer's victims. One possibility is that the protein itself remains unreactive, but as it collects over many years, it attracts other types of molecules to the deposits. These other molecules may then damage surrounding neurons and glia. Another hypothesis suggests that after the amyloid protein reaches critical concentrations, it directly damages surrounding neurons and glia or makes them more vulnerable to subtle injurious processes that can occur in the brain.



PET SCANS OF BRAINS from a healthy aged adult (left) and a patient with Alzheimer's disease (right) differ markedly. The excess of dark shading in the second image shows that brain activity is impaired.

In any case, the work of several neuroanatomists—including Damasio, Bradley T. Hyman and Gary W. Van Hoesen and their co-workers at the University of Iowa and John Morrison of the Mount Sinai School of Medicine and his colleagues—has shown that a buildup of amyloid protein, combined with the formation of neurofibrillary tangles and other structural changes in neurons and their extensions, contributes to a progressive disconnection of neuronal circuits serving memory and thinking. Over years the limbic system and the association cortices, which are vital to organizing mental processes, appear to become increasingly out of touch with other neuronal areas. This disconnection helps lead to the impaired memory, judgment, abstraction and language that is all too familiar in Alzheimer's patients. Because most motor and sensory functions are spared until rather late in the disease, the changes give rise to the classical, tragic picture of a

person who can walk, talk and eat but cannot make sense of the world.

In spite of recent progress, many central problems need to be addressed. How do mutations in the beta-APP gene lead to accelerated amyloid deposition compared with the slow rate encountered in normal aged humans? Why is such deposition largely confined to the brain, when virtually all tissues synthesize the amyloid precursor? Which cells actually secrete the devastating amyloid fragments? Why do some neurons in certain brain regions, such as the hippocampus, show striking reaction to the presence of amyloid protein, yet other areas, such as the cerebellum, exhibit little or none? Most important, how can the terrible destruction be blocked?

As these questions are being pursued, so is the problem of how to bolster and protect the aging mind. The fact that no

single compound is likely to block all the potential ravages of great longevity is underscored by results of many clinical trials of vitamins, minerals and various other compounds thought to "enhance" biochemical reactions in the brain or increase blood flow. These substances have yielded little or no cognitive improvement in either demented or functional elderly people.

One reasonable "home remedy" would be to stay physically fit. Robert E. Dustman and his colleagues at the University of Utah and other investigators have demonstrated that older subjects who regularly do aerobic exercise perform better on cognitive tests than do sedentary individuals of the same age with low aerobic fitness.

I would additionally advise against ingesting agents, including alcohol, that interfere with the activity of the nervous system. Similarly, I would urge physicians to be cautious when prescribing for elderly patients medications that act on the brain. Extensive experimental and clinical evidence has shown that people older than about 60 years are particularly sensitive to benzodiazepines (such as the sedative Valium) and many other depressants and stimulants of the central nervous system. Compared with young adults, older people suffer a greater decline in reasoning while such drugs are in their system, are affected longer and react to low doses more strongly. These undesirable effects on cognition are even more pronounced in those who already suffer from a dementing illness.


Researchers continue to engage in

READER SERVICE DIRECTORY

The advertisers listed below are making additional information available, *free*. Just circle the corresponding number on the adjacent card, detach and mail. Offer expires December 15, 1992.

1. **ACADEMIC PRESS DICTIONARY OF SCIENCE AND TECHNOLOGY** The largest original dictionary of *any type* produced in the US since the 1960's, this volume features 124,000 definitions, encompassing every field of science today. (800-321-5068)
2. **AMERITECH** Circle #2 for more information on Ameritech's commitment to improving the quality of life in the Great Lakes Region it serves.
3. **ARC SCIENCE SIMULATIONS** An interactive, visual model of the dynamic sky and solar system through time.
4. **AUDIO-FORUM LEARN A FOREIGN LANGUAGE ON YOUR OWN!** Cassette/book courses in 76 languages from Audio-Forum. Free catalog.
5. **BIO-IMAGING TECHNOLOGIES, INC.** The new measure for drug and device development. For business inquiries call 609-883-2000.
6. **BOOKS ON TAPE, INC.**
7. **BRAINMAKER** Find patterns and hidden relationships in data and learn from the past. Complete neural network development system. California Scientific Software. 800-284-8112, 916-478-9040.
8. **CALCULUS BY AND FOR YOUNG PEOPLE** by Don Cohen. Book, worksheets and two videotapes. Telephone 1-217-356-4555 FAX 1-217-356-4593
9. **COURVOISIER COGNAC** Discover the pleasures of cognac and enhance your knowledge. Send for a FREE booklet outlining the most "Frequently Asked Questions on Cognacs & Courvoisier."
10. **DAVIS INSTRUMENTS** The world's smallest and most affordable weather stations designed for use anywhere. PC option too! Free catalog. (800-618-3669).
11. **EDUCALC MAIL STORE** The EduCALC catalog offers the latest in calculators, Palmtop computers and *Informators* at the best discount prices available. Comprehensive selection of books, peripherals and software.
12. **EMI BRONZE REPRODUCTIONS** Remington, Russell, Chiparus, Moreau, Degas. Cast in the highest quality for discerning bronze collectors. (800-777-8126)
13. **EPIC SYSTEMS NEURALYST FOR EXCEL** Neuralyst integrates neural network technology into Microsoft Excel (PC or Macintosh), providing for "intelligent" spreadsheet data analysis capabilities.
14. **GLOBAL INTUITION NETWORK** Promotes research and applied use of intuition in decision making. Free information.
15. **HOECHST-ROUSSEL PHARMACEUTICALS, INC.** Somerville, New Jersey 1-800-451-4455.
16. **HORSTMANN SOFTWARE DESIGN CORPORATION** ChiWriter scientific/multilingual word processor for the IBM PC and compatible. Enables the editing of scientific and foreign language text in completely WYSIWYG fashion. (800-736-8886)
17. **HUGHES COMMUNICATIONS, INC.** Brochure describes services offered in C-band, Ku-band and high-powered DBS satellite communications for corporate networks, cable, broadcasting and other applications.
18. **ISCAN, INC.** World's leading manufacturer of eye movement monitoring systems for medical research, human factors and advanced man/machine interface applications.
19. **JOHN NUVEEN & CO., INC.** To learn more about tax-free investments, send for a free informative brochure which will be sent promptly with a prospectus.
20. **MAZDA MOTOR OF AMERICA** To find out more about the New Mazda and our whole new line of exciting cars and trucks, call 1-800-639-1000.
21. **MEDIA MAGIC** Offers a free 96 page catalog of unique educational and entertaining Software, Videotapes, and Books, about Computers in Science & Art.
22. **NILES & ASSOCIATES** Professors, students and researchers! Organize your references and create bibliographies instantly with EndNote Plus. Call 1-800-554-3049.
23. **PHARMACEUTICAL MANUFACTURERS ASSOCIATION** To learn more about the critical role of pharmaceutical research, call for a free "Good Medicine" brochure. 1-800-538-2692
24. **PRESCIENCE** For information on *Theorist*[®] and *Expressionist*[®], an easy to use equation typesetter, call 415-543-2252.
25. **RAND JOURNAL OF ECONOMICS** A leader in the field of microeconomics, a focus on industrial organization, contracts, and regulation.
26. **SCHERING-PLOUGH RESEARCH INSTITUTE** A research-based company engaged in the discovery, development, manufacturing and marketing of pharmaceutical and health care products worldwide.
27. **SA MEDICINE** A cost-effective and efficient way for practitioners to keep current in the ever-changing field of medicine.
28. **STANFORD GRAPHICS FOR WINDOWS** Delivers stunning presentation graphics, plus powerful data analysis, specifically designed for the business, statistical and technical user.
29. **TOURISM CANADA** Canada's Official Travellers Guide. Everything you need to know plus telephone numbers. Order your free copy today. 1-800-VISIT92.
30. **WOLFRAM RESEARCH/ MATHEMATICA** Mathematica is a system for doing numerical, symbolic and graphical computation used both as an interactive calculation tool and as a programming language. Call 1-800-441-MATH.

If the card is missing, write on a postcard the name and corresponding number of the advertiser(s) you would like information from and send to: Scientific American, September 1992, P.O. Box 5147, Pittsfield, MA 01203-9827.



MEASURES OF PROTEIN CHEMISTRY in the brains of gerbils suggest that certain age-related changes may be reversible. John M. Carney of the University of Kentucky and Robert A. Floyd of the Oklahoma Medical Research Foundation find that the number of carbonyl groups on brain proteins (a marker of protein oxidation) in old gerbils (blue bar in a) is typically higher than in young gerbils (brown). But the levels decline

in response to a drug that inactivates certain oxidizing chemicals (b). Geriatric gerbils are also less efficient than youngsters at negotiating a radial maze (c), such as the one shown in the photograph. After treatment, the number of errors drops (d). The apparent improvement in short-term memory encourages hope that antioxidants may one day help to somewhat protect the minds of aging humans.



lively debate over whether maintaining or increasing mental activity can protect against cognitive decline late in life. Unfortunately, rigorous data on the subject remain elusive.

The value to brain function of dietary restriction—a well-publicized potential antidote to aging—is similarly unclear. A nutritionally balanced but very low calorie diet has been shown to delay many age-related diseases and to increase the life span in a variety of lower mammals. In some studies, rats fed restricted diets exhibited fewer neurochemical changes in their brains late in life than did their better-fed counterparts, and they were more successful in old age at learning to find their way through a maze.

Similarly, Alan Peters and his colleagues at the Boston University School of Medicine maintained rats on a very low calorie regimen that enabled them to live for as long as four years, perhaps a year longer than usual. The team found that the animals lost neurons and developed various age-related neuronal and glial alterations later in life than did control animals. On the other hand, the eventual occurrence of such alterations suggests that calorie control may delay, but will not prevent, senescence of the brain.

No one really understands the mechanism by which marked calorie restriction leads to prolonged survival in test animals. Nor does anyone know the extent to which it might retard cognitive impairment in humans. It is evident, however, that to have any benefit, the approach would probably have to be practiced throughout much of life. Moreover, sudden and severe nutritional deprivation in the elderly could lead to symptoms of dementia, making calorie restriction a risky undertaking if it is attempted without professional guidance.

A more palatable alternative (both literally and figuratively) to serious dietary denial might one day be prolonged administration of antioxidants, such as vitamin E. This vitamin has been shown to extend longevity and retard some age-associated systemic diseases in rodents, but benefits for the aging human brain have not been proved.

For now, the most rational approach to developing treatments for cognitive failure late in life is to decipher the molecular mechanisms underlying dementing diseases and then to design drugs that block one or more critical steps. In Alzheimer's disease, for example, it appears likely that therapies will ultimately be directed at inhibiting the enzymes that liberate the beta-amyloid protein from its precursor, blocking the delivery of the amyloid protein into cerebral tissue or preventing the inflammatory and neurotoxic responses that the protein apparently initiates. Such treatments might also prove helpful for combating mild to moderate forgetfulness in some older people who do not have full-blown dementia. This seems likely because the amyloid plaques and neurofibrillary tangles that characterize Alzheimer's disease do form in areas important for memory and learning during healthy aging, albeit to a much smaller degree. Several therapies are also under study for preventing Parkinson's disease and for preventing or treating stroke [see "Stroke Therapy," by Justin A. Zivin and Dennis W. Choi; *SCIENTIFIC AMERICAN*, July 1991].

During the next three decades, rigorous molecular and clinical examination of brain aging will become more common as the developed nations confront a huge surge in the numbers of very

old people. An estimated three million Americans are 85 years or older today, and that number may double by the year 2020. Discovery of ways to block age-related disorders of higher cortical function without necessarily prolonging life will undoubtedly enable many of the elderly to remain independent and enjoy life well beyond the eighth decade. Successful aging of the body and mind will bring about profound economic and sociological consequences that will require great creativity and vigor to address. Fortunately, at that point, society will possess a valuable resource to help solve those problems: the sharp minds and accessible wisdom of many of its oldest citizens.

FURTHER READING

- NEURON NUMBERS AND DENDRITIC EXTENT IN NORMAL AGING AND ALZHEIMER'S DISEASE. P. D. Coleman and D. G. Flood in *Neurobiology of Aging*, Vol. 8, No. 6, pages 521-545; November/December 1987.
- THE RETARDATION OF AGING AND DISEASE BY DIETARY RESTRICTION. Richard Weindruch and Roy L. Walford. Springfield, Ill., Charles C. Thomas, 1988.
- RNA AND PROTEIN METABOLISM IN THE AGING BRAIN. Caleb E. Finch and David G. Morgan in *Annual Review of Neuroscience*, Vol. 13. Edited by W. M. Cowan et al. Annual Reviews Inc., 1990.
- LONGEVITY, SENESCENCE, AND THE GENOME. Caleb E. Finch. University of Chicago Press, 1990.
- THE MOLECULAR PATHOLOGY OF ALZHEIMER'S DISEASE. Dennis J. Selkoe in *Neuron*, Vol. 6, No. 4, pages 487-498; April 1991.
- BRAIN AGING AND ALZHEIMER'S DISEASE: "WEAR AND TEAR" VS. "USE IT OR LOSE IT." D. F. Swaab in *Neurobiology of Aging*, Vol. 12, No. 4, pages 317-324; July/August 1991.

SECRETS *from a Lifetime of Practice . . .*



Very up-to-date mini-textbooks in Q & A format

Each book in the Secrets Series is a practical, clinical guide that addresses key questions commonly raised by teachers during rounds, in the OR, in the clinic, and on oral exams. SECRETS are manageable, readable textbooks for students and residents, and succinct reviews and references for practitioners.

MEDICAL SECRETS

Edited by ANTHONY ZOLLO, Jr., M.D.

The "Secrets" approach is applied in this book to review most of what a student should know about general internal medicine. Organized by subspecialty areas of internal medicine, the book is an ideal review for examinations, rounds, and clinical discussions.

• June 1991, 550 pages, illustrated

Q. *How can I obtain these SECRETS?*

A. *Send your order today.*

OB/GYN SECRETS

Edited by HELEN L. FREDERICKSON, M.D.
and LOUISE WILKINS-HAUG, M.D., Ph.D.

This mini-textbook of obstetrics and gynecology in question and answer format encompasses general gynecology, reproductive endocrinology and infertility, gynecologic oncology, general obstetrics, maternal complications, the fetus, the placenta, and labor and delivery.

• May 1991, 308 pages, illustrated

CRITICAL CARE SECRETS

Edited by POLLY E. PARSONS, M.D.
and JEANINE P. WIENER-KRONISH, M.D.

The fundamentals required to care for critically ill patients—whether in the intensive care unit, the operating room, the recovery room, or the emergency room—are presented in questions and answers designed both to provide factual information and to stimulate discussion.

• February 1992, 480 pages, illustrated

SURGICAL SECRETS, 2nd edition

Edited by CHARLES M. ABERNATHY, M.D.
and ALDEN H. HARKEN, M.D.

The vanguard book of the "Secrets" series that achieved best-seller status is now in its second edition, and is even better. The text has been revised and updated and includes a *new* Atlas of Technical Tips that contains graphic, original artwork depicting methods of tying sutures, handling instruments, and other basics.

• July 1991, 330 pages, illustrated

PEDIATRIC SECRETS

Edited by RICHARD A. POLIN, M.D.
and MARK F. DITMAR, M.D.

Here is a collection of answers, tips and "secrets" from the experts on a wide range of diagnostic and therapeutic problems that pediatricians face. This book provides both the right question and its answer to help students, residents, and practitioners hone their skills for practice and board exams.

• 1989, 447 pages, illustrated

- ☐ **CRITICAL CARE SECRETS**, \$32.95 U.S.; \$37.95 elsewhere. 480 pages, illustrated, February 1992.
- ☐ **MEDICAL SECRETS**, \$33.95 U.S.; \$38.95 elsewhere. 558 pages, illustrated, 1991.
- ☐ **OB/GYN SECRETS**, \$27.95 U.S.; \$32.95 elsewhere. 308 pages, illustrated, 1991.
- ☐ **PEDIATRIC SECRETS**, \$30.95 U.S.; \$35.95 elsewhere. 450 pages, illustrated, 1989.
- ☐ **SURGICAL SECRETS**, 2nd Edition, \$28.95 U.S.; \$33.95 elsewhere. 330 pages, illustrated, 1991.

Name _____

Street _____

City/State/Zip _____

I enclose payment: ☐ Check ☐ Visa ☐ MasterCard

Credit Card # _____


Expiration Date _____

Signature _____

HANLEY & BELFUS, INC., Medical Publishers

210 South 13th Street, Philadelphia, PA 19107

215-546-7293



How Neural Networks Learn from Experience

Networks of artificial neurons can learn to represent complicated information. Such neural networks may provide insights into the learning abilities of the human brain

by Geoffrey E. Hinton


The brain is a remarkable computer. It interprets imprecise information from the senses at an incredibly rapid rate. It discerns a whisper in a noisy room, a face in a dimly lit alley and a hidden agenda in a political statement. Most impressive of all, the brain learns—without any explicit instructions—to create the internal representations that make these skills possible.

Much is still unknown about how the brain trains itself to process information, so theories abound. To test these hypotheses, my colleagues and I have attempted to mimic the brain's learning processes by creating networks of artificial neurons. We construct these neural networks by first trying to deduce the essential features of neurons and their interconnections. We then typically program a computer to simulate these features.

Because our knowledge of neurons is incomplete and our computing power is limited, our models are necessarily gross idealizations of real networks of neurons. Naturally, we enthusiastically debate what features are most essential in simulating neurons. By testing these features in artificial neural networks, we have been successful at ruling out all kinds of theories about how the brain processes information. The models are also beginning to reveal how the brain may accomplish its remarkable feats of learning.

In the human brain, a typical neuron collects signals from others through a host of fine structures called dendrites. The neuron sends out spikes of electrical activity through a long, thin strand known as an axon, which splits into thousands of branches. At the end of each branch, a structure called a synapse converts the activity from the axon into electrical effects that inhibit or excite activity in the connected neurons. When a neuron receives excitatory input that is sufficiently large compared with its inhibitory input, it sends a spike of electrical activity down its axon. Learning occurs by changing the effectiveness of the synapses so that the influence of one neuron on another changes.

Artificial neural networks are typically composed of interconnected "units," which serve as model neurons. The function of the synapse is modeled by a modifiable weight, which is associated with each connection. Most artificial networks do not reflect the detailed geometry of the dendrites and ax-



ons, and they express the electrical output of a neuron as a single number that represents the rate of firing—its activity.

Each unit converts the pattern of incoming activities that it receives into a single outgoing activity that it broadcasts to other units. It performs this conversion in two stages. First, it multiplies each incoming activity by the weight on the connection and adds together all these weighted inputs to get a quantity called the total input. Second, a unit uses an input-output function that transforms the total

input into the outgoing activity [see "The Amateur Scientist," page 170].


The behavior of an artificial neural network depends on both the weights and the input-output function that is specified for the units. This function typically falls into one of three categories: linear, threshold or sigmoid. For linear units, the output activity is proportional to the total weighted input. For threshold units, the output is set at one of two levels, depending on whether the total input is greater than or less than some threshold value. For sigmoid units, the output varies continuously but not linearly as the input changes. Sigmoid units bear a greater resemblance to real neurons than do linear or threshold units, but all three must be considered rough approximations.

To make a neural network that performs some specific task, we must choose how the units are connected to one another, and we must set the weights on the connections appropriately. The connections determine whether it is possible for one unit to influence another. The weights specify the strength of the influence.

The commonest type of artificial neural network consists of three groups, or layers, of units: a layer of input units is connected to a layer of "hidden" units, which is connected to a layer of output units. The activity of the input units represents the raw information that is fed into the network. The activity of each hidden unit is determined by the activities of the input units and the weights on the connections between

NETWORK OF NEURONS in the brain provides people with the ability to assimilate information. Will simulations of such networks reveal the underlying mechanisms of learning?

GEOFFREY E. HINTON has worked on representation and learning in artificial neural networks for the past 20 years. In 1978 he received his Ph.D. in artificial intelligence from the University of Edinburgh. He is currently the Noranda Fellow of the Canadian Institute for Advanced Research and professor of computer science and psychology at the University of Toronto.




IDEALIZATION OF A NEURON processes activities, or signals. Each input activity is multiplied by a number called the weight. The “unit” adds together the weighted inputs. It then computes the output activity using an input-output function.

the input and hidden units. Similarly, the behavior of the output units depends on the activity of the hidden units and the weights between the hidden and output units.

This simple type of network is interesting because the hidden units are free to construct their own representations of the input. The weights between the input and hidden units determine when each hidden unit is active, and so by modifying these weights, a hidden unit can choose what it represents.

We can teach a three-layer network to perform a particular task by using the following procedure. First, we present the network with training exam-



COMMON NEURAL NETWORK consists of three layers of units that are fully connected. Activity passes from the input units (green) to the hidden units (gray) and finally to the output units (yellow). The reds and blues of the connections represent different weights.

ples, which consist of a pattern of activities for the input units together with the desired pattern of activities for the output units. We then determine how closely the actual output of the network matches the desired output. Next we change the weight of each connection so that the network produces a better approximation of the desired output.

For example, suppose we want a network to recognize handwritten digits. We might use an array of, say, 256 sensors, each recording the presence or absence of ink in a small area of a single digit. The network would therefore need 256 input units (one for each sensor), 10 output units (one for each kind of digit) and a number of hidden units. For each kind of digit recorded by the sensors, the network should produce high activity in the appropriate output unit and low activity in the other output units.

To train the network, we present an image of a digit and compare the actual activity of the 10 output units with the desired activity. We then calculate the error, which is defined as the square of the difference between the actual and the desired activities. Next we change the weight of each connection so as to reduce the error. We repeat this training process for many different images of each kind of digit until the network classifies every image correctly.

To implement this procedure, we need to change each weight by an amount that is proportional to the rate at which the error changes as the weight is changed. This quantity—called the error derivative for the weight, or simply

the EW—is tricky to compute efficiently. One way to calculate the EW is to perturb a weight slightly and observe how the error changes. But that method is inefficient because it requires a separate perturbation for each of the many weights.

Around 1974 Paul J. Werbos invented a much more efficient procedure for calculating the EW while he was working toward a doctorate at Harvard University. The procedure, now known as the back-propagation algorithm, has become one of the more important tools for training neural networks.

The back-propagation algorithm is easiest to understand if all the units in the network are linear. The algorithm computes each EW by first computing the EA, the rate at which the error changes as the activity level of a unit is changed. For output units, the EA is simply the difference between the actual and the desired output. To compute the EA for a hidden unit in the layer just before the output layer, we first identify all the weights between that hidden unit and the output units to which it is connected. We then multiply those weights by the EAs of those output units and add the products. This sum equals the EA for the chosen hidden unit. After calculating all the EAs in the hidden layer just before the output layer, we can compute in like fashion the EAs for other layers, moving from layer to layer in a direction opposite to the way activities propagate through the network. This is what gives back propagation its name. Once the EA has been computed for a unit, it is straightforward to compute the EW for each incoming connection of the unit. The EW is the product of the EA and the activity through the incoming connection.

For nonlinear units, the back-propagation algorithm includes an extra step. Before back-propagating, the EA must be converted into the EI, the rate at which the error changes as the total input received by a unit is changed. (The details of this calculation are given in the box on page 148.)

The back-propagation algorithm was largely ignored for years after its invention, probably because its usefulness was not fully appreciated. In the early 1980s David E. Rumelhart, then at the University of California at San Diego, and David B. Parker, then at Stanford University, independently rediscovered the algorithm. In 1986 Rumelhart, Ronald J. Williams, also at the University of California at San Diego, and I popularized the algorithm by demonstrating that it could teach the hidden units to produce interesting rep-

representations of complex input patterns.

The back-propagation algorithm has proved surprisingly good at training networks with multiple layers to perform a wide variety of tasks. It is most useful in situations in which the relation between input and output is non-linear and training data are abundant. By applying the algorithm, researchers have produced neural networks that recognize handwritten digits, predict currency exchange rates and maximize the yields of chemical processes. They have even used the algorithm to train networks that identify precancerous cells in Pap smears and that adjust the mirror of a telescope so as to cancel out atmospheric distortions.

Within the field of neuroscience, Richard Andersen of the Massachusetts Institute of Technology and David Zipser of the University of California at San Diego showed that the back-propagation

algorithm is a useful tool for explaining the function of some neurons in the brain's cortex. They trained a neural network to respond to visual stimuli using back propagation. They then found that the responses of the hidden units were remarkably similar to those of real neurons responsible for converting visual information from the retina into a form suitable for deeper visual areas of the brain.



Yet back propagation has had a rather mixed reception as a theory of how biological neurons learn. On the one hand, the back-propagation algorithm has made a valuable contribution at an abstract level. The algorithm is quite good at creating sensible representations in the hidden units. As a result, researchers gained confidence in learning procedures in which weights are gradually adjusted to reduce errors. Previously, many workers had assumed that such

methods would be hopeless because they would inevitably lead to locally optimal but globally terrible solutions. For example, a digit-recognition network might consistently home in on a set of weights that makes the network confuse ones and sevens even though an ideal set of weights exists that would allow the network to discriminate between the digits. This fear supported a widespread belief that a learning procedure was interesting only if it were guaranteed to converge eventually on the globally optimal solution. Back propagation showed that for many tasks global convergence was not necessary to achieve good performance.

On the other hand, back propagation seems biologically implausible. The most obvious difficulty is that information must travel through the same connections in the reverse direction, from one layer to the previous layer. Clearly,

How a Neural Network Represents Handwritten Digits

A neural network—consisting of 256 input units, nine hidden units and 10 output units—has been trained to recognize handwritten digits. The illustration below shows the activities of the units when the network is presented with a handwritten 3. The third output unit is most active. The nine panels at the right represent the 256 incoming weights and the 10 outgoing weights for each of the nine hidden units. The red regions indicate weights that are excitatory, whereas yellow regions represent weights that are inhibitory.



this does not happen in real neurons. But this objection is actually rather superficial. The brain has many pathways from later layers back to earlier ones, and it could use these pathways in many ways to convey the information required for learning.

A more important problem is the speed of the back-propagation algorithm. Here the central issue is how the time required to learn increases as the network gets larger. The time taken to calculate the error derivatives for the weights on a given training example is proportional to the size of the network because the amount of computation is proportional to the number of weights. But bigger networks typically require more training examples, and they must update the weights more times. Hence, the learning time grows much faster than does the size of the network.

The most serious objection to back propagation as a model of real learning is that it requires a teacher to supply the desired output for each training example. In contrast, people learn most things without the help of a teacher. Nobody presents us with a detailed description of the internal representations of the world that we must learn to extract from our sensory input. We learn to understand sentences or visual scenes without any direct instructions.

How can a network learn appropriate internal representations if it starts with no knowledge and no teacher? If a network is presented with a large set of patterns but is given no information about what to do with them, it apparently does not have a well-defined problem to solve. Nevertheless, researchers have developed several general-purpose, unsupervised procedures that can adjust the

weights in the network appropriately.

All these procedures share two characteristics: they appeal, implicitly or explicitly, to some notion of the quality of a representation, and they work by changing the weights to improve the quality of the representation extracted by the hidden units.

In general, a good representation is one that can be described very economically but nonetheless contains enough information to allow a close approximation of the raw input to be reconstructed. For example, consider an image consisting of several ellipses. Suppose a device translates the image into an array of a million tiny squares, each of which is either light or dark. The image could be represented simply by the positions of the dark squares. But other, more efficient representations are

The Back-Propagation Algorithm

To train a neural network to perform some task, we must adjust the weights of each unit in such a way that the error between the desired output and the actual output is reduced. This process requires that the neural network compute the error derivative of the weights (EW). In other words, it must calculate how the error changes as each weight is increased or decreased slightly. The back-propagation algorithm is the most widely used method for determining the EW.

To implement the back-propagation algorithm, we must first describe a neural network in mathematical terms. Assume that unit j is a typical unit in the output layer and unit i is a typical unit in the previous layer. A unit in the output layer determines its activity by following a two-step procedure. First, it computes the total weighted input x_j , using the formula

$$x_j = \sum_i y_i w_{ij},$$

where y_i is the activity level of the i th unit in the previous layer and w_{ij} is the weight of the connection between the i th and j th unit.

Next, the unit calculates the activity y_j using some function of the total weighted input. Typically, we use the sigmoid function:

$$y_j = \frac{1}{1 + e^{-x_j}}.$$

Once the activities of all the output units have been determined, the network computes the error $\mathcal{E}$, which is defined by the expression

$$\mathcal{E} = \frac{1}{2} \sum_j (y_j - d_j)^2,$$

where y_j is the activity level of the j th unit in the top layer and d_j is the desired output of the j th unit.

The back-propagation algorithm consists of four steps:

1. Compute how fast the error changes as the activity of an output unit is changed. This error derivative (EA) is

the difference between the actual and the desired activity.

$$EA_j = \frac{\partial \mathcal{E}}{\partial y_j} = y_j - d_j$$

2. Compute how fast the error changes as the total input received by an output unit is changed. This quantity (EI) is the answer from step 1 multiplied by the rate at which the output of a unit changes as its total input is changed.

$$EI_j = \frac{\partial \mathcal{E}}{\partial x_j} = \frac{\partial \mathcal{E}}{\partial y_j} \frac{dy_j}{dx_j} = EA_j y_j (1 - y_j)$$

3. Compute how fast the error changes as a weight on the connection into an output unit is changed. This quantity (EW) is the answer from step 2 multiplied by the activity level of the unit from which the connection emanates.

$$EW_{ij} = \frac{\partial \mathcal{E}}{\partial w_{ij}} = \frac{\partial \mathcal{E}}{\partial x_j} \frac{\partial x_j}{\partial w_{ij}} = EI_j y_i$$

4. Compute how fast the error changes as the activity of a unit in the previous layer is changed. This crucial step allows back propagation to be applied to multilayer networks. When the activity of a unit in the previous layer changes, it affects the activities of all the output units to which it is connected. So to compute the overall effect on the error, we add together all these separate effects on output units. But each effect is simple to calculate. It is the answer in step 2 multiplied by the weight on the connection to that output unit.

$$EA_i = \frac{\partial \mathcal{E}}{\partial y_i} = \sum_j \frac{\partial \mathcal{E}}{\partial x_j} \frac{\partial x_j}{\partial y_i} = \sum_j EI_j w_{ij}$$

By using steps 2 and 4, we can convert the EAs of one layer of units into EAs for the previous layer. This procedure can be repeated to get the EAs for as many previous layers as desired. Once we know the EA of a unit, we can use steps 2 and 3 to compute the EWs on its incoming connections.

also possible. Ellipses differ in only five ways: orientation, vertical position, horizontal position, length and width. The image can therefore be described using only five parameters per ellipse.

Although describing an ellipse by five parameters requires more bits than describing a single dark square by two coordinates, we get an overall savings because far fewer parameters than coordinates are needed. Furthermore, we do not lose any information by describing the ellipses in terms of their parameters: given the parameters of the ellipse, we could reconstruct the original image if we so desired.


Almost all the unsupervised learning procedures can be viewed as methods of minimizing the sum of two terms, a code cost and a reconstruction cost. The code cost is the number of bits required to describe the activities of the hidden units. The reconstruction cost is the number of bits required to describe the misfit between the raw input and the best approximation to it that could be reconstructed from the activities of the hidden units. The reconstruction cost is proportional to the squared difference between the raw input and its reconstruction.

Two simple methods for discovering economical codes allow fairly accurate reconstruction of the input: principal-components learning and competitive learning. In both approaches, we first decide how economical the code should be and then modify the weights in the network to minimize the reconstruction error.

A principal-components learning strategy is based on the idea that if the activities of pairs of input units are correlated in some way, it is a waste of bits to describe each input activity separately. A more efficient approach is to extract and describe the principal components—that is, the components of variation shared by many input units. If we wish to discover, say, 10 of the principal components, then we need only a single layer of 10 hidden units.

Because such networks represent the input using only a small number of components, the code cost is low. And because the input can be reconstructed quite well from the principal components, the reconstruction cost is small.

One way to train this type of network is to force it to reconstruct an approximation to the input on a set of output units. Then back propagation can be used to minimize the difference between the actual output and the desired output. This process resembles supervised learning, but because the desired output is exactly the same as the input, no teacher is required.




TWO FACES composed of eight ellipses can be represented as many points in two dimensions. Alternatively, because the ellipses differ in only five ways—orientation, vertical position, horizontal position, length and width—the two faces can be represented as eight points in a five-dimensional space.

Many researchers, including Ralph Linsker of the IBM Thomas J. Watson Research Center and Erkki Oja of Lappeenranta University of Technology in Finland, have discovered alternative algorithms for learning principal components. These algorithms are more biologically plausible because they do not require output units or back propagation. Instead they use the correlation between the activity of a hidden unit and the activity of an input unit to determine the change in the weight.

When a neural network uses principal-components learning, a small num-

ber of hidden units cooperate in representing the input pattern. In contrast, in competitive learning, a large number of hidden units compete so that a single hidden unit is used to represent any particular input pattern. The selected hidden unit is the one whose incoming weights are most similar to the input pattern.

Now suppose we had to reconstruct the input pattern solely from our knowledge of which hidden unit was chosen. Our best bet would be to copy the pattern of incoming weights of the chosen hidden unit. To minimize the recon-



COMPETITIVE LEARNING can be envisioned as a process in which each input pattern attracts the weight pattern of the closest hidden unit. Each input pattern represents a set of distinguishing features. The weight patterns of hidden units are adjusted so that they migrate slowly toward the closest set of input patterns. In this way, each hidden unit learns to represent a cluster of similar input patterns.

struction error, we should move the pattern of weights of the winning hidden unit even closer to the input pattern. This is what competitive learning does. If the network is presented with training data that can be grouped into clusters of similar input patterns, each hidden unit learns to represent a different cluster, and its incoming weights converge on the center of the cluster.

Like the principal-components algorithm, competitive learning minimizes the reconstruction cost while keeping the code cost low. We can afford to use many hidden units because even with a million units it takes only 20 bits to say which one won.

In the early 1980s Teuvo Kohonen of Helsinki University introduced an important modification of the competitive learning algorithm. Kohonen showed how to make physically adjacent hidden units learn to represent similar input patterns. Kohonen's algorithm adapts not only the weights of the winning hidden unit but also the weights of the winner's neighbors. The algorithm's ability to map similar input patterns to nearby hidden units suggests that a procedure of this type may be what the brain uses to create the topographic maps found in the visual cortex [see "The Visual Image in Mind and Brain," by Semir Zeki, page 68].

Unsupervised learning algorithms can be classified according to the type of representation they create. In principal-components methods, the hidden units cooperate, and the representation of each input pattern is distributed across all of them. In competitive methods, the hidden units compete, and the representation of the input pattern is localized in the single hidden unit that is selected. Until recently, most work on unsupervised learning focused on one or another of these two techniques, probably because they lead to simple rules for changing the weights. But the most interesting and powerful algorithms probably lie somewhere between the extremes of purely distributed and purely localized representations.

Horace B. Barlow of the University of Cambridge has proposed a model in which each hidden unit is rarely active and the representation of each input pattern is distributed across a small number of selected hidden units. He and his co-workers have shown that this type of code can be learned by forcing hidden units to be uncorrelated while also ensuring that the hidden code allows good reconstruction of the input.

Unfortunately, most current methods of minimizing the code cost tend to eliminate all the redundancy among the activities of the hidden units. As a result, the network is very sensitive to the malfunction of a single hidden unit. This feature is uncharacteristic of the brain, which is generally not affected greatly by the loss of a few neurons.

The brain seems to use what are known as population codes, in which information is represented by a whole population of active neurons. That point was beautifully demonstrated in the experiments of David L. Sparks and his co-workers at the University of Alaba-

rameters of the face being represented by that population code. In abstract terms, each face cell represents a particular point in a multidimensional space of possible faces, and any face can then be represented by activating all the cells that encode very similar faces, so that a bump of activity appears in the multidimensional space of possible faces.


Population coding is attractive because it works even if some of the neurons are damaged. It can do so because the loss of a random subset of neurons has little effect on the population average. The same reasoning applies if some neurons are overlooked when the

system is in a hurry. Neurons communicate by sending discrete spikes called action potentials, and in a very short time interval many of the "active" neurons may not have time to send a spike. Nevertheless, even in such a short interval, a population code in one part of the brain can still give rise to an approximately correct population code in another part of the brain.

At first sight, the redundancy in population codes seems incompatible with the idea of constructing internal representations that minimize the code

cost. Fortunately, we can overcome this difficulty by using a less direct measure of code cost. If the activity that encodes a particular entity is a smooth bump in which activity falls off in a standard way as we move away from the center, we can describe the bump of activity completely merely by specifying its center. So a fairer measure of code cost is the cost of describing the center of the bump of activity plus the cost of describing how the actual activities of the units depart from the desired smooth bump of activity.

Using this measure of the code cost, we find that population codes are a convenient way of extracting a hierarchy of progressively more efficient encodings of the sensory input. This point is best illustrated by a simple example. Consider a neural network that is presented with an image of a face. Suppose the network already contains one set of units dedicated to representing noses, another set for mouths and another set for eyes. When it is shown a particular face, there will be one bump of activity in the nose units, one in the mouth units and two in the eye units. The location of each of these activity bumps represents the spatial parameters of the feature encoded by the bump. Describing the four activity bumps is cheaper than describing the raw image, but it



POPULATION CODING represents a multiparameter object as a bump of activity spread over many hidden units. Each disk represents an inactive hidden unit. Each cylinder indicates an active unit, and its height depicts the level of activity.

ma. While investigating how the brain of a monkey instructs its eyes where to move, they found that the required movement is encoded by the activities of a whole population of cells, each of which represents a somewhat different movement. The eye movement that is actually made corresponds to the average of all the movements encoded by the active cells. If some brain cells are anesthetized, the eye moves to the point associated with the average of the remaining active cells. Population codes may be used to encode not only eye movements but also faces, as shown by Malcolm P. Young and Shigeru Yamane at the RIKEN Institute in Japan in recent experiments on the inferior temporal cortex of monkeys.

For both eye movements and faces, the brain must represent entities that vary along many different dimensions. In the case of an eye movement, there are just two dimensions, but for something like a face, there are dimensions such as happiness, hairiness or familiarity, as well as spatial parameters such as position, size and orientation. If we associate with each face-sensitive cell the parameters of the face that make it most active, we can average these parameters over a population of active cells to discover the pa-


would obviously be cheaper still to describe a single bump of activity in a set of face units, assuming of course that the nose, mouth and eyes are in the correct spatial relations to form a face.

This raises an interesting issue: How can the network check that the parts are correctly related to one another to make a face? Some time ago Dana H. Ballard of the University of Rochester introduced a clever technique for solving this type of problem that works nicely with population codes.

If we know the position, size and orientation of a nose, we can predict the position, size and orientation of the face to which it belongs because the spatial relation between noses and faces is roughly fixed. We therefore set the weights in the neural network so that a bump of activity in the nose units tries to cause an appropriately related bump of activity in the face units. But we also set the thresholds of the face units so that the nose units alone are insufficient to activate the face units. If, however, the bump of activity in the mouth units also tries to cause a bump in the same place in the face units, then the thresholds can be overcome. In effect, we have checked that the nose and mouth are correctly related to each other by checking that they both predict the same spatial parameters for the whole face.

This method of checking spatial relations is intriguing because it makes use of the kind of redundancy between different parts of an image that unsupervised learning should be good at finding. It therefore seems natural to try to use unsupervised learning to discover hierarchical population codes for extracting complex shapes. In 1986 Eric Saund of M.I.T. demonstrated one method of learning simple population codes for shapes. It seems likely that with a clear definition of the code cost, an unsupervised network will be able to discover more complex hierarchies by trying to minimize the cost of coding the image. Richard Zemel and I at the University of Toronto are now investigating this possibility.

By using unsupervised learning to extract a hierarchy of successively more economical representations, it should be possible to improve greatly the speed of learning in large multilayer networks. Each layer of the network adapts its incoming weights to make its representation better than the representation in the previous layer, so weights in one layer can be learned without reference to weights in subsequent layers. This strategy eliminates many of the interactions between weights that make back-propagation learning very slow in deep multilayer networks.



BUMPS OF ACTIVITY in sets of hidden units represent the image of a nose and a mouth. These population codes will cause a bump in the face units if the nose and mouth have the correct spatial relation (left). If not, the active nose units will try to create a bump in the face units at one location while the active mouth units will do the same at a different location. As a result, the input activity to the face units does not exceed a threshold value, and no bump is formed in the face units (right).

All the learning procedures discussed thus far are implemented in neural networks in which activity flows only in the forward direction from input to output even though error derivatives may flow in the backward direction. Another important possibility to consider is networks in which activity flows around closed loops. Such recurrent networks may settle down to stable states, or they may exhibit complex temporal dynamics that can be used to produce sequential behavior. If they settle to stable states, error derivatives can be computed using methods much simpler than back propagation.

Although investigators have devised some powerful learning algorithms that are of great practical value, we still do not know which representations and learning procedures are actually used by the brain. But sooner or later computational studies of learning in artificial neural networks will converge on

the methods discovered by evolution. When that happens, a lot of diverse empirical data about the brain will suddenly make sense, and many new applications of artificial neural networks will become feasible.

FURTHER READING

LEARNING REPRESENTATIONS BY BACK-PROPAGATING ERRORS. David E. Rumelhart, Geoffrey E. Hinton and Ronald J. Williams in *Nature*, Vol. 323, No. 6188, pages 533-536; October 9, 1986.
CONNECTIONIST LEARNING PROCEDURES. Geoffrey E. Hinton in *Artificial Intelligence*, Vol. 40, Nos. 1-3, pages 185-234; September 1989.
INTRODUCTION TO THE THEORY OF NEURAL COMPUTATION. J. Hertz, A. Krogh and R. G. Palmer. Addison-Wesley, 1990.
THE COMPUTATIONAL BRAIN. Patricia S. Churchland and Terrence J. Sejnowski. The MIT Press/Bradford Books, 1992.



The Problem of Consciousness

It can now be approached by scientific investigation of the visual system. The solution will require a close collaboration among psychologists, neuroscientists and theorists


by Francis Crick and Christof Koch

The overwhelming question in neurobiology today is the relation between the mind and the brain. Everyone agrees that what we know as mind is closely related to certain aspects of the behavior of the brain, not to the heart, as Aristotle thought. Its most mysterious aspect is consciousness or awareness, which can take many forms, from the experience of pain to self-consciousness.

In the past the mind (or soul) was often regarded, as it was by Descartes, as something immaterial, separate from the brain but interacting with it in some way. A few neuroscientists, such as Sir John Eccles, still assert that the soul is distinct from the body. But most neuroscientists now believe that all aspects of mind, including its most puzzling attribute—consciousness or awareness—are likely to be explainable in a more materialistic way as the behavior of large sets of interacting neurons. As William James, the father of American psychology, said a century ago, consciousness is not a thing but a process.

Exactly what the process is, however, has yet to be discovered. For many years after James penned *The Principles of Psychology*, consciousness was a taboo concept in American psychology because of the dominance of the behaviorist movement. With the advent of cognitive science in the mid-1950s, it became possible once more for psychologists to consider mental processes as opposed to merely observing behavior. In spite of these changes, until recently most cognitive scientists ignored consciousness, as did almost all neuroscientists. The problem was felt to be either purely “philosophical” or too elusive to study experimentally. It would not have been easy for a neuroscientist to get a grant just to study consciousness.

In our opinion, such timidity is ridiculous, so a few years ago we began to think about how best to attack the problem scientifically. How to explain mental events as being caused by the firing of large sets of neurons? Although there are those who believe such an approach is hopeless, we feel it is not productive to worry too much over aspects of the problem that cannot be solved scientifically or, more precisely, cannot be solved solely by using existing scientific ideas. Radically new concepts may indeed be needed—recall the



VISUAL AWARENESS primarily involves seeing what is directly in front of you, but it can be influenced by a three-dimensional representation of the object in view retained by the brain. If you see the back of a person's head, the brain infers that there is a face on the front of it. We know this is true because we would be very startled if a mirror revealed that the front was exactly like the back, as in this painting, *Reproduction Prohibited* (1937), by René Magritte.

modifications of scientific thinking forced on us by quantum mechanics. The only sensible approach is to press the experimental attack until we are confronted with dilemmas that call for new ways of thinking.

There are many possible approaches to the problem of consciousness. Some psychologists feel that any satisfactory theory should try to explain as many aspects of consciousness as possible, including emotion, imagination, dreams, mystical experiences and so on. Although such an all-embracing theory will be

necessary in the long run, we thought it wiser to begin with the particular aspect of consciousness that is likely to yield most easily. What this aspect may be is a matter of personal judgment. We selected the mammalian visual system because humans are very visual animals and because so much experimental and theoretical work has already been done on it [see “The Visual Image in Mind and Brain,” by Semir Zeki, page 68].

It is not easy to grasp exactly what we need to explain, and it will take many careful experiments before visual consciousness can be described scientifically. We did not attempt to define consciousness itself because of the dangers of premature definition. (If this seems like a cop-out, try defining the word “gene”—you will not find it easy.) Yet the experimental evidence that already exists provides enough of a glimpse of the nature of visual consciousness to guide research. In this article, we will attempt to show how this evidence opens the way to attack this profound and intriguing problem.

Visual theorists agree that the problem of visual consciousness is ill posed. The mathematical term “ill posed” means that additional constraints are needed to solve the problem. Although the main function of the visual system is to perceive objects and events in the world around us, the in-

FRANCIS CRICK and CHRISTOF KOCH share an interest in the experimental study of consciousness. Crick is the co-discoverer, with James Watson, of the double helical structure of DNA. While at the Medical Research Council Laboratory of Molecular Biology in Cambridge, he worked on the genetic code and on developmental biology. Since 1976, he has been at the Salk Institute for Biological Studies in San Diego. His main interest lies in understanding the visual system of mammals. Koch was awarded his Ph.D. in biophysics by the University of Tübingen. After spending four years at the Massachusetts Institute of Technology, he joined the California Institute of Technology, where he is now associate professor of computation and neural systems. He is studying how single brain cells process information and the neural basis of motion perception, visual attention and awareness. He also designs analog VLSI vision chips for intelligent systems.

formation available to our eyes is not sufficient by itself to provide the brain with its unique interpretation of the visual world. The brain must use past experience (either its own or that of our distant ancestors, which is embedded in our genes) to help interpret the information coming into our eyes. An example would be the derivation of the three-dimensional representation of the world from the two-dimensional signals falling onto the retinas of our two eyes or even onto one of them.

Visual theorists also would agree that seeing is a constructive process, one in which the brain has to carry out complex activities (sometimes called computations) in order to decide which interpretation to adopt of the ambiguous visual input. "Computation" implies that the brain acts to form a symbolic representation of the visual world, with a mapping (in the mathematical sense) of certain aspects of that world onto elements in the brain.

Ray Jackendoff of Brandeis University postulates, as do most cognitive scientists, that the computations carried out by the brain are largely unconscious and that what we become aware of is the result of these computations. But while the customary view is that this awareness occurs at the highest levels of the computational system, Jackendoff has proposed an intermediate-level theory of consciousness.

What we see, Jackendoff suggests, relates to a representation of surfaces that are directly visible to us, together with their outline, orientation, color, texture and movement. (This idea has similarities to what the late David C. Marr of the Massachusetts Institute of Technology called a "2½-dimensional sketch." It is more than a two-dimensional sketch because it conveys the orientation of the visible surfaces. It is less than three-dimensional because depth information is not explicitly represented.) In the next stage this sketch is processed by the brain to produce a three-dimensional representation. Jackendoff argues that we are not visually aware of this three-dimensional representation.

An example may make this process clearer. If you look at a person whose back is turned to you, you can see the back of the head but not the face. Nevertheless, your brain infers that the person has a face. We can deduce as much because if that person turned around and had no face, you would be very surprised.

The viewer-centered representation that corresponds to the visible back of the head is what you are vividly aware of. What your brain infers about the front would come from some kind of

three-dimensional representation. This does not mean that information flows only from the surface representation to the three-dimensional one; it almost certainly flows in both directions. When you imagine the front of the face, what you are aware of is a surface representation generated by information from the three-dimensional model.

It is important to distinguish between an explicit and an implicit representation. An explicit representation is something that is symbolized without further processing. An implicit representation contains the same information but requires further processing to make it explicit. The pattern of colored dots on a television screen, for example, contains an implicit representation of objects (say, a person's face), but only the dots and their locations are explicit. When you see a face on the screen, there must be neurons in your brain whose firing, in some sense, symbolizes that face.

We call this pattern of firing neurons an active representation. A latent representation of a face must also be stored in the brain, probably as a special pattern of synaptic connections between neurons [see "How Neural Networks Learn from Experience," by Geoffrey E. Hinton, page 144]. For example, you probably have a representation of the Statue of Liberty in your brain, a representation that usually is inactive. If you do think about the Statue, the representation becomes active, with the relevant neurons firing away.

An object, incidentally, may be represented in more than one way—as a visual image, as a set of words and their related sounds, or even as a touch or a smell. These different representations are likely to interact with one another. The representation is likely to be distributed over many neurons, both locally, as discussed in Geoffrey E. Hinton's article, and more globally. Such a representation may not be as simple and straightforward as uncritical introspection might indicate. There is suggestive evidence, partly from studying how neurons fire in various parts of a monkey's brain and partly from examining the effects of certain types of brain damage in humans, that different aspects of a face—and of the implications of a face—may be represented in different parts of the brain.

First, there is the representation of a face as a face: two eyes, a nose, a mouth and so on. The neurons involved are usually not too fussy about the exact size or position of this face in the visual field, nor are they very sensitive to small changes in its orientation. In monkeys,

there are neurons that respond best when the face is turning in a particular direction, while others seem to be more concerned with the direction in which the eyes are gazing.

Then there are representations of the parts of a face, as separate from those for the face as a whole. Further, the implications of seeing a face, such as that person's sex, the facial expression, the familiarity or unfamiliarity of the face, and in particular whose face it is, may each be correlated with neurons firing in other places.

What we are aware of at any moment, in one sense or another, is not a simple matter. We have suggested that there may be a very transient form of fleeting awareness that represents only rather simple features and does not require an attentional mechanism. From this brief awareness the brain constructs a viewer-centered representation—what we see vividly and clearly—that does require attention. This in turn probably leads to three-dimensional object representations and thence to more cognitive ones.

Representations corresponding to vivid consciousness are likely to have special properties. William James thought that consciousness involved both attention and short-term memory. Most psychologists today would agree with this view. Jackendoff writes that consciousness is "enriched" by attention, implying that while attention may not be essential for certain limited types of consciousness, it is necessary for full consciousness.

Yet it is not clear exactly which forms of memory are involved. Is long-term memory needed? Some forms of acquired knowledge are so embedded in the machinery of neural processing that they are almost certainly used in becoming aware of something. On the other hand, there is evidence from studies of brain-damaged patients (such as H.M., described in "The Biological Basis of Learning and Individuality," by Eric R. Kandel and Robert D. Hawkins, page 78) that the ability to lay down new long-term episodic memories is not essential for consciousness.

It is difficult to imagine that anyone could be conscious if he or she had no memory whatsoever of what had just happened, even an extremely short one. Visual psychologists talk of iconic memory, which lasts for a fraction of a second, and working memory (such as that used to remember a new telephone number) that lasts for only a few seconds unless it is rehearsed. It is not clear whether both of these are essential for consciousness. In any case, the division of short-term memory into these two categories may be too crude.

If these complex processes of visual awareness are localized in parts of the brain, which processes are likely to be where? Many regions of the brain may be involved, but it is almost certain that the cerebral neocortex plays a dominant role. Visual information from the retina reaches the neocortex mainly by way of a part of the thalamus (the lateral geniculate nucleus); another significant visual pathway from the retina is to the superior colliculus, at the top of the brain stem.

The cortex in humans consists of two intricately folded sheets of nerve tissue, one on each side of the head. These sheets are connected by a large tract of about half a billion axons called the corpus callosum. It is well known that if the corpus callosum is cut, as is done for certain cases of intractable epilepsy, one side of the brain is not aware of what the other side is seeing.

In particular, the left side of the brain (in a right-handed person) appears not to be aware of visual information received exclusively by the right side. This shows that none of the information required for visual awareness can reach the other side of the brain by traveling down to the brain stem and, from there, back up. In a normal person, such information can get to the other side only by using the axons in the corpus callosum.

A different part of the brain—the hippocampal system—is involved in one-shot, or episodic, memories that, over weeks and months, it passes on to the neocortex, as described in the article by Eric R. Kandel and Robert D.

Hawkins. This system is so placed that it receives inputs from, and projects to, many parts of the brain.

Thus, one might suspect that the hippocampal system is the essential seat of consciousness. This is not the case: evidence from studies of patients with damaged brains shows that this system is not essential for visual awareness, although naturally a patient lacking one, such as H.M., is severely handicapped in everyday life because he cannot remember anything that took place more than a minute or so in the past.


In broad terms, the neocortex of alert animals probably acts in two ways. By building on crude and somewhat redundant wiring, produced by our genes and

by embryonic processes [see "The Developing Brain," by Carla J. Shatz, page 60], the neocortex draws on visual and other experience to slowly "rewire" itself to create categories (or "features") it can respond to. A new category is not fully created in the neocortex after exposure to only one example of it, although some small modifications of the neural connections may be made.

The second function of the neocortex (at least of the visual part of it) is to respond extremely rapidly to incoming signals. To do so, it uses the categories it has learned and tries to find the combinations of active neurons that, on the

has seen to it that they form as fast as possible; otherwise, no animal could survive. The brain is handicapped in forming neuronal coalitions rapidly because, by computer standards, neurons act very slowly. The brain compensates for this relative slowness partly by using very many neurons, simultaneously and in parallel, and partly by arranging the system in a roughly hierarchical manner.

If visual awareness at any moment corresponds to sets of neurons firing, then the obvious question is: Where are these neurons located in the brain, and in what way are they firing? Visual awareness is highly unlikely to occupy all the neurons in the neocortex that happen to be firing above their background rate at a particular moment. We would expect that, theoretically, at least some of these neurons would be involved in doing computations—trying to arrive at the best coalitions—while others would express the results of these computations, in other words, what we see.



WILLIAM JAMES, the father of American psychology, observed that consciousness is not a thing but a process.

basis of its past experience, are most likely to represent the relevant objects and events in the visual world at that moment. The formation of such coalitions of active neurons may also be influenced by biases coming from other parts of the brain: for example, signals telling it what best to attend to or high-level expectations about the nature of the stimulus.

Consciousness, as James noted, is always changing. These rapidly formed coalitions occur at different levels and interact to form even broader coalitions. They are transient, lasting usually for only a fraction of a second. Because coalitions in the visual system are the basis of what we see, evolution

Fortunately, some experimental evidence can be found to back up this theoretical conclusion. A phenomenon called binocular rivalry may help identify the neurons whose firing symbolizes awareness. This phenomenon can be seen in dramatic form in an exhibit prepared by Sally Duensing and Bob Miller at the Exploratorium in San Francisco.

Binocular rivalry occurs when each eye has a different visual input relating to the same part of the visual field. The early visual system on the left side of the brain receives an input from both eyes but sees only the part of the visual field to the right of the fixation point. The converse is true for the right side. If these two conflicting inputs are rivalrous, one sees not the two inputs superimposed but first one input, then the other, and so on in alternation.

In the exhibit, called "The Cheshire Cat," viewers put their heads in a fixed place and are told to keep the gaze fixed. By means of a suitably placed mirror [see box on next two pages], one of the eyes can look at another person's face, directly in front, while the other eye sees a blank white screen to the side. If the viewer waves a hand in front of this plain screen at the same location

in his or her visual field occupied by the face, the face is wiped out. The movement of the hand, being visually very salient, has captured the brain's attention. Without attention the face cannot be seen. If the viewer moves the eyes, the face reappears.

In some cases, only part of the face disappears. Sometimes, for example, one eye, or both eyes, will remain. If the viewer looks at the smile on the person's face, the face may disappear, leaving only the smile. For this reason, the effect has been called the Cheshire Cat effect, after the cat in Lewis Carroll's *Alice's Adventures in Wonderland*.

Although it is very difficult to record activity in individual neurons in a human brain, such studies can be done in monkeys. A simple example of binocular rivalry has been studied in a monkey by Nikos K. Logothetis and Jeffrey D. Schall, both then at M.I.T. They trained a macaque to keep its eyes still and to signal whether it is seeing upward or downward movement of a horizontal grating. To produce rivalry, upward movement is projected into one of the monkey's eyes and downward movement into the other, so that the two images overlap in the visual field. The monkey signals that it sees up and down movements alternatively, just as humans would. Even though the motion stimulus coming into the monkey's eyes is always the same, the monkey's percept changes every second or so.

Cortical area MT (which Semir Zeki calls in his article V5) is an area mainly concerned with movement. What do the neurons in MT do when the monkey's percept is sometimes up and sometimes down? (The researchers studied only the monkey's first response.) The simplified answer—the actual data are rather more messy—is that whereas the firing of some of the neurons correlates with the changes in the percept, for others the average firing rate is relatively unchanged and independent of which direction of movement the monkey is seeing at that moment. Thus, it is unlikely that the firing of all the neurons in the visual neocortex at one particular moment corresponds to the monkey's visual awareness. Exactly which neurons do correspond remains to be discovered.

We have postulated that when we clearly see something, there must be neurons actively firing that stand for what we see. This might be called the activity principle. Here, too, there is some experimental evidence. One example is the firing of neurons in cortical area V2 in response to illusory con-

tours, as described by Zeki. Another and perhaps more striking case is the filling in of the blind spot. The blind spot in each eye is caused by the lack of photoreceptors in the area of the retina where the optic nerve leaves the retina and projects to the brain. Its location is about 15 degrees from the fovea (the visual center of the eye). Yet if you close one eye, you do not see a hole in your visual field.

Philosopher Daniel C. Dennett of Tufts University is unusual among philosophers in that he is interested both in psychology and in the brain. This interest is much to be welcomed. In a recent book, *Consciousness Explained*, he has argued that it is wrong to talk about filling in. He concludes, correctly, that "an absence of information is not the same as information about an absence." From this general principle he argues that the brain does not fill in the blind spot but rather ignores it.

Dennett's argument by itself, however, does not establish that filling in does not occur; it only suggests that it might not. Dennett also states that "your brain has no machinery for [filling in] at this location." This statement is incorrect. The primary visual cortex (V1) lacks a direct input from one eye, but normal "machinery" is there to deal with the input from the other eye.

Ricardo Gattass and his colleagues at the Federal University of Rio de Janeiro have shown that in the macaque some of the neurons in the blind-spot area of V1 do respond to input from both eyes, probably assisted by inputs from other parts of the cortex. Moreover, in the case of simple filling in, some of the neurons in that region respond as if they were actively filling in.

Thus, Dennett's claim about blind spots is incorrect. In addition, psychological experiments by Vilayanur S. Ramachandran [see "Blind Spots," *SCIENTIFIC AMERICAN*, May] have shown that what is filled in can be quite complex depending on the overall context of the visual scene. How, he argues, can your brain be ignoring something that is in fact commanding attention?

Filling in, therefore, is not to be dismissed as nonexistent or unusual. It probably represents a basic interpolation process that can occur at many levels in the neocortex. It is, incidentally, a good example of what is meant by a constructive process.

How can we discover the neurons whose firing symbolizes a particular percept? William T. Newsome and his colleagues at Stanford University have done a series of brilliant experiments on neurons in cortical area MT of the macaque's brain. By studying a neuron

The Cheshire Cat Experiment

This simple experiment with a mirror illustrates one aspect of visual awareness. It relies on a phenomenon called binocular rivalry, which occurs when each eye has a different input from the same part of the visual field. Motion in the field of one eye can cause either the entire image or parts of the image to be erased. The movement captures the brain's attention.



in area MT, we may discover that it responds best to very specific visual features having to do with motion. A neuron, for instance, might fire strongly in response to the movement of a bar in a particular place in the visual field, but only when the bar is oriented at a certain angle, moving in one of the two directions perpendicular to its length within a certain range of speed.

It is technically difficult to excite just a single neuron, but it is known that neurons that respond to roughly the same position, orientation and direction of movement of a bar tend to be located near one another in the cortical sheet. The experimenters taught the monkey a simple task in movement discrimination using a mixture of dots, some moving randomly, the rest all in one direction. They showed that electrical stimulation of a small region in the right place in cortical area MT would bias the monkey's motion discrimination, almost always in the expected direction.

Thus, the stimulation of these neurons can influence the monkey's behavior and probably its visual percept. Such experiments do not, however, show decisively that the firing of such neurons is the exact neural correlate of the percept. The correlate could be only a subset of the neurons being activated. Or perhaps the real correlate is the firing

of neurons in another part of the visual hierarchy that are strongly influenced by the neurons activated in area MT.

These same reservations apply also to cases of binocular rivalry. Clearly, the problem of finding the neurons whose firing symbolizes a particular percept is not going to be easy. It will take many careful experiments to track them down even for one kind of percept.

It seems obvious that the purpose of vivid visual awareness is to feed into the cortical areas concerned with the implications of what we see; from there the information shuttles on the one hand to the hippocampal system, to be encoded (temporarily) into long-term episodic memory, and on the other to the planning levels of the motor system. But is it possible to go from a visual input to a behavioral output without any relevant visual awareness?

That such a process can happen is demonstrated by the remarkable class of patients with "blindsight." These patients, all of whom have suffered damage to their visual cortex, can point with fair accuracy at visual targets or track them with their eyes while vigorously denying seeing anything. In fact, these patients are as surprised as their doctors by their abilities. The amount of information that "gets through," how-

ever, is limited: blindsight patients have some ability to respond to wavelength, orientation and motion, yet they cannot distinguish a triangle from a square.

It is naturally of great interest to know which neural pathways are being used in these patients. Investigators originally suspected that the pathway ran through the superior colliculus. Recent experiments suggest that a direct albeit weak connection may be involved between the lateral geniculate nucleus and other cortical areas, such as V4. It is unclear whether an intact V1 region is essential for immediate visual awareness. Conceivably the visual signal in blindsight is so weak that the neural activity cannot produce awareness, although it remains strong enough to get through to the motor system.

Normal-seeing people regularly respond to visual signals without being fully aware of them. In automatic actions, such as swimming or driving a car, complex but stereotypical actions occur with little, if any, associated visual awareness. In other cases, the information conveyed is either very limited or very attenuated. Thus, while we can function without visual awareness, our behavior without it is rather restricted.

Clearly, it takes a certain amount of time to experience a conscious percept. It is difficult to determine just how much time is needed for an episode of visual awareness, but one aspect of the problem that can be demonstrated experimentally is that signals received close together in time are treated by the brain as simultaneous.


A disk of red light is flashed for, say, 20 milliseconds, followed immediately by a 20-millisecond flash of green light in the same place. The subject reports that he did not see a red light followed by a green light. Instead he saw a yellow light, just as he would have if the red and the green light had been flashed simultaneously. Yet the subject could not have experienced yellow until after the information from the green flash had been processed and integrated with the preceding red one.

Experiments of this type led psychologist Robert Efron, now at the University of California at Davis, to conclude that the processing period for perception is about 60 to 70 milliseconds. Similar periods are found in experiments with tones in the auditory system. It is always possible, however, that the processing times may be different in higher parts of the visual hierarchy and in other parts of the brain. Processing is also more rapid in trained, compared with naive, observers.

Because it appears to be involved in some forms of visual awareness, it

To observe the effect, a viewer divides the field of vision with a mirror placed between the eyes (*a*). One eye sees the cat; the other eye a reflection in the mirror of a white wall or background. The viewer then waves the hand that corresponds to the eye looking at the mirror so that the hand passes through the area in which the image of the cat appears in the other eye (*b*). The result is that the cat may disappear. Or if the viewer was attentive to a specific feature before the hand was waved, those parts—the eyes or even a mocking smile—may remain (*c*).





OPTICAL ILLUSION devised by Vilayanur S. Ramachandran illustrates the brain's ability to fill in, or construct, visual information that is missing because it falls on the blind spot of the eye. When you look at the patterns of broken green bars, the visual system produces two illusory contours defining a vertical strip. Now shut your right eye and focus on the white square in the green series of bars. Move the page toward your eye until the blue dot disappears (roughly six inches in front of your nose). Most observers report seeing the vertical strip completed across the blind spot, not the broken line. Try the same experiment with the series of just three red bars. The illusory vertical contours are less well defined, and the visual system tends to fill in the horizontal bar across the blind spot. Thus, the brain fills in differently depending on the overall context of the image.

would help if we could discover the neural basis of attention. Eye movement is a form of attention, since the area of the visual field in which we see with high resolution is remarkably small, roughly the area of the thumbnail at arm's length. Thus, we move our eyes to gaze directly at an object in order to see it more clearly. Our eyes usually move three or four times a second. Psychologists have shown, however, that there appears to be a faster form of attention that moves around, in some sense, when our eyes are stationary.

The exact psychological nature of this faster attentional mechanism is at present controversial. Several neuroscientists, however, including Robert Desimone and his colleagues at the National Institute of Mental Health, have shown that the rate of firing of certain neurons in the macaque's visual system depends on what the monkey is attending to in the visual field. Thus, attention is not solely a psychological concept; it also has neural correlates that can be observed. Several researchers have found that the pulvinar, a region of the thalamus, appears to be involved in visual attention. We would like to believe that the thalamus deserves to be called "the organ of attention," but this status has yet to be established.

The major problem is to find what activity in the brain corresponds directly to visual awareness. It has been speculated that each cortical area produces awareness of only those visual features that are "columnar," or arranged in the stack or column of neurons perpendicular to the cortical surface. Thus, area V1 could code for orientation and area MT for motion. So far, as Zeki has explained, experimentalists have not found one particular region in the brain where all the information needed for visual awareness appears to come together. Dennett has dubbed such a hypothetical place "The Cartesian Theater." He argues on theoretical grounds that it does not exist.

Awareness seems to be distributed not just on a local scale, as in some of the neural nets described by Hinton, but more widely over the neocortex. Vivid visual awareness is unlikely to be distributed over every cortical area because some areas show no response to visual signals. Awareness might, for example, be associated with only those areas that connect back directly to V1 or alternatively with those areas that project into each other's layer 4. (The latter areas are always at the same level in the visual hierarchy.)

The key issue, then, is how the brain forms its global representations from

visual signals. If attention is indeed crucial for visual awareness, the brain could form representations by attending to just one object at a time, rapidly moving from one object to the next. For example, the neurons representing all the different aspects of the attended object could all fire together very rapidly for a short period, possibly in rapid bursts.


This fast, simultaneous firing might not only excite those neurons that symbolized the implications of that object but also temporarily strengthen the relevant synapses so that this particular pattern of firing could be quickly recalled—a form of short-term memory. (If only one representation needs to be held in short-term memory, as in remembering a single task, the neurons involved may continue to fire for a period, as described by Patricia S. Goldman-Rakic in "Working Memory and the Mind," page 110.)

A problem arises if it is necessary to be aware of more than one object at exactly the same time. If all the attributes of two or more objects were represented by neurons firing rapidly, their attributes might be confused. The color of one might become attached to the shape of another. This happens sometimes in very brief presentations.

Some time ago Christoph von der Malsburg, now at the Ruhr-Universität Bochum, suggested that this difficulty would be circumvented if the neurons associated with any one object all fired in synchrony (that is, if their times of firing were correlated) but out of synchrony with those representing other objects. More recently, two groups in Germany reported that there does appear to be correlated firing between neurons in the visual cortex of the cat, often in a rhythmic manner, with a frequency in the 35- to 75-hertz range, sometimes called 40-hertz, or γ , oscillation.

Von der Malsburg's proposal prompted us to suggest that this rhythmic and synchronized firing might be the neural correlate of awareness and that it might serve to bind together activity in different cortical areas concerning the same object. The matter is still undecided, but at present the fragmentary experimental evidence does rather little to support such an idea. Another possibility is that the 40-hertz oscillations may help distinguish figure from ground [see "The Legacy of Gestalt Psychology," by Irvin Rock and Stephen Palmer; *SCIENTIFIC AMERICAN*, December 1990] or assist the mechanism of attention.

Are there some particular types of neurons, distributed over the visual neocortex, whose firing directly symbolizes the content of visual awareness? One



BRIEF FLASHES of colored light enable researchers to infer the minimum time required for visual awareness. A disk of red light is projected for 20 milliseconds (a), followed immediately by a 20-millisecond flash of green light (b). But the observer reports seeing a single flash of yellow (c), the color that would be apparent if red and green were projected simultaneously. The subject does not become aware of red followed by green until the length of the flashes is extended to 60 to 70 milliseconds.

very simplistic hypothesis is that the activities in the upper layers of the cortex are largely unconscious ones, whereas the activities in the lower layers (layers 5 and 6) mostly correlate with consciousness. We have wondered whether the pyramidal neurons in layer 5 of the neocortex, especially the larger ones, might play this latter role.

These are the only cortical neurons that project right out of the cortical system (that is, not to the neocortex, the thalamus or the claustrum). If visual awareness represents the results of neural computations in the cortex, one might expect that what the cortex sends elsewhere would symbolize those results. Moreover, the neurons in layer 5 show a rather unusual propensity to fire in bursts. The idea that the layer 5 neurons may directly symbolize visual awareness is attractive, but it still is too early to tell whether there is anything in it.

Visual awareness is clearly a difficult problem. More work is needed on the psychological and neural basis of both attention and very short term memory. Studying the neurons when a percept changes, even though the visual input is constant, should be a powerful experimental paradigm. We need to con-

struct neurobiological theories of visual awareness and test them using a combination of molecular, neurobiological and clinical imaging studies.

We believe that once we have mastered the secret of this simple form of awareness, we may be close to understanding a central mystery of human life: how the physical events occurring in our brains while we think and act in the world relate to our subjective sensations—that is, how the brain relates to the mind.

FURTHER READING

- PERCEPTION. Irvin Rock. Scientific American Library, 1984.
CONSCIOUSNESS AND THE COMPUTATIONAL MIND. Ray Jackendoff. The MIT Press/Bradford Books, 1987.
COLD SPRING HARBOR SYMPOSIA ON QUANTITATIVE BIOLOGY, Vol. LV: THE BRAIN. Cold Spring Harbor Laboratory Press, 1990.
TOWARDS A NEUROBIOLOGICAL THEORY OF CONSCIOUSNESS. Francis Crick and Christof Koch in *Seminars in the Neurosciences*, Vol. 2, pages 263-275; 1990.
THE COMPUTATIONAL BRAIN. Patricia S. Churchland and Terrence J. Sejnowski. The MIT Press/Bradford Books, 1992.



Hot Potato

Will safety questions curb public appetite for gene-spliced food?

While some companies were focusing on the enormous potential of gene splicing to produce new drugs from modified cells, other entrepreneurs saw in biotechnology the promise of a second "Green Revolution." A group of small start-ups, agricultural chemical giants, seed producers and food processors devoted their efforts to developing plants with new genetic traits of value to consumers and farmers, such as better nutrition and taste, longer shelf life and resistance to diseases and pesticides.

Nearly a decade after the first drug produced by biotechnology received marketing approval from the Food and Drug Administration, genetically engineered foods still await their debut. Agricultural biotechnology firms are finding that the same science that lifts hopes in the clinic raises fears around the kitchen table. Top New York City chefs are loudly threatening to boycott what one wag has dubbed "Franken-food," and groups critical of biotechnology, such as Jeremy Rifkin's Foundation on Economic Trends, promise legal challenges to the FDA.

Apprehensions about the safety of genetically engineered foods began flaring like a grease fire in a deep fryer when the FDA announced in late May that plants modified by genetic engineering will in most cases be treated just like any ordinary garden variety. New plants, whether they are produced through techniques of biotechnology or conventional breeding, do not require any special review to gain market approval, the agency decreed. The FDA also decided that genetically engineered foods need not be labeled as such simply because they are produced by that process. Changes that result in products significantly different from what consumers expect—for instance, high-fiber potatoes or oranges low in vitamin C—will warrant some kind of informational labeling, the FDA says.

Critics charge that the public has the right to know what it is buying. They point out that the FDA currently leaves the responsibility for food safety to producers, and they question the wisdom



JOHN C. SORENSON displays yellow squash genetically engineered by Asgrow to resist viruses. Stunted plants nearby are unaltered. Photo: Peter Yates.

of extending that trust to purveyors of unproved technology. "The FDA is counting on companies' coming to them; otherwise, they're saying you don't have to talk to us," declares Rebecca J. Goldberg, senior scientist at the Environmental Defense Fund (EDF).

The New York-based environmental group has proposed its own ideas for safety and labeling rules under the title *A Mutable Feast: Assuring Food Safety in the Era of Genetic Engineering*. "While I don't think most genes will be hazardous, I do think we have to look for exceptions, not the rule," Goldberg says. The EDF is particularly concerned about plants engineered to contain pesticides; it also fears that the potential for allergy problems will rise as genes are transferred among organisms. Rifkin plans to emphasize allergenicity in his Pure Food campaign as well.

For their part, companies say the technology to insert new genes permits them to do what plant breeders have been doing for thousands of years—only more rapidly and with greater specificity. "Single-gene modifications are actually less likely to cause unexpected negative traits than is traditional breed-

ing because far less genetic material is exchanged," asserts Vic C. Knauf, vice president for research at Calgene in Davis, Calif.

The first genetically engineered vegetable is slated to appear at the produce counter in mid-1993 or early 1994. The Flavor Saver tomato engineered by Calgene contains "antisense" RNA sequences to block production of an enzyme that hastens rotting. Other food crops that various companies are currently field-testing include squash, cantaloupe, cucumbers, potatoes, corn and such oilseeds as cotton and canola [see box on page 162].

Beyond the first wave of bioengineered plants is a "more complex and subtle second generation" that will not approach market for another five to 10 years, says John C. Sorenson, executive director of vegetable research for Asgrow Seed Company in Kalamazoo, Mich. "We're now moving to understand the metabolic basis of plant disease and to interfere with that process," he notes. In addition, some laboratories—for instance, the U.S. Department of Agriculture's Agricultural Research Service in Lubbock, Tex.—are experiment-

Get the RAINFOREST LIBRARY


for just
\$2.95

as your introduction to the
Natural Science Book Club

You simply agree to buy three more books -
at significant savings - within the next 12 months.

Rainforest facts...

- Tropical forests cover only 6% of the earth's land surface, yet shelter up to 30 million species— 90% of our planet's total
- Rainforests contain fully 80% of earth's terrestrial vegetation
- 57 acres of tropical moist forest are cut down every minute of every day



An
\$89.95
Value!

Rainforests are the richest source of life on the planet. This intriguing three-volume set reveals these fragile ecosystems through startling photographs, incredible facts and scientific explanation, and tells how they can be preserved.

Tropical Rainforest: A World Survey of Our Most Valuable and Endangered Habitat with a Blueprint for Its Survival


By Arnold Newman

Follow Newman's stunning odyssey into the richness and diversity of rainforest life. You'll discover the factors that threaten these incredible environments. And what can be done to save them! Includes over 200 magnificent color photographs and illustrations, educational resource list, bibliography, and complete index.

Publisher's Price: \$40.00

THE NATURAL SCIENCE BOOK CLUB is a unique book club devoted to bringing you exciting, refreshingly different books on all areas of the natural sciences. For over twenty years we have brought enthusiasts the books they need to explore the worlds of physics, astronomy, archaeology, wildlife, and much more. All books are available to you at significant savings, with prices starting as low as \$9.95.

MEMBERSHIP BENEFITS: In addition to getting the Rainforest Library for only \$2.95 when you join, you'll also receive discounts on books you choose to purchase. Discounts generally range from 15% to 30% off the publishers' prices, and occasionally even more. • Also, you will immediately become eligible to participate in our Bonus Book Plan, with savings of more than 50% off the publishers' prices. • Moreover, our books are always identical to the publisher's editions. You'll never receive an "economy edition" with inferior paper and bindings from us. • At 3-4 week intervals (15 times per year), you will receive the Natural Science Book Club News, describing the coming Main Selection and Alternate Selections, together with a dated reply card. • In addition, up to three times a year, you may receive offers of Special Selections which will be made available to a group of select members. • If you want the Main Selection, do nothing, and it will be sent to you automatically. • If you prefer another selection, or no book at all, simply indicate your choice on the card and return it by the date specified. • You will have at least 10 days to decide. If, because of late mail delivery of the News, you should receive a book you do not want, we guarantee return postage.



An Introduction to Tropical Rainforests

By T.C. Whitmore

Find out how rainforests function and their value to man. Learn about the history and ecological dynamics of these incredible habitats. Uncover new and fascinating interdependencies among plants, insects, and even mammals, some never before seen by man. Includes 190 illustrations and photographs.

Publisher's Price: \$35.00

Lessons of the Rainforest

Edited by Suzanne Head and Robert Heinzman

Join 24 leading authorities in a variety of fields as they present alternatives to rainforest decimation. Bringing together the insights of biologists, ecologists, historians, activists, even lawyers, these in-depth articles examine the many interrelated issues surrounding tropical deforestation, and offer strategies for slowing the destruction that threatens this fragile ecosystem.

Publisher's Price: \$14.95

Natural Science Book Club

4-CR6

A Newbridge Book Club
3000 Cindel Drive
Delran, NJ 08370-0001

Please accept my application for trial membership and send me the Rainforest Library (00169) billing me only \$2.95, plus shipping and handling. I agree to purchase at least three additional Selections or Alternates over the next 12 months. Savings generally range from 15% to 30% off the publishers' prices. My membership is cancelable any time after I buy these three additional books. A shipping and handling charge is added to all shipments.

No-Risk Guarantee: If I am not satisfied—for any reason—I may return the Rainforest Library within 10 days. My membership will be canceled, and I will owe nothing.

Name _____

Address _____ Apt. _____

City _____

State _____ Zip _____

(Books purchased for professional purposes may be a tax-deductible expense. Offer good in Continental U.S. and Canada only. Prices slightly higher in Canada.)

©Newbridge Communications, Inc.

Scientific American 9/92

Major Developers of Genetically Engineered Foods

Asgrow Seed Company, Kalamazoo, Mich. Field-testing yellow squash, cucumbers and cantaloupes genetically vaccinated against four common plant viruses. Also developing genes for resistance to fungus and bacteria to protect these and other fresh-market vegetables, including lettuce and carrots.

Calgene, Davis, Calif. Flavor Saver tomato with "antisense" RNA that allows vine ripening and longer shelf life will debut commercially, possibly in 1993. Testing canola (rapeseed) engineered to produce laurates and stearates found in tropical oils, for food processing and soap manufacture. Has also modified canola to produce a highly hydrogenated oil for margarine.

DNA Plant Technologies, Cinnaminson, N.J. Field-testing tomatoes that produce the enzyme chitinase, which repels fungi responsible for post-harvest rot, and tomatoes with an enzyme that extends vine ripening. Developing tomatoes with "antifreeze" genes from winter flounder. Also working with Du Pont on a genetically engineered canola that produces a high-temperature frying oil that is low in saturated fat.

Monsanto, St. Louis, Mo. By incorporating genes from soil bacteria, the company is developing a high-solids, low-moisture potato that will absorb less fat during deep frying and a tomato with delayed ripening. Also developing potatoes and tomatoes engineered to resist viral disease as well as potatoes protected against insects. Testing herbicide-resistant cottonseed for oil.

Pioneer Hi-Bred, Des Moines, Iowa Testing sunflowers and canola with altered amino acid profiles to make oils that are more nutritious and lower in saturated fat. Also developing high-methionine soybeans for livestock feed, corn resistant to insects and disease, and alfalfa resistant to alfalfa mosaic virus.

ing with "thermostat" genes believed to determine the narrow ranges of temperature at which plants grow best. Transfer of these traits and others, including resistance to drought, salt and even heavy metals, could help crops grow on land that is currently not arable.

Now that scientists are beginning to reap the benefits of agricultural biotechnology, companies and regulators alike are eager to soothe public concerns. "It isn't that we're not asserting authority and giving everything away," argues Eric L. Flamm, a microbiologist in the FDA's Office of Biotechnology. "It just happens that most of the modifications we know of are trivial from a public health point of view." Although regulators insist genetically engineered plants are not inherently dangerous, neither are the foods inevitably safe, they concede. The same legal provisions allowing the FDA to prevent sales of contaminated shellfish could also be used to yank a genetically engineered product, Flamm says. "Everybody knows—one phone call and it's off the market," he adds.

To fend off critics, the FDA has already issued guidelines on potential problem areas for genetically engineered crops. Companies should first consider the basic characteristics of a new substance, the agency advises. Something substantially similar to the

proteins, fats and carbohydrates already in the diet would generally be considered safe. But in some cases—for instance, when a protein's function is unknown or unusual—that substance might be subject to regulation as a food additive. Two sweeteners being considered for development in plants, thaumatin and monellin, are likely to incur such scrutiny, Flamm says. Unlike most sweet substances, which are carbohydrates, these are proteins.

In addition, Flamm warns that companies should be aware of, and look for, natural toxins likely to be found in both the parent plant and the organism donating genetic material. For example, plant breeders routinely monitor new varieties of squash, celery and potatoes, which are notorious for producing toxic metabolites.

The FDA also wants companies to ensure that engineered plants have the same range of nutrients as do conventional varieties. The agency says it will come down hardest on potential allergens. "Our concept is if you're moving a gene from an allergen, you are moving the allergen," Flamm declares. Plants modified with DNA from wheat, nuts, legumes, milk, eggs and shellfish will require labeling unless companies can conclusively prove the new food inherits no allergenicity, he says.

Some developers of genetically engi-

neered plants do not object to labeling requirements. Calgene, for one, plans to highlight the fact so consumers know why Flavor Saver will cost three to four times the going rate for a store-bought tomato. But labeling agricultural commodities, such as grains, would present problems, says Edward W. Raleigh, Du Pont's manager of biotechnology regulatory affairs. "It's one thing to label an identity-preserved product and quite another to label something that is processed industrially, like corn," he notes. There are so many truckloads, grain elevators and vats between the farm and the consumer that tracking a shipment of gene-spliced corn all the way to a tortilla chip or bottle of oil would be almost impossible.

Moreover, it is still unclear precisely what kind of information consumers want from labeling of genetically engineered food, says Alan Goldhammer, director of technical affairs for the Industrial Biotechnology Association in Washington, D.C. Do they want to know if a gene put into a tomato comes from a soybean? Do they want to know that an enzyme called polygalacturonase is being blocked with antisense RNA? "There is now a way of assuring safety of genetically engineered plants," Goldhammer declares. "It is easy to communicate on a scientific level; our challenge now is to communicate to the society at large," he says.

Even the Agricultural Biotechnology Research Advisory Committee, which reviews field-trial requests submitted to the USDA, could use some tutoring, observes Alvin Young, director of the USDA's Office of Agricultural Biotechnology. "From a scientist's point of view, a gene is just a functional unit that causes certain reactions," Young says. He acknowledges that there are deeper issues to be addressed. "We need some really good thinkers" to assist regulators in their responses to societal concerns about agricultural biotechnology, he emphasizes.

To get the ball rolling, Young has asked W. Steven Burke of the North Carolina Biotechnology Center in Research Triangle Park to prepare a set of draft guidelines. Young intends to use the document, when it is submitted next March, to open up the first major discussion on ethics and personal, cultural and community beliefs surrounding genetically engineered food. If education makes consumers more comfortable with the idea of moving genes, then public awareness programs may well ease acceptance of agricultural biotechnology. Yet logic is rarely a match for instinct, especially when it comes to food. —Deborah Erickson

The Hole Story

Fine-pore membranes remove viruses from biological drugs

One need not look too far into history to find examples of biological drugs that have given people other diseases. Some batches of yellow fever vaccine, for instance, were later discovered to contain avian leukosis virus. Just a few years ago growth hormone derived from human sources transmitted Creutzfeldt-Jakob's disease, a rare degenerative nerve condition. Although such occurrences are rare, the risk of viral contamination is climbing as more therapeutic products are extracted from tissues or grown in mammalian cells.

Membranes with ultra-tiny pores may be just what pharmaceutical firms need to eliminate the last traces of contamination. That is the contention of several U.S. companies now developing the products, including Millipore, Pall and Micron Separations.

Drugmakers currently rely on heat, ultraviolet light, low pH, irradiation and harsh chemicals to kill known potential contaminants such as the viruses that cause AIDS, hepatitis B and hepatitis C. "But to be absolutely sure, the best thing to do is to purify, kill, kill again and purify again," declares Curtis L. Scribner, chief of the hematologic products branch of the Food and Drug Administration. When firms come to the agency seeking marketing approval for biological drugs, he explains, "we look for the ability of their purification steps to remove several thousandfold more virus than they would expect to have in their initial product."

Because viruses are always larger than proteins, membranes with fine pores can retain the contaminants while product flows through. The membranes are most effective when the drug and contaminant are not close in size. Even in the best of circumstances, not all the protein comes through, says Paul J. Sekhri, product manager for the Viresolve membrane developed by Millipore, a bioseparations and analysis company in Bedford, Mass. Millipore works with pharmaceutical manufacturers to ensure that they can recover at least 90 percent of the protein they are making.

The secret of the new membranes is extremely uniform pore size. Traditional membranes, in which pores are formed when solvent evaporates from a polymer, inevitably contain a small number of large pores. They might retain 99 percent of viral contaminants, "but that's not good enough for the FDA," notes

Silver Lining

Serendipity has been an infrequent visitor in the laboratories of high-temperature superconductor researchers struggling to turn the brittle ceramic materials into wires for real-world applications. But she apparently did drop by one day at the Superconductivity Technology Center at Los Alamos National Laboratory.

There researchers were busily perfecting a new wire-manufacturing method they had devised. By packing superconducting ceramic powder into a silver tube and then drawing and heating it, they had managed to bond the materials into a single solid thread. Unfortunately, their wire suffered the same fundamental flaws as had other designs. To conduct electricity without resistance, the wire would have required a cumbersome jacket to carry cooling liquid nitrogen. Nor had they solved the biggest challenge—cracking. Pressing the thread into a tape gave a bit more stability, but the scientists were still far from a commercial product.

"Suddenly we got the idea to turn the powder-in-a-tube design inside out," recalls Dean E. Peterson, director of the Los Alamos center. Instead of stuffing the tube with superconductor, Peterson, along with David Korzekwa, John Bingert and Haskell Sheinberg, decided to use the ceramic as a coating. That way, they could benefit from the natural mechanical strength of a tube and use the hollow core as a passageway for coolant. To protect the semiconductor coating, the team decided to slip on another concentric silver tube; all three layers would be bonded by heating and the whole tube lengthened by drawing. "Until we actually built one, it was just a neat idea we weren't sure would work," Bingert says.


The researchers are now confident that their tube-within-a-tube forms a stronger, simpler wire that promises to perform well in the real world. Because the metal layers protect the ceramic from within and without, the superconductor should be able to withstand the rigors of shipping and installation. Unlike solid wires or tapes, which can be severed by an impact, cracks must completely encircle the tube to halt the flow of current.

The team hopes to boost current-carrying capacity by building up concentric layers, like the rings of a tree. "We can potentially make miles of wire of any diameter," Peterson says. But joining lengths of multilayered wire may be a difficult endeavor, he admits, because any misalignment would cause energy loss.

The team's choice of a bismuth-based ceramic the researchers call Bisco is critical to the tube's success, notes James L. Smith, chief scientist at the Los Alamos center. When the researchers tested yttrium-based ceramic powders, the silver melted before the material formed the planar structure necessary for superconductivity. Other metals such as gold, copper, steel and platinum rendered the Bisco ceramic unable to superconduct. "The fact that Bisco and silver get along so well is a gift of nature," Smith declares.

The tubes have many potential applications, the workers say, ranging from delivering current to powerful miniaturized motors to wiring up superconducting magnets for maglev transportation. "We expect companies will come up with other ideas and improvements," Korzekwa offers. If the work can be commercialized successfully, it appears that superconducting wire, once a wisp of a dream, will be round for a while.

—Deborah Erickson



SUPERCONDUCTING WIRE manufactured at Los Alamos National Laboratory benefits from tubular geometry. Photo: Fred Rick.

Jerold M. Martin, vice president of marketing for Pall Ultrafine Filtration Company in East Hills, N.Y.

Millipore's Viresolve is different enough to have won a patent. The composite of two membranes—a supporting layer with a fine skin cast over it—is also the first validatable product designed specifically for virus removal. Users verify the membrane's performance by presenting it with two immiscible fluids, Sekhri explains. The ra-

tio of fluids that pass through shares a one-to-one correlation with the amount of contaminant that will be removed.

"The trick is going from biotechnology to bioprocessing," says David W. Hann, vice president of marketing and sales for Micron Separations in Westborough, Mass. Certain products that could benefit from viral screening—for instance, blood plasma and its derivatives—are produced in volumes of hundreds of liters, Hann points out. But

most of the market for virus-removing membranes will be biopharmaceuticals given in small doses. A one-liter batch of interferon, for example, has a market value of \$1.5 million. Millipore's Viresolve currently accommodates runs of 20 to 30 liters.

Still, this fall Millipore plans to offer industry 10-foot-square membranes for large-volume applications. Sometimes, going with the flow requires a good deal of effort. —Deborah Erickson

Underground Allies

Public and legislative pressure to reduce use of chemical pesticides has companies scrambling to develop alternatives. Biosys in Palo Alto, Calif., has come up with one that wriggles. The firm is breeding microscopic nematodes to battle insects underground, where 30 percent of the world's major agricultural enemies spend at least part of their lives. Some types of nematodes are themselves destructive parasites of plants, but the *Steinernema* strain Biosys has selected feeds only on grubs and larvae that damage crops.

Most biological pesticides, such as those based on bacteria, are inactive until an insect comes along to eat them, observes Venkat S. Sohoni, chief executive officer of Biosys. But the tiny, wormlike organisms his company began selling this year seek out their prey, guided by the carbon dioxide emissions of soil-borne insects. After entering through a body opening, the nematodes cause a lethal infection, assisted by symbiotic bacteria that also aid digestion. The host dies in a day or two and continues to provide food and lodging for several generations of nematodes. When the food is gone, the nematodes go forth in search of the next meal. They will travel as far as 12 to 14 inches, never harming earthworms, whose cuticle is impenetrable to them. "If there is no pest, there is no progeny," Sohoni explains.

Researchers have known about beneficial nematodes and the bacteria they carry for 50 years, but no one be-

fore Biosys had been able to solve the breeding and scale-up problems necessary to create a marketable product. Commercializing the tiny predators "required looking at the world from the nematode's point of view," says Milton J. Friedman, the firm's vice president for research. The creatures are far more sensitive than are bacteria to temperature and pH, and they need more oxygen than do cultured cells. Because nematodes reproduce through male-female partnering, they require a fairly calm environment to couple. To complicate matters further, nematodes are pathogenic to insects only during the so-called infective juvenile stage of their development, which takes 16 to 18 days to reach.

When Biosys began commercialization, "one of the first things we did was sit around thinking what we'd make if a nematode came over to dinner," Friedman recalls. The research team decided on something like pancake batter. The emulsion, which includes raw egg yolks, soy flour and lecithin, supplies the desired fat and protein. The thick mixture can be churned throughout a fermenter, incorporating oxygen without becoming too turbulent in a worm's immediate environment. The billions of infectious nematodes strained from a 75,000-liter fermenter are enough to cover 5,000 acres of cropland with some 250 organisms per square inch.

At the moment, shipping and storing the living, breathing nematodes is accomplished by immobilizing them in a gel the consistency of solidified rubber cement. By preventing movement, the matrix reduces their oxygen needs and keeps the product fresher, longer. By next planting season, Biosys hopes to have developed a more convenient slurry formulation. The firm is also using DNA analysis to develop strains more tolerant of ultraviolet light.

Over the next five years, many financial analysts expect Biosys, which was founded in 1983 with initial funding from Silicon Valley pioneers William Hewlett and David Packard, to capture a good bit of the \$2-billion global market for antidotes to subterranean pests. The firm is courting such niche markets as citrus and cranberry farms. Meanwhile Ortho distributes the nematodes for consumer lawns and gardens, and Ciba-Geigy sells them to commercial growers of turf and ornamental plants.

A third big company is becoming familiar with Biosys technology—corn processor Archer Daniels Midland has a contract to produce the nematodes at its new fermentation facility through 1996. "It's possible that we might be a takeover candidate," Sohoni hints. If nematodes do as well as expected against conventional pesticides, it is quite likely a corporate giant will find Biosys as tasty as nematodes find grubs. —Deborah Erickson



VENKAT S. SOHONI of Biosys peers into a bioreactor full of nematodes. Photo: Douglas L. Peck.

**THIS SEASON
WE'LL POISON
A VIOLINIST
CREATE A
MONSTER
AND WATCH A
CATERPILLAR
TURN INTO A
CHICKEN**

And we have the videotape to prove it. Watch each episode of Scientific American Frontiers on PBS for stories you have to see to believe. Meet the concert violinist injected with poison to relieve his muscle disorder. Learn the secrets behind the making of the silver screen's scariest monsters. And see the Panamanian caterpillar that screams for help when attacked. In fact, the only thing more incredible than our stories is the fact that they're all true. Sponsored by GTE Corporation.



YOU JUST CAN'T MAKE THIS STUFF UP.



MINNEAPOLIS AFTER DARK is shown in this image from a Honeywell infrared camera that is mounted on a car (left).



The car lights in the distance, as they appear through the windshield, can be seen in the photograph at the right.

Defensive Driving

Declassified infrared sensors may be an automotive option

The U.S. Army's M1 tank is equipped with forward-looking infrared detectors for spotting targets and enemy forces. During the Persian Gulf War, FLIRs, as they are called in military parlance, permitted the Allied tanks to roll into combat in pitch-blackness. The army buys these exotic thermal-imaging devices at costs that can climb to well above \$100,000 per unit.

Two companies are beating electronic swords into plowshares. Texas Instruments and Honeywell are readying infrared systems for the commercial market. By the turn of the decade, these systems might be sold in a consumer appliance store for the price of a camcorder. They could be used in night-vision systems for automobiles, home security monitors and devices that ferret out insulation leaks.

For more than five years, the companies have held separate contracts with the army and other federal agencies to develop a new generation of lightweight, inexpensive infrared cameras that might one day be mounted on the barrel of an M16 rifle. The army's night vision and electro-optics directorate recently declassified the program in hopes that a large civilian market for these systems will drive down the price of the cameras and because work on these systems is also progressing in Great Britain and Australia.

Borrowing on their expertise in integrated-circuit manufacturing, Texas Instruments and Honeywell were able to build a single microchip bearing 80,000 thermal detector elements, each of which can pick up a variation in tem-

perature (infrared radiation) of a tenth of a degree Celsius or less. The devices are sensitive enough and the contrast between the thermal pixels sharp enough to yield a ghostly image that can reveal facial features, show the arteries in the neck or produce "thermal prints," pictures of heat traces left after touching a book or another object.

Still, the cameras produce only a tenth of the thermal sensitivity provided by the expensive sensors mounted on an M1 tank or military aircraft. These so-called quantum detectors are built of materials such as mercury cadmium telluride. Unlike the soon-to-be commercial cameras, mercury cadmium telluride detectors require costly and relatively unreliable cryogenic systems to eliminate thermal noise. "The number of dollars that have gone into making this older technology affordable is unbelievable—and we still don't have an inexpensive system," says Norman A. Foss, who heads sensor research and development at Honeywell.

Honeywell's device is one of the first near-commercial uses of a relatively new fabrication process called surface micromachining. Employing the photolithographic, etching and other techniques of integrated-circuit manufacturers, surface micromachining sculpts silicon into suspended structures on a microchip. Honeywell etched an array of 336 by 240 pixels out of vanadium oxide. Measuring about 50 microns on a side, each pixel is positioned between supporting structures by tiny posts that channel a current to and from the detector. The arrangement also provides thermal isolation.

Infrared lenses focus the image on the detector array. A transistor on the same chip, one below each pixel, turns an electric current on and off 30 times a second. A change in incident thermal radiation on a pixel alters the resistance of the vanadium oxide, which

in turn modifies the flow of the current.


The technology chosen by Texas Instruments relies on a barium strontium titanate ceramic. A laser—the same type used in semiconductor facilities to pare down the size of resistors and capacitors—etches out the individual pixels. The ceramic is a ferroelectric material, one on which an electrical charge changes when exposed to either an electric field or—in the case of this chip—variations in temperature. When a pixel heats up or cools down, the change in charge, or capacitance, is sensed by various circuit elements.

For the moment, the Honeywell manufacturing process is potentially less costly than laser etching because it relies on standard chip-making methods. But Texas Instruments is developing a similar production approach, and its camera is closer to the commercial market. The company has compressed lenses, detectors and signal-processing electronics into a single four-pound box. The Honeywell system consists of several small boxes connected by wires.

Both Texas Instruments and Honeywell are working closely with a major automobile company, although neither will provide additional details. (Texas Instruments has a working agreement with General Motors and the automaker's defense subsidiary, Hughes Aircraft.) A camera with a dashboard-mounted display would not only double the field of view afforded by automobile headlights, from about 15 to 30 degrees in the horizontal axis, but also allow a driver to see several times the distance ahead, from 500 to, at most, over 3,000 feet.

Interest is also coming from insurance companies, which could use handheld cameras to check house walls for "hot spots" caused by faulty electrical wiring. Infrared cameras could also be useful in home security systems, drug interdiction, police work and border

PREMIER EDITION



The VINTAGE SCIENTIFIC AMERICAN 1993 WALL CALENDAR

Featuring Artwork and Stories of Ingenuity
and Invention from the Pages of
SCIENTIFIC AMERICAN in the 19th Century.

- ◆ Great for home or office
- ◆ Pages suitable for framing
- ◆ The perfect gift
- ◆ Oversized — 13" x 22" open
- ◆ Ample room for daily notes

ONLY
\$9.⁹⁵

The premier edition of the VINTAGE SCIENTIFIC AMERICAN WALL CALENDAR features fascinating facts and impressive engravings of America's greatest technological triumphs:

- ◆ The building of the Brooklyn Bridge
- ◆ The construction of the Statue of Liberty
- ◆ The first "vertical merry-go-round" built by George Ferris
- ◆ Electric motor vehicles
- ◆ Advancements in steam locomotives

Fill out and mail the form below to explore the archives of SCIENTIFIC AMERICAN. Most pictures and stories haven't been seen for over 100 years—and now they are faithfully recreated on replica parchment for your enjoyment all year round.



CUT ALONG LINE AND MAIL TODAY!

Order now in time for the holidays!

WC3SA9

YES! Please rush me _____ copies of The VINTAGE SCIENTIFIC AMERICAN 1993 WALL CALENDAR for \$9.95* plus \$1.50 postage and handling.

Send orders with payment to:

SCIENTIFIC AMERICAN

1993 Wall Calendar
415 Madison Avenue
New York, NY 10017-1111

NAME (PLEASE PRINT) _____

ADDRESS _____

APT _____

CITY _____

STATE _____

ZIP _____

☐ My check/money order for \$_____ is enclosed.

Charge my: ☐ Visa ☐ MasterCard

CARD NO _____

EXPIRES _____

SIGNATURE _____

*Add appropriate sales tax in: CA, IA, IL, MA, MI, NY. In Canada, add GST (No. R127387652). Outside North America, add additional \$2.50 for surface mail. Allow 4-6 weeks for delivery. Photocopies of order form accepted.

patrols. In fact, the army agonized over declassification because these devices might find their way into the hands of drug dealers or terrorists. "Drug dealers have more money than police departments," comments James E. Miller, a physicist with the army's night vision and electro-optics directorate.

Further shrinking the sensors and electronics might lead to infrared glasses that would make it possible to take a stroll on the blackest of nights. Existing night-vision goggles have a drawback. They enhance only moonlight or other low-illumination sources. Despite the depiction in the popular movie "The Silence of the Lambs," these goggles cannot function in total darkness. A pair of infrared glasses, though, would let a wearer extract an eerie light from the heat of the night.

—Gary Stix

See-Through View

Virtual reality may guide physicians' hands

The age-old dream of seeing through solid objects, a notion well ensconced in comic books and science fiction, has now inched a step closer to reality. Or rather a virtual reality. Henry Fuchs, a professor of computer science at the University of North Carolina at Chapel Hill, and two graduate students, Mike Bajura and Ryutarou Ohbuchi, have drawn on the technology for creating simulated environments controlled by a computer to develop what may be a precursor to "x-ray vision" glasses for physicians.

The view so far is not x-ray, but ultrasound. The investigators took the stock-in-trade of virtual reality, a head-mounted display, to superimpose ultrasonic images of a fetus onto a live video image of a pregnant woman's abdomen. The video, on a liquid-crystal screen within the helmetlike display, shows what the viewer would normally see. The ultrasound image, placed over the video image of the abdomen, provides a window into the body.

With such a system, a physician could gain an accurate spatial perspective of the position of a probe inside the body. To perform amniocentesis, for example, a doctor must make the difficult visual correlation between the sonogram displayed on a bedside video monitor and the actual position of a needle as it enters the uterine cavity and moves into the amniotic sac. Similarly, physicians performing a needle biopsy could be more accurately guided by peering through the ultrasonic porthole.

This technique differs from most earlier forms of virtual reality in which the observer looks at a fantasy world of three-dimensional graphic images generated by a computer.

Fuchs and his graduate students have taken virtual reality one step further by adding real-world images. In the first test, on a volunteer in her 38th week of pregnancy, a researcher wearing the display observed the woman on the video while a technician moved an ultrasonic scanning instrument over her abdomen to produce two-dimensional images of the fetus. The ultrasonic images were projected onto the video in the proper spatial orientation. Coordination of the imagery was possible because a magnet-

ic sensor on the ceiling tracked the position of the researcher's head and the ultrasonic scanning device. The different images, with their correct spatial coordinates, were combined by a specialized graphics-processing computer.

A more refined version of such a system could be a valuable tool for physicians. But the North Carolina team must first address the difficulties of fusing different kinds of live imagery. One problem with the early system was that the ultrasound picture appeared to be pasted artificially on top of the abdomen. Fuchs and his students responded by creating a "virtual hole" in a subject's body. A sonogram can now reside within what looks like a recessed pit, which is constructed with three-dimensional graphics.

A viewer looks at the mixed video-ultrasonic picture of the subject only through the left eye (the right liquid-crystal screen on the display is blank). The display will become more realistic when a second camera is placed on the head-mounted display in order to produce stereoscopic imagery. The ultrasound depiction also came from a two-dimensional scanner, although work on three-dimensional ultrasound is moving ahead at other academic institutions, such as Duke University. What the viewer sees in the ultrasound picture is a narrow slice of gray imagery on a black background that is suspended in space. "It's like looking through a keyhole," Fuchs says.

Other needs are improved head-mounted display resolution and a better sensor tracking system, especially one able to predict where the viewer may be about to move the head so that the ultrasound image moves smoothly along with the video.


Better and faster imagery will also come with continued improvements in the already blazing speed of the specialized graphics computer used to create the images. The University of North Carolina already has one of the world's most powerful computers devoted to presenting graphic images in real time. Called Pixel-Planes 5, it is a parallel computer that contains more than 320,000 one-bit processors.

See-through displays may also prove their worth outside the medical field. A composite image, from video and millimeter-wavelength radar, could see through the smoke of a burning building to find someone trapped inside. Graphic elements superimposed onto the fuselage of an aircraft could show a service technician the correct placement of hydraulic lines or electrical wiring. So virtual reality might be used to enhance the real thing.

—Gary Stix



WORLD WITHIN STANDS OUT by using a head-mounted display (left) that might one day let a physician look through an expectant mother's abdomen by merging ultrasound and video imagery (above).



Economic Growth Factors

To hear this election season's campaign speeches, it sounds as if President George Bush and Governor Bill Clinton agree on at least one point: an expanding economy is all-important. Clinton has made growth the centerpiece of his package of economic goals. Political pundits warn Republicans that Bush will have a dour November if he does not find ways to pump up the gross domestic product.

Economists, in contrast, have not hopped en masse onto that bandwagon. "Growth itself shouldn't be a goal," demurs Allan H. Meltzer of Carnegie Mellon University. Adds Samuel Bowles of the University of Massachusetts at Amherst: "A trite question, but one worth asking is: Growth of what and for whom?"

These economists point out what the slogans omit. Designing policies that will bolster people's quality of life is tricky business. "The attractiveness of growth is that it supports the optimism that the future will be better than the past," Bowles observes. But increasingly economists are clashing over what strategies can fulfill that vision.

As Rudiger W. Dornbusch of the Massachusetts Institute of Technology points out, most economists were educated to believe that steady improvements in growth—as measured by a robust gross domestic product—served all, albeit not equally. Many economists looked to the "rising tide" of growth to lift all boats. "The old view would have been that trickle-down works," Dornbusch says.

In the wake of the past decade, Dornbusch believes economists must reassess that faith in rising tides. During the 1980s, "growth had a savage redistribution effect on the economy," he suggests. "Growth per se isn't enough."

Instead, Dornbusch argues, policies should aim to build future resources. "That means education is important," he says. "You need a better and more explicit focus on not leaving people off the bus." Improving workers' skills and bolstering health care policies are also on his list. "Our poor economic performance is not the result of taxation but of an aggressive lack of attention to education and skills," he asserts.

Although such points may be on every

politician's lips this season, Dornbusch is more willing to own up to the costs of such programs. "If money is intelligently spent on education, I wouldn't worry too much about the budget deficit," he says. The national debt now amounts to about 50 percent of the GDP, he points out; it topped 120 percent in 1951. He underlines the need for fundamental plans to improve education: "The worst would be to increase teachers' pay and do nothing else."

Other economists argue that inadequate attention to growth—and a mushrooming deficit—is responsible for the sluggish U.S. economy. "Policies since 1980 have been well constructed to discourage growth in productive capacity," suggests Henry J. Aaron of the Brookings Institution in Washington, D.C. Because of slow growth "people have gradually been coming to the realization that their incomes are not increasing and

Politicians proclaim growth is a key goal, but economists harbor doubts.

they're not likely to outstrip their parents," he maintains.

The government's inattentiveness to training workers, educating children and supporting science and technology is particularly disturbing, Aaron says. "There are competing theories about what's most important, but advance of science and technology and innovative management are on everybody's list."

The implications of even a modest boost in growth, Aaron suggests, are dramatic. Should growth rise from the annual 2 percent watermark to an average rate of 2.5 percent over 20 years, national income would expand by \$900 billion more than it would otherwise. Such higher growth would mean fewer losers and many fewer large losers, he says. "It's not accidental that social discontent and divisiveness have increased as growth has slowed," he asserts. "Growth makes it easier, not harder, to deal with inequality and poverty."

That not all people benefit from rising economic tides is true but hardly unique

to the 1980s, Meltzer counters. "People not in the labor force, in a 'culture of poverty,' have been around for a long time," he says. Instead he advocates that the government loosen further the policies that constrict consumers' personal choices to save or to spend.

Both the tax code and health care subsidies exemplify such stumbling blocks, Meltzer suggests. Because depreciation is not pegged to rising inflation, investors are reluctant to make long-term investments, he says. Americans also overindulge in health care services, he argues, because "we're giving away health care, so no one can see the cost of it. If we put less emphasis on preserving the life of the aged and more on youngsters and mothers, we'd be using our resources better," he contends.

Despite gloomy prognostications, the U.S. economy is growing, Meltzer maintains. "Our children and grandchildren will be richer than we are. We're the poor relations of our children," he declares. "We shouldn't do things that subsidize them, and we shouldn't penalize them."

Such arguments encounter stiff rebuttals. There are fundamental differences between this and previous generations, Bowles insists. The belief that an economy can grow endlessly is crumbling, he observes. Limited growth should force people to confront the problems caused by inequities in wealth.

One result of the growth in inequity since the 1970s, Bowles states, has been what he calls "invidious consumption," an appetite for consuming noticeably more than the Joneses. Rather than the reasonable aim of boosting the productivity of labor—more output per hour—people have mistakenly seen growth as an increase in total output, he declares.

Bowles points to a curious historical note: two traditions of "growth" converged in the late 1700s—Adam Smith's theories of economic growth and the ideas about personal development that emerged from the Enlightenment. "The idea that economic growth should provide the inputs to human growth is morally and environmentally defensible," Bowles says. But in recent years, he adds, society and policies have treated economic growth as the goal. "This inversion of ends and means," Bowles asserts, "lies at the root of our problems." —Elizabeth Corcoran

Jane, at age 8, worked on the tower puzzle; this led her to the exponential function $2^x - 1$ and its graph (Ch.8).



Calculus By and For Young People (ages 7, yes 7 and up) by Don Cohen

- 1 Book: ISBN 09621674-1-x; \$13.95 (reviewed in the Dec. '88 *Scientific American* & Oct. '89 *Math Teacher*..)
- 2 Worksheets: ISBN 09621674-5-2; \$22.95
- 3 VT#1: Infinite Series; ISBN -2-8; 24 min.
- 4 VT#2: Iteration of Inf. Seq's.; ISBN -4-4; 38 min. Video tapes \$42.95 each, go with Ch. 1 & 8; with 6 and 7 yr. olds.

All 4 items ordered together: \$107.

M/C, V, s&h incl. in US; others add 20% USD


Don Cohen, 809 Stratford Dr.,

Champaign, IL 61821; Tel: 217-356-4555

Fax: 217-356-4593

BOOKS ON TAPE®

World's Largest Selection of Audio Books



Box 7900-SA, Newport Beach, CA 92658

Reg. U.S. Pat. & T.M. Off.

More than 2,500 best sellers on cassette, all full length. Rentals by mail. Prepaid delivery and return. Your satisfaction guaranteed. Call (800) 626-3333 toll-free for color brochure. Free.

(800) 626-3333



"BrainMaker is the most fascinating computer software I've ever seen."
-John C. Dvorak
PC Magazine

Neural networks simulate the brain's ability to learn by example. BrainMaker is a complete neural network development system that will find patterns and hidden relationships in your data on any PC-compatible or Macintosh.

Every day, BrainMaker users solve problems in stock and commodity trading, biotechnology, medicine, physics, and astronomy. To build a neural network, just arrange data in rows and columns. Training and testing are automatic - BrainMaker tries every possibility automatically.


Call **800-284-8112** to order BrainMaker or a free catalog. BrainMaker is \$195; Professional is \$795. 30-day money back guarantee. Free text *Introduction to Neural Networks and Expert Systems*.

California Scientific Software
916-478-9040 sales

10024 Newtown Rd, Nevada City, CA 95959

RockWare

The Scientific Software Catalog



**the source
for affordable
scientific software**


**Over 300 programs!
Free catalog available!**

4251 Kipling St., Suite 595
Wheat Ridge, CO 80033
(303) 423-5645 • FAX (303) 423-6171
RockFAX (303) 423-7112
(24 hr FAX product info service)



RockWare
Scientific Software

Intuitive Computer Mathematics



Theorist® is a powerful symbolic algebra and graphing program with a graphical user interface that makes mathematics easy to use on the computer. If you need to incorporate typeset quality equations into your documents, try our easy-to-use equation editor, Expressionist®.

"Theorist... surpasses the highly rated Mathematica... in interface and execution."

- MacUser magazine, Editor's Choice Award, Best Math Program of 1989.

For more information, please call:

1-800-UC-R-MATH

Prescience

939 Howard St. #105, San Francisco, CA 94103

The RAND Journal of Economics

The RAND Journal of Economics is a leader in the field of microeconomics, focusing on industrial organization, contracts, and regulation. The peer-reviewed Journal published by RAND includes topics on market structure, entry and exit, collusion, R&D, patents, innovation, pricing, product and employee liability, health economics, regulatory mechanisms, and the effectiveness of regulation.

Subscriptions for 1992 are \$50 for individuals, \$142 for institutions. Foreign subscribers please add \$15 surface mail postage, \$28 air mail postage. Please send subscription requests to: The RAND Journal of Economics, P.O. Box 1897, Lawrence, KS 66044-1987.

Name _____

Street Address _____ Apt/P.O. Box _____

City _____ State/Province _____

Zip _____ Country _____

Payment: ☐ Bill Me ☐ Enclosed (Checks must be in U.S. funds and drawn on a U.S. bank)

Charge my: ☐ VISA ☐ Master Card

Exp. date _____

Card No. _____

Signature _____

Automated Bibliographies

EndNote Plus builds bibliographies in research papers automatically. It maintains a database of up to 32,000 references and supports searching and other tools for organizing references.

EndNote Plus comes with pre-defined styles such as Science, Nature and APA. It also allows the user to specify new styles. DOS and Mac versions available. Works with Microsoft Word, WordPerfect and other word processors. For information, call 510-649-8176 or fax your request to 510-649-8179.

"Recommended."
Byte Magazine, July 1992

"Top Choice"
MacUser Magazine, September 1991



Niles & Associates, Inc.
2000 Hearst Ave. # 200
Berkeley, CA 94709

1-800-554-3049

SCIENTIFIC AMERICAN
PROPOSES A UNIQUE
OPPORTUNITY FOR
YOUR ADVERTISING

MARKETPLACE
offers 1/6 to 1 page units.
Reader Service included.

For extensive visibility and brand equity with influential Scientific American readers, contact:

Tonia Wendt
Scientific American
1554 S. Sepulveda, Ste. 212
Los Angeles, CA 90025
Phone: (310) 477-9299
Fax: (310) 477-9179

December is the only remaining **MARKETPLACE** section in 1992. Don't miss this exciting advertising opportunity!



**HEWLETT
PACKARD**
*puts the computer
where it belongs!*



The amazing HP 95LX
1MB PC: The first Palmtop computer that comes complete with Real DOS, Real Lotus, and 1 Megabyte of RAM Built-in! In our catalog you'll find both HP 95 versions (Half Meg & 1 Meg), and the most comprehensive list of software, bookware, and accessories, including Derive, PC links, application manuals, and memory cards. You'll see all of the HP Handhelds with clear and definitive descriptions, and receive personalized technical support.

Send for FREE Catalog:
EduCALC
27953 Cabot Rd.
Laguna Niguel, CA 92677
(714) 582-2637

Brain transplant while-u-wait

Give your Excel™ spreadsheet a new brain! Integrate *Neuralyst™* Neural Network software with Excel on Macintosh™ or Windows™ systems while retaining full access to Excel features.

Neuralyst analyzes data sets or time series in your spreadsheets to learn their varied patterns and complex inter-relationships. Once trained, *Neuralyst* becomes an "expert" advisor for your use.

Neuralyst is \$195

New Version 1.3 available.
More capacity, more graphics!

Call to order or for more information
Visa/Mastercard accepted



EPIC
SYSTEMS GROUP

3814 E. Colorado Blvd, Suite 101
Pasadena, CA 91107
Ph: (818)564-0383 FAX: (818)564-0322


COMPUTERS IN SCIENCE



Virtual Reality




Satellite Imaging



The Solar System



Computer Animation




Chaos/Artificial Life



Medical Imaging



Visualization



Fractals

A free 96 page Catalog of
Software, Videotapes,
Books, CD-ROMS, & more...

write to:
MEDIA MAGIC
P.O. BOX 598-SA
Nicasio, CA 94946

or circle our number on the reader service card

**Want to
brush up
on a
foreign
language?**




With Audio-Forum's intermediate and advanced materials, it's easy to maintain and sharpen your foreign-language skills.

Besides intermediate and advanced audio-cassette courses – most developed for the U.S. State Department – we offer foreign-language mystery dramas, dialogs recorded in Paris, games, music, and many other helpful materials. And if you want to learn a new language, we have beginning courses for adults and for children.

We offer introductory and advanced materials in most of the world's languages: French, German, Spanish, Italian, Japanese, Mandarin, Greek, Russian, Arabic, Korean, and others. 215 courses in 76 languages. Our 20th year.

Call 1-800-551-6300 for FREE 36-page catalog, or write:

AUDIO-FORUM®
Room F925, 96 Broad Street,
Guilford, CT 06437 (203) 453-9794



Neurons for Computers

Artificial neural networks are computer models inspired by the structure and behavior of real neurons. Like the brain, they can recognize patterns, reorganize data and, most interestingly, learn. The artificial networks are made up of objects called units, which represent the bodies of the neurons. The units are connected by links, which act as the axons and dendrites. The link multiplies the output from a unit by a weighting factor, a value analogous to the connection strength at a synapse. The link then passes the weighted output value to another unit, which sums up the values passed to it by all other incoming links. If the total input value exceeds some threshold value, the unit fires.

Modifications in the firing pattern constitute the learning. In real neurons, learning is thought to occur in the synapses: when the connection strengths between synapses change, the firing behavior of the network changes. In artificial networks the learning occurs when the weighting factors on the links change [see "How Neural Networks Learn from Experience," by Geoffrey E. Hinton, page 144].

Artificial neural networks are made up of three types of units. The input units take in information from the outside world. The output units send out signals

that are visible to the external world. The hidden units act as go-betweens from the input to the output units; they neither receive input directly from the outside nor produce a visible output.

An illustration of a simple network appears below. The numbers in the units indicate threshold values. The values along the connecting lines are the link weights. Note that there can be shortcuts: some input connections can bypass the hidden unit.

If one input unit is presented with 1 and the other with 0, then the input to the hidden unit will be $(1 \times 1) + (0 \times 1) = 1$. But because this value is less than the threshold value, the hidden unit will not fire (that is, it will have an output of 0). The values to the output unit will be $(1 \times 1) + (0 \times (-2)) + (0 \times 1) = 1$, which is greater than the threshold value of 0.5. The output unit will then fire.

You may recognize that the logic involved mimics an "exclusive or" (XOR) gate. Specifically, if just one input unit is given the value of 1, then the network will produce an output value of 1. Otherwise, it will produce 0—that is, it will not fire. You will find that the other three possible input patterns lead to the appropriate output results for an XOR gate as well.


That you can model an XOR gate

with an artificial neural network is not particularly exciting. By carefully selecting the weights and thresholds of a suitably complex network, you can model any logic function. What is exciting about such networks is that you do not have to choose the weights and thresholds. For our XOR gate, we could have started with any weight and threshold. Then, by repeatedly being shown the patterns of inputs and outputs, the network would learn the weights necessary to implement the XOR gate. Even more interesting, the network can generalize what it has learned. For large data sets, it can recognize patterns that it has not seen before.

To create such a network, we need to make a few modifications to our artificial neurons. First, the mathematics used to train the network becomes much easier if we do not have to change both weights and thresholds during the learning. This is actually simple to do. Any unit with a positive threshold, T , and a certain number, n , of incoming links can be replaced by a unit with a 0 threshold and $n + 1$ links, where the extra link has the weight of $-T$ and comes from a unit that always fires (that is, produces an output of 1). This trick is called biasing. Artificial neural networks often introduce the bias unit, which is usually connected to every unit in the network, to change the threshold into a weight. Representing the output of a unit as a sigmoid function also simplifies the math. The sigmoid is just a smooth approximation of a threshold function.

Although there are several ways to train a network, I chose a method from a particular class of algorithms called supervised learning (other types include unsupervised learning and reinforcement learning). In supervised learning the weights of the network are adjusted in a manner that causes the actual outputs of the net to move closer to the desired outputs. One of the most successful of such training methods is the back-propagation algorithm.

Hinton's article gives a more detailed



ARTIFICIAL NEURAL NETWORK shown here represents an "exclusive or" (XOR) gate: the output unit fires only if an input unit is presented with a 1. The numbers along the connections are weights, and those inside the units are threshold values.

DREW VAN CAMP is a computer programmer and researcher at the University of Toronto. He specializes in creating neural-network simulators for various applications.

description of the technique. Briefly, for those who have a tolerance for elementary calculus, you are taking the derivative of a function in order to find the direction that minimizes the network's error. The function most commonly used for the error is the sum of the squared errors of the output units. The box to the right lists the equations and the steps needed to implement the back-propagation algorithm.

My first implementation was an auto-encoder network. Such devices compress patterns. For example, think of a network that has four input units, two hidden units and four output units (a 4-2-4 encoder). When any input pattern is presented to the network, all four input values must be combined into the two hidden units. These two values must then be able to reproduce the original pattern of the four values in the output units. The entire four-valued input pattern thus must be encoded in the two hidden units.

In the 4-2-4 encoder, note that every input is connected to every hidden unit. Similarly, every hidden unit is connected to every output unit. The bias unit is connected to all the hidden and output units (there is no need to connect the bias unit to the input units, as they will just reproduce the inputs). Each of the four patterns I wanted to train the network to recognize has one unit on and the three remaining units off.

Before I started training, I assigned random values between -1 and 1 for the initial weights. When I presented the four patterns to this untrained network, the total error (the value of E) varied between three and five (depending on the initial weights).

Training the network consisted of repeatedly presenting the patterns to the network and updating the weights after each presentation. I decided that the network reached an acceptable performance level when the total error of the four patterns was less than 0.1. Using this criterion, I found it took between 800 and 2,000 presentations before the network learned the task, depending on the step size, δ , for changing the weights. Indeed, one of the trickiest parts in training the network was choosing δ . If the value chosen is too low, the network will take a long time to converge. If the value is too high, the network behavior will be unstable and may never converge. Through trial and error, I came to use a value of 0.5. Although the network did not train quickly, it never blew up.

Once the network was trained, I decided to see what kind of internal patterns it used to encode the presented patterns. The hidden units contained

Training an Artificial Neural Network

A network learns by successive repetitions of a problem, making smaller errors with each iteration. The most commonly used function for the error is the sum of the squared errors of the output units:

$$E = 1/2 \sum (y_i - d_i)^2$$

The variable d_i is the desired output of unit i , and y_i is its actual output, where y_i is the sigmoid function $1/(1 + e^{-x})$. To minimize the error, take the derivative of the error with respect to w_{ij} , the weight between units i and j :

$$\frac{\partial E}{\partial w_{ij}} = y_i y_j (1 - y_j) \beta_j$$

where $\beta_j = (y_j - d_j)$ for output units and $\beta_j = \sum_k w_{jk} y_k (1 - y_k) \beta_k$ for hidden units (k represents the number of units in the next layer that unit j is connected to). Note that $y_j (1 - y_j)$ is the derivative of the sigmoid function [see illustration on next page].

As you can see, the error can be calculated directly for the links going into the output units. For hidden units, however, the derivative depends on values calculated at all the layers that come after it. That is, the value β must be back-propagated through the network to calculate the derivatives.


Using these equations, we can state the back-propagation algorithm as follows:

- Choose a step size, δ (used to update the weights).
- Until the network is trained,
 - For each sample pattern,
 - Do a forward pass through the net, producing an output pattern.
 - For all output units, calculate $\beta_j = (y_j - d_j)$.
 - For all other units (from last layer to first), calculate β using the calculation from the layer after it:

$$\beta_j = \sum_k w_{jk} y_k (1 - y_k) \beta_k$$


- For all weights in the network, change the weight by

$$\Delta w_{ij} = -\delta y_i y_j (1 - y_j) \beta_j$$



INPUT PATTERNS				HIDDEN UNIT OUTPUTS		ACTUAL OUTPUTS			
1	0	0	0	0.03	0.97	0.91	0.10	0.00	0.07
0	1	0	0	0.98	0.96	0.07	0.88	0.06	0.00
0	0	1	0	0.91	0.02	0.00	0.10	0.91	0.06
0	0	0	1	0.03	0.07	0.07	0.00	0.09	0.90

4-2-4 ENCODER NETWORK compresses patterns. After the training period, the network accurately reproduced the input by representing the patterns essentially as binary code (as revealed by the hidden units).



SIGMOID FUNCTION, defined by the equation $y = 1/(1 + e^{-x})$, produces almost the same output as an ordinary threshold (a step function) but is mathematically simpler. Its derivative is $dy/dx = y(1 - y)$.

values such as 0.03, 0.97, 0.98 and 0.07. Essentially, the network had developed a binary code for the patterns.

Finally, I decided to build a network that could learn arithmetic—specifically, the addition of two three-bit binary numbers. The input to the network would be the six binary digits specifying the two numbers to add. The output would be the four digits to which they summed (four digits are necessary in case a number carries over).

Before building this network, I generated a training set of 64 patterns. To be sure that the representations of the numbers in three-bit form were correct, I wrote a short program that did the calculations and output all the training patterns. Next, I had to determine the architecture of the network. Once again, I decided to use a single hidden layer, but I was not sure how many units it should contain. That is because of a

common problem that arises during the training of artificial neural networks: their performance on the training data will always continue to improve (assuming a stable procedure to update the weights). But if you give patterns that are not from the training set, you will notice that the network's performance on these patterns will first improve, then get worse.

The process is called overfitting. It occurs after the network has learned some general rules about the data. As you train the network, it learns more and more about the anomalies in the training set. It then tries to generalize these anomalies to other data and, as a result, produces a large error.

To avoid serious overfitting, the number of weights in a network should be much less than the number of bits required to specify the desired output for all the training examples. For the binary addition task, there should be far fewer than 256 weights (64 patterns times four digits of output). Using this rule of thumb, I decided to try 15 hidden units with six input units, four output units and a bias unit, which gave 169 connections. This network took significantly longer to train than the previous networks. In fact, it took more than 30,000 iterations.

Once I knew the network could be trained, I tried something different. I removed four of the patterns from the training data, randomized the weights and retrained the network on the remaining 60 patterns. Once it was trained, I tested it on the four patterns it had never seen. It produced the correct answers for these patterns. The network had learned to do binary addition.

Many modifications would greatly enhance the network's learning rate. In particular, you could use more complex methods for choosing the direction to move the weights or use line searches to determine how large a step to take. There are also many more complex algorithms to try, including constructive algorithms that add hidden units as they train.

For a copy of the encoder program (written in ANSI C language), send a formatted double-density disk (3½ or 5¼ inches) with a stamped, self-addressed mailer to: *The Amateur Scientist*, Scientific American, 415 Madison Avenue, New York, NY 10017-1111.

Network Parameters and Data Structures

```
/*Network parameters*/
#define NUM_INPUT      4      /*number of input units*/
#define NUM_HIDDEN     2      /*number of hidden units*/
#define NUM_OUTPUT     4      /*number of output units*/
#define NUM_BIAS       1      /*number of bias units*/
#define STEP_SIZE      0.5    /*step size for updating weights*/

/* Network data structures: Unit, Net */
typedef struct Unit {
    double    input;          /*total input*/
    double    output;         /*total output (input through
                               the sigmoid function)*/
    double    target;         /*target output (for output units)*/
    double    beta;           /*error derivative (backpropagated)*/
} Unit;

typedef struct Net {
    Unit      input [NUM_INPUT]; /*layer of input units*/
    Unit      hidden [NUM_HIDDEN]; /*layer of hidden units*/
    Unit      output [NUM_OUTPUT]; /*layer of output units*/
    Unit      bias [NUM_BIAS]; /*layer of bias units*/

    /*Connections between layers (note order of indices)*/
    double    i2h[NUM_HIDDEN][NUM_INPUT]; /*input to hidden*/
    double    h2o[NUM_OUTPUT][NUM_HIDDEN]; /*hidden to output*/
    double    b2h[NUM_HIDDEN][NUM_BIAS]; /*bias to hidden*/
    double    b2o[NUM_OUTPUT][NUM_BIAS]; /*bias to output*/

    FILE      *fpPatterns; /*circular file of patterns*/
    double    error; /*network error*/
} Net;

/*Useful macros: sigmoid (derivative), random, square*/
#define sigmoid(x)      (1.0 / (1.0 + exp(-(double)(x))))
#define sigmoidDerivative(x) ((double)(x) * (1.0 - (x)))
#define random(x)      ((double)rand()/(RAND_MAX))
#define square(x)      ((x)*(x))
```

FURTHER READING

EXPLORATIONS IN PARALLEL DISTRIBUTED PROCESSING: A HANDBOOK OF MODELS, PROGRAMS, AND EXERCISES. David E. Rumelhart and James L. McClelland. The MIT Press, 1988.

Transform Your Library of SCIENTIFIC AMERICAN into a Powerful Research Tool



SCIENTIFIC
AMERICAN
SciDex™



■ Easy to Use ■ Fully Cross-Referenced ■ Article
Abstracts ■ 530 Issues ■ Over 4,300 Feature Articles
Indexed ■ Over 5,000 Authors ■ Print Your Search
■ Over 43,000 Topical Entries ■ Annual Renewal

SYSTEM REQUIREMENTS AND GENERAL INFORMATION: The Macintosh version is available immediately (allow 4-6 weeks for delivery). SciDex™ can be used on any Macintosh (Mac Plus or better) with 2Mb of RAM (4Mb recommended), running under System v6.0.5 or later and a hard disk with 5Mb of free space. SciDex™ comes compressed on 2 high-density 1.4Mb or 4 double-density 800K 3-1/2" disks. Please specify disk format.

The IBM and compatible versions available for shipment in October, running under Windows or MS-DOS. The Windows version requires an IBM PC, AT or 100% compatible (80386 recommended) with 2Mb of RAM (4Mb recommended); Microsoft Windows 3.0 or later; hard disk drive; and an EGA, VGA or higher resolution monitor. The DOS version requires IBM PC, AT or 100% compatible with a minimum of 256K of RAM; MS-DOS 3.0 or later; hard disk drive; and a monochrome, EGA, VGA or higher resolution monitor.

Apple® and Macintosh® are registered trademarks of Apple Computer, Inc. IBM® is a registered trademark of International Business Machines Corp. Windows™ is a trademark of Microsoft Corp. All other brand and product names mentioned are trademarks or registered trademarks of their respective holders. SciDex™ and its contents © 1992 by Scientific American. All rights reserved. Software program (Macintosh version) © 1992 Catherine Cetrangolo.

SCIENTIFIC AMERICAN CUMULATIVE INDEX 1948-1992 ELECTRONIC DATABASE ON COMPUTER DISK

More than four decades of science and technology are at your fingertips with this exciting new product. SciDex™ is the most powerful research tool you can use to harness the wealth of information from the brightest minds in the pages of *Scientific American*.

Perfect for researchers, libraries, students, educators and you—for use in conjunction with your *Scientific American* collection. The information in this index is available from no other source.

Within seconds, SciDex™ will give you the issue and page number you need to find any article you want on any scientific subject. So easy to use—just type in a topic, author, issue date, or title (or even a word from the title or topic), and the article reference appears.

Then you can cross-reference to other articles. SciDex™ includes listings for every feature article in *Scientific American* from May 1948 to June 1992.

All this power is yours for only \$49.95. Price includes full documentation.

GUARANTEE: If your copy of SciDex™ is defective, return it with your registration number within 90 days and we will promptly replace it free of charge.



SCIENTIFIC
AMERICAN
SciDex

415 Madison Avenue
New York, NY
10017-1111

Please send me _____ copies of SciDex™, the *Scientific American* electronic index from May 1948 to June 1992 at \$49.95. Add \$3.00 for domestic shipping and handling.* Corporate orders accepted if accompanied by authorized purchase order. I understand the Macintosh® version is available immediately (allow 4 to 6 weeks for delivery). Windows™ and MS-DOS® versions available in October '92. Select version and disk format below.

Name _____
Organization _____
Address _____
City _____ State _____
Zip _____ Fax: () _____
Tel: () _____ Please Ship:

☐ Macintosh® ☐ Windows™ ☐ MS-DOS® version as
☐ 3-1/2" DS/HD, ☐ 3-1/2" DS/DD
☐ 5-1/4" DS/HD, ☐ 5-1/4" DS/DD disks
☐ My check/money order is enclosed for \$ _____
Charge my ☐ VISA ☐ MasterCard
☐ ACCESS ☐ EuroCard Exp. Date _____

Card No. _____
Signature _____

*Add applicable sales tax for IA, IL, MA, MI, CA, NY, PA. DS/SA9
Outside the U.S. remit \$49.95 in U.S. funds drawn on a U.S. bank or by credit card and add \$5.00 for surface delivery and handling or check here ☐ and add \$15 for air delivery and handling.



Bohr's Century

NIELS BOHR'S TIMES, IN PHYSICS, PHILOSOPHY, AND POLITY, by Abraham Pais. Oxford University Press (Clarendon Press), 1991 (\$35).

Bohr and Einstein were in their... sixties...when I first met them," Abraham Pais writes. "Since I am one (perhaps the last) of those who knew both men rather well personally...I have been asked off and on how... they compared." And then, across three or four warm pages, Pais lists many comparisons between the two lifelong friends—Bohr was six years the younger and died in 1962, seven years after Einstein.

In two important respects, the physicists were "extreme opposites." To Bohr, "one and only one place was home: Denmark. Einstein...would call himself a gypsy, or a bird of passage"; he lived, not as visitor only but as resident, in nine cities of five countries. Most striking was "Einstein's apartness, Bohr's conjointness.... Einstein was not a lone-

ly figure.... Nevertheless it was his deepest need to think separately.... Bohr...craved togetherness, in life and in thought. Bohr created a major school; Einstein did not." There were profound similarities, too. "Both had a deep need for simplicity, in thought and in behavior. Each had a lifelong boyish—not juvenile, boyish—curiosity, and pleasure in play. They took science very seriously, but to them it was ultimately a game."

So out of old memories and long careful inquiry writes Professor Pais, himself a distinguished contributor to postwar theoretical physics. Pais has now published ample biographies of "arguably the two leading figures in science in this century." His two thick, vivid books—in 1982 *Subtle Is the Lord...*, on Einstein, and in 1991 the book on Bohr—do justice to career, character and ideas, not always the case in chronicles of modern scientists for the non-specialized reader. Here we tease only a few strands out of the long tapestry of Bohr's century and our own.

Young Niels entered the University of Copenhagen in 1903 as a physics ma-

jor. The next year his brother, Harald, a year and a half younger, joined him on campus. The two were inseparable companions. "I have never known people to be as close as they are," a fellow student wrote. They were mutual confidants, linked by teasing and by private games, "thinking *à deux*" and sharing delight in science, poetry, shopwork and the soccer field. Halfback Harald became a national figure in the sport; slower, stronger Niels played goalie, sometimes visibly inattentive. Harald went on to win an international reputation in mathematics; he died at 63. Niels later said of him: "He was in all respects more clever than I...a great mathematician, you know." The boys' father was a physiology professor, later to become University rector, widely judged wise and friendly; their mother was a loving, energetic woman from a wealthy Jewish family admired in tolerant Denmark for its public spirit. The family was not at all religious; the boys were baptized Lutherans who left the church in maturity.

In 1911 Niels went as a postdoc to



NIELS BOHR and his wife, Margrethe (at left), were photographed at the Copenhagen airport in August of 1945, the day they returned to Denmark from an exile that had been



imposed by the war. At right, lifelong friends Bohr and Albert Einstein stroll down a street in Brussels during the 1930 Solvay Conference.

J. J. Thomson at the Cavendish Laboratory, a brief and disappointing encounter. He soon joined Ernest Rutherford at Manchester, to open there a fulfilling, lifelong friendship. Home he came after a year or so with his stunningly original idea for quantized electron orbits around Rutherford's new nucleus. In its unreasonably successful "triumph over logic," the quantum theory that Bohr published in 1913 still defines the public icon of the atom.

Within a few years, Bohr was well known, became Professor at Copenhagen and was ready to seek and indeed soon to receive for his own the small, new Institute for Theoretical Physics, which had both public and government support. That beloved home for quantum theorists has borne its founder's name since 1965. Bohr's passion, integrity and insight nourished dozens of younger physicists who came to the institute. Besides his tireless determination to understand "conjointly," he never failed to provide the practical essentials, support and a place to work among the best of peers, the celebrated names that ornament the major findings of quantum mechanics—among them Heisenberg, Schrödinger, Dirac, Pauli, Kramers, Klein, Landau, Gamow, Franck. About 1,200 papers came from the institute during Bohr's time; more than 400 visitors from 35 countries had worked there.

Bohr raised enough Danish and international support (much from American foundations) to make his institute a training ground and a refuge. His palatial Residence of Honor, a lifetime grant from the Carlsberg Foundation, he made into a family hearth, its spirit strong enough to sustain intense argument, unmarred by arrogance or conceit. "He utters his opinions," Einstein wrote, "like one perpetually groping and never like one who believes he is in possession of definite truth."

Philosophy, though not in the sense of the professional literature, was what Bohr practiced, much more than most of his pragmatic friends. In a way what he did begins with form, not content; his papers were usually less mathematical and technical than most by a great deal. (That does not make them easy: they are intricate, but their language, always qualified and subtle, not to say tentative, is still the natural language we all share.) His best-known single idea is complementarity. He devised a coat-of-arms when he was given the Danish Order of the Elephant: it displays the entwined Chinese forms of the monad, opposite and yet completing each other. "Opposites are complementary," it reads. The poet William



If you're a Jack Daniel's drinker, drop us a line. We enjoy hearing from our good friends.

AN EMPTY ROCKER in Jack Daniel's Hollow reminds us we're making a bit more Jack Daniel's these days.

Call for our Tennessee Whiskey is higher than ever right now, so we've had to forego some sitting time. Mind you, we're not making Jack Daniel's faster. The old slow way we've always used is the way we know you like. But be assured we are working longer to supply your needs—no matter how much we miss our rockers.

SMOOTH SIPPIN'
TENNESSEE WHISKEY



Tennessee Whiskey • 40-43% alcohol by volume (80-86 proof) • Distilled and Bottled by Jack Daniel Distillery, Lem Motlow, Proprietor, Route 1, Lynchburg (Pop 361), Tennessee 37352
Placed in the National Register of Historic Places by the United States Government.

ChiWriter

The definite tool.

Now – Version 4.1

Full-featured scientific word processing

ChiWriter gives you all the features of an advanced scientific word processor and more! The WYSIWYG screen display lets you enter and edit text and scientific notation exactly as you want it on your printout. Add easy math editing, font design, multiple undo and redo, automatic formatting, import graphics, mouse support, macros, foreign language capability, spell checker, and support for dot-matrix and laser printers — *all for only \$349.95* plus shipping and handling.*


Call today to order your **ChiWriter** package. Full 30-day money back guarantee. Bankcards welcome.

* Chemistry Fonts \$59.95,
Russian Fonts \$29.95

20% educational discount.

To order, call toll free

1 • 800 • 736 • 8886



FEATURES

- Screen display corresponds exactly to printout — no encoding or preview mode
- Easy formula editing. Symbols and formulas can be anywhere in text
- Multiple undo and redo — easy to fix mistakes, or change your mind
- Automatic formatting of text, equations and tables
- Simultaneous editing of up to 10 documents
- Graphics import — PCX and TIFF files
- Font designer — modify symbols or design your own
- Foreign language capability
- Mouse support and menu driven for ease of use
- Unlimited levels of superscripts and subscripts
- All standard math and scientific symbols included
- Spell checker included
- DOS shell



HORSTMANN SOFTWARE DESIGN CORPORATION

4 N. 2nd St., Ste. 500 • P.O. Box 1807 • San Jose, CA 95109-1807, USA

Phone: 408 • 298 • 0828 FAX: 408 • 298 • 6157

IBM is a registered trademark of International Business Machines.

Sometimes the PREDICTABLE THINGS IN LIFE are the most unexpectedly REASSURING.



A DEPENDABLE, RELIABLE source of monthly income.  If that sounds like something you'd find predictably reassuring, ask your financial adviser about the benefits of a Nuveen Tax-Free* Unit Trust.  And rest assured, the return on your Nuveen investment will remain the same until bonds in the portfolio mature, are called or are sold.[†]  But please, request a prospectus with more complete information (including charges and expenses), and read it carefully before you invest or send money. Or call us toll-free at...

1-800-262-3423.

NUVEEN

Quality Tax-Free Investments Since 1898.

*Income may be subject to state and local taxes. Save additional taxes with Nuveen state trusts which are also exempt from state and local income taxes. †Unit trusts have a fluctuating redemption value

Blake put the same idea thus: "Negations exist not, but contraries mutually exist." Thus wave and particle, thus uncertainties of place and motion, and much more. But the concept gives no general solution to real problems; it is to be used sparingly as a case-by-case hint to the direction of specific and hard-won progress.

Pais is honestly impatient with the present revival of interest in the controversy about the local nature of the properties of atomic systems. The matter was first raised by Einstein, Podolsky and Rosen in 1935 and answered by Bohr: no change needed, he said. It hangs on the strange fact—Einstein called it "spooky action at a distance"—that certain arrangements allow the experimenter to delay a measurement that can fix at option either member of a complementary pair of unknowns. Yet you cannot learn both results, even though it often appears that the particle in question has long before had its properties settled by a prior interaction. The remarkable experiments of today all confirm the Bohr view. Pais sums up by saying that Bohr simply redefined the concept of phenomenon: it must include "both the object of study and the mode of observation...including an account of the whole experiment," all the way through.

An illustration here in what Bohr called the pseudorealistic style is a start toward understanding. A slit position is well defined not by symbols but by heavy brass bolts that secure it to some stable optical bench. If you measure momentum by collision, you will instead hang the test mass almost freely on a delicate fiber and surrender to wobble all precise information about just where it is. That is complementarity expressed in hardware. The modern critics will hardly be satisfied with Pais's judgment, but his case is strong if summary and supported by every experiment so far.

Peer into one of the last pages of Bohr's physics: nuclear fission. During the 1930s, Bohr had built the institute in Copenhagen into an important center for nuclear experiment. Otto Robert Frisch and Lise Meitner, working with Bohr in 1938, gave a physical explanation of the astonishing radiochemical demonstration that the uranium nucleus could be split nearly in half by slow neutrons. Frisch was the first to see the energetic fragments of the fissioned nuclei. Recognition of the unstable balance between the repulsive electric forces among protons and the attractive bonds that hold neutrons close to their proton partners owed a good deal to Bohr's 1936 modeling of all heavy nuclei as

drops of a charged quantum fluid. That persuasive approach in turn bears an uncanny parallel to the essay that undergraduate Niels had written in 1906. Using the resources of his father's physiology lab, he had measured photographically the surface vibrations that break up a fine jet of water into droplets. His essay was awarded the gold medal of the Danish Academy; "a lovely experiment," it was the first and the last experiment Bohr ever performed.

Bohr brought the burning-hot news of fission to America when he arrived on a long-planned visit to Princeton in 1939. By early February of that year he published a compelling argument to show that the unstable uranium could be only a rather rare isotope of that element. That nucleus was both less stable against fission and energetically more excited on neutron capture than the much more abundant one. By September he and American theorist John Archibald Wheeler, an old boy of the Copenhagen institute, offered a pretty full theory of the fission process. In early 1940, experiment confirmed the prediction; from then onward it was uranium 235 that was the superexplosive sought around the warring world.

In 1943 Niels and Margrethe Bohr (and their sons) fled Denmark one night by sea. They were at mortal risk, among 7,000 Danish Jews and refugees whom the Danes spirited to neutral Sweden only days before the planned Nazi roundup for deportation. No other occupied country so successfully saved its own from the camps of the Third Reich.

Internationalist Bohr, from a small and traditionally neutral land, who had himself first clarified fission, fled before the Germany he knew was somehow engaged in the release of nuclear energy, all the way to wartime Los Alamos. Fission, though unexpected, is no profound puzzle for physics. It is for polity that it is of decisive importance; the nuclear weapons it alone enables hang heavy over the last half of the century. Naturally Bohr was seized with this knot of knots. He had the start of a solution, one he pressed with no success on our wartime leaders. He made it public in 1950 and again in 1956, in long letters to the United Nations. It seemed utopian to many and never caught the attention of the world, prepossessed by power, but "fell through the cracks in the ice of the cold war." And yet his idea is in a way regnant.

For Bohr was the "pioneer of glasnost." In 1948 his memo to Secretary of State George Marshall proposed "an offer...of immediate measures towards openness on a mutual basis" to allow all to form judgments about the real

YOU'RE SERIOUS ABOUT TREATING BLEEDING GUMS.



SO ARE WE.

Viadent® Toothpaste and Oral Rinse is the only dental regimen with sanguinaria. This combination is clinically proven to reduce bleeding associated with gingivitis, an early reversible gum disease. So, if your dental professional tells you to take better care of your gums, listen. And start an oral care program including Viadent, flossing, and professional care.

Viadent treatment. Not just toothpaste.

© 1992 Viadent, Inc.

THE HUNTINGTON BRONZE COLLECTION

Fine art bronze sculpture



Bronco Buster by Frederic Remington

This is one of over 300 "lost wax" bronze sculptures in our collection. Call for our full color catalog today! 1-800-777-8126. All sculptures are guaranteed and made in the USA. Quality, personalized service.



401 E. Cypress Ave. Visalia, CA 93277
(209) 732-8126 • Fax (209) 732-5961
MC/VISA

Science Connection

Science Connection is a North America-wide singles network for science professionals and others interested in science or natural history.

Begun in spring 1991, the network operates by facilitating contacts among members, not by "matchmaking". Members are from all parts of the United States and Canada (every state and 8 provinces) and range in age from twenties to late seventies.

For further information, please contact us by mail or phone.

Science Connection Inc.

PO Box 188, Youngstown, NY 14174
(800) 667-5179 (U.S. only)
(519) 583-2858



state of affairs. Nobody was ready, until, after a long time of change, there came Mikhail Sergeyevitch Gorbachev. It would be ingenuous to suppose that the mere notion from an outsider could have meant much early on, but it is the case that thereabouts was found a hopeful path that seems to be leading us toward the edge of the grim forest. No better memorial could we raise to Niels Bohr.

Was It Murder?

MAMMOTHS, MASTODONTS, AND ELEPHANTS: BIOLOGY, BEHAVIOR, AND THE FOSSIL RECORD, by Gary Haynes. Cambridge University Press, 1991 (\$69.50).

The scholarly title obscures the real topic. In fact, this volume is a brilliant murder mystery. The murder indictment itself is nothing new. The defendants are the first Americans, whose presence on our continent is signaled by ample evidence, those bold and skillful immigrant Paleo-Indian hunters we call the Clovis culture (after a defining archaeological site). The prosecution argues that they killed off within a few centuries all the wild elephants then abundant in North America. Their co-defendants are the redoubtable mammoth hunters of northern Eurasia, charged with extinguishing the great woolly beasts there at about the same time.

The extinction of temperate-region elephantkind is plain enough. Today only two near-tropical species remain: the African elephants, whose broad ears resemble the map of Africa, and the Asian species, whose smaller ears evoke the map of India. After millions of years of elephantine lineage, the end for most species came within a few millennia; in the Americas at least, it followed hard on the entry of the hunters. Direct evidence is grim: the dated and identified bones of mammoth and mastodont are heaped up in killing and butchering sites, at dozens of locations in North America and more than 1,000 across Eurasia. At more than a dozen American sites, several of the distinctive fluted dart points fashioned by the Clovis people have been found with the rib cages of fossil mammoth and mastodont. The prosecutors do not hesitate to cite many another instance of human overkill of big game. We cannot ignore the precipitate decline of the African elephant population during our own century, although a circumspect judge might rule out such guilt by association.

Enter the author, an energetic anthropologist at the University of Nevada.

His brief is too scrupulous to be called advocacy; put him down as friend of the court. Neither rough coincidence in time nor plausible generalizations can justify a firm conclusion for him. Forensics must be called on to test the chief inspector's keen inferences. Wild elephants still die daily. What can we learn from modern remains to assess conclusions about relics a hundred centuries old?

Every year since 1982 Haynes has sought out elephant dead in the wildest quarter of the Zimbabwean national park named Hwange, on the Botswana border. Death by human poachers has been effectively eliminated by the park management, although they themselves have systematically culled the herd to control the population in Hwange, an area twice the size of Yellowstone. Elephant bones accumulate around water holes, although the bones weather away or become buried after some decades. The driest months find elephants dying within the network of meandering shallow sandy channels, called seeps, that are watercourses during the wet season there at the edge of the Kalahari. The elephants come to dig and redig wells, seeking scarce water down a meter or two in the sands. One of five or so die; the younger, smaller ones find the water table sunken out of reach and die preferentially. A map of the largest of the die-off sites the author watched shows the places of death over nine years of 100-odd elephants along a winding seep, all within a couple of hundred acres.

Haynes observed dozens of mass kills, too; sometimes herds of 30 to 50 elephants were spotted by aircraft, then stalked downwind, to die within a minute before the rifles of a few marksmen. The kills are analyzed for demographic information about age classes, and then local men butcher the carcasses. Only a few large males accompany each typical matriarchal herd. These herd kills center in the uplands among abundant browse, on much-used elephant trails, just as die-off clusters are found at lowland seeps in the sandy scrubland.

For the archaeologist, the riveting experience was "disturbing...but it was instructive." The epigraph of this chapter cites a rueful old professional hunter: "I should not greatly care to kill any more elephants. They are too big, too old, and too wise to be classed as mere game." Yet the Pygmy metaphor for their rare elephant prey is "a hill of meat," and Clovis hunters may have shared the attitude.


From these witnessed deaths among elephants, one can begin to infer events long past. Zimbabwe is not glacier-edge

New Mexico, and the African elephant is no woolly mammoth. The author has a flair for expressing open doubt as well as for dispelling it. He carefully shows that his models, approximate and never certain, offer a simple and strong interpretation of the record we find in worked points, old broken bones, glacial till and soil. Some Clovis sites repeat the age pattern of the Hwange die-offs: subadults predominant, and very few of their elders, especially in those sites where stone artifacts are present. Some sites are patternless, the data too sparse; none of them shows fair age samples of elephant population. "It is deceptively easy to find a... relationship between... human presence and an overrepresentation of young animals." But at the dry seeps during Zimbabwe winters, it became manifest that such an inference is not necessary.

The huge piles of mammoth tusks that mark the most spectacular of the Eurasian sites seem to tell us that those old hunters dwelled in huts walled and floored in ivory. But did they in fact hunt the dangerous elephants for that exotic architectural material? Or did they mine the stuff from the abundant remains of mass deaths of mammoths and make dwellings from the bones, perhaps when wood was in short supply? After all, some 50,000 mammoth tusks have been collected across the land within the past century or so. The more romantic image fades, even though it is not strictly disproved.

But what of the extinction of the American mammoths by Clovis hunters? The multiple kill sites just don't look like the ethnographic record. "I believe," the author concludes, that the large Clovis mammoth assemblages are the remains of die-offs exploited by the people who made fluted points. Elephant die-offs everywhere provided opportunities for the newcomers, who were indeed skilled hunters, come to easy pickings. The rapid spread of the Clovis culture over the hemisphere was not the cause of mammoth extinction, but rather more the effect of it.

One cannot reject the possibility that the hunters added the last straw to a heavy climatic burden. Perhaps they were the unique new agency behind sudden extinction, critically increasing the natural mortality of the hard-pressed herds. The verdict on Clovis may have to be moderated to something like murder in the second degree. The free-roaming African elephants who now face their last few decades—we will probably manage to preserve the species, though only under close watch—can show us how the end came to the societies of the elephant long ago.



Computers & Weather: Together at Last!

Finally, you can bring the weather home! Combine our Weather Monitor II with our Weatherlink Software for the most complete, state-of-the-art weather station available. Use the power of your PC to record, summarize, and graph the weather—from today, yesterday, or a year ago!

WEATHER MONITOR II FEATURES:

- Inside & Outside Temps
- Wind Speed & Direction
- Inside Humidity
- Time & Date
- Barometer
- Wind Chill
- Alarms
- Outside Humidity & Dew Point Option
- Highs & Lows
- Instant Metric Conversion
- Rainfall Option

WEATHERLINK FEATURES:

- Create daily, weekly, monthly, or yearly plots
- Compare two weather conditions on the same graph
- Compare two days, weeks, months, or years on the same graph
- Store information for up to 4 months before transferring to the PC
- Monitor conditions in remote locations using a modem



The Ultimate In Weather Monitoring Systems!

Call Today: 1-800-678-3669 • M-F 7 a.m. to 5:30 p.m. Pacific Time • FAX 1-510-670-0589

One Year Warranty • 30-day money-back guarantee • M/C and VISA • SC321L

Weatherlink system requirements: IBM compatible, 640K RAM, Hercules, CGA, EGA, or VGA

Weather Monitor II also sold separately • Weatherlink also works with Davis' Perception II and Weather Wizard II

DAVIS INSTRUMENTS 3465 DIABLO AVE., HAYWARD, CA 94545

How Air Support Gives You A Better Night's Sleep. Guaranteed.



Select Comfort's unique air cushion design conforms perfectly to your body's contours and provides total body support.

Adjust firmness to your exact needs

Select Comfort is a revolutionary air sleep system that allows you to choose your comfort and support level simply by pushing a button. In seconds, you can set it extra firm or as soft as a feather bed. And because there are dual air chambers, your partner can do the same.

Select Comfort provides superior support.


Select Comfort's unique air-cushion design conforms perfectly to the contours of your body. It provides total body comfort and total body support, regardless of your weight or shape. And because Select Comfort provides excellent spinal support, it helps people with back pain sleep better.

90 Night Trial Program

We're so confident you'll sleep better on a Select Comfort Sleep System that we offer a 90 Night Trial Program. We guarantee you'll sleep great. If you don't, we'll gladly refund your money.

"What a breakthrough in sleeping! I no longer toss and turn to find a comfortable sleep position."

—R. Johnson, Valley City, ND



To find out how you can get a better night's sleep, call for a

FREE VIDEO AND BROCHURE
1-800-831-1211

☐ **YES!** Please send me a FREE information packet and videotape.

Name _____

Address _____

City _____ State _____ Zip _____

Evening Phone () _____

Select Comfort

9301 75th Avenue North, Brooklyn Park, MN 55428
1-800-831-1211

©1992 Select Comfort

Dept. 621



Trouble in Mind

Einstein once said that the most incomprehensible fact about nature was that it was comprehensible. It seems to me that the mysterious thing about nature is not that it is comprehensible but that it contains such a thing as comprehension at all, that is, the mind itself—the very idea!

Of course, ever since Darwin we have been able to consider the mind as something emerging from nature by virtue of certain arrangements developed through chance and necessity. Although Darwin is rightly credited with the discovery of the biological process that yielded such developments, their nature remained a mystery until the middle of this century. Thus, when early 19th-century physiologists brazenly invoked “material organization” as an explanation for both Life and Mind, it was still little more than hand-waving.

Today all that has changed. In the once unoccupied gap between the bare necessities of Matter and the enigmatic peculiarities of Mind, there is now an elaborate construction site of mediating concepts whose existence lends weight to the claim that the functions of mind are implemented by purely physical means. Neurons and their synapses *do* have the capacity to transact what is required to realize seeing and foreseeing. Conversely, at the other end of the psychophysical gap the image of mind itself has been radically redesigned. Under the influence of cognitive science, information theory and artificial intelligence, we can now envisage mental functions in a form that literally demands physical implementation.

The theoretical structures that reach toward one another across the great divide are developing so rapidly it is tempting to imagine that it's only a matter of time before they join up and that when that happens the terms once applied to mental life will be supplanted by a different sort of language altogether. In some quarters, this belief is already upheld as an article of faith; although paradoxically, among those who subscribe to it, the very concept of belief is regarded as questionable, along with many of the other psychological categories that figure in common parlance. Hence, the disparaging phrase “folk psychology.” According to its critics, folk psychology is a hopelessly sloppy theory of human conduct, featuring terms such as “belief” and “desire,”

which are said to be not all that different from “phlogiston” and “caloric.” Philosopher Paul M. Churchland insists that “we need therefore an entirely new kinematics and dynamics with which to understand human cognitive activity, one drawn perhaps from computational neuroscience and connectionist A.I. Folk Psychology could then be put aside in favor of descriptively more accurate...portrayals of the reality within.”

No one can deny that the sciences to which Churchland refers will furnish more inclusive accounts than the ones we have at the moment. But that doesn't mean that folk psychology is nothing other than interim discipline. Nor is it a jury-rigged theory, good enough for ordinary folk and acceptable to experts as long as they are off duty. It is not and never was a theory, and to speak of it, as Churchland does, as an explanation “which fails on an epic scale” is to make a serious category mistake. The vernacular language of beliefs and desires is the expression of a form of life peculiar to creatures whose consciousness leads them to believe—yes, believe!—that others have experiences of the same kind.

In fact, it is hard to imagine how a radically different language would even begin to do a better job. It could and has been made more expressive, as any student of literature knows, and because of imaginative writers such as Shakespeare, say, or Proust, we can now convey psychological subtleties that were once inexpressible. A hard-core eliminativist would probably jeer that such revisions are doomed to failure and, like the epicycles added to Ptolemaic theory, are futile attempts to salvage a derelict system.

That, I venture to say, is nonsense. Consciousness may be implemented by neurobiological processes—how else?—but the language of neurobiology does not and cannot convey what it's *like* to be conscious. If, as philosopher Thomas Nagel says, there is something it's *like* to be a bat, there is something even more interesting it's like to be one of us, and the language of folk psychology, pidgin though it be in some respects, is the best medium for expressing this.

And that, of course, brings me to consciousness, which is quite rightly regarded as the most difficult problem in nature, more puzzling perhaps than the one which worried Einstein. And yet there are scientists prepared to insist that consciousness itself will yield to analysis, just as the problem of life has yielded. But there's the rub!—or rather the double bind. Although conscious-

ness exists by virtue of some physical property of the brain, just as bioluminescence exists by virtue of some chemical property of certain specialized cells, it is *not*, as bioluminescence is, an observable property of living matter. It isn't a brain glow. Nor is it, on the other hand, an *invisible* property, less readily detectable than other biological processes. It is detectable to anyone who has it. The difficulty is that the method by which consciousness is detected is logically different from the method by which bioluminescence is detected. To put it bluntly, consciousness is not *detected* at all, because that would imply that it could pass *undetected*, and that doesn't make sense. *Your* consciousness may pass undetected by me, but *my* consciousness, if I have it at all, is self-evidently self-evident to me.

For that reason, we invoke a special kind of observation to which we can give the name “introspection.” And yet introspection is not, as the word seems to imply, a peculiar form of gaze, directed at something inaccessible to the more typical forms of looking. On the contrary, introspection is one of the many forms consciousness itself can assume, so that it represents a significant part of what we are trying to explain. Indeed, the method by which we are acquainted with consciousness is so fundamentally different from the method by which we acquaint ourselves with brains that I suspect, as philosopher Colin McGinn does, that although we don't have to invoke anything other than brain—no magic that contravenes the laws of nature—we will never fully understand the connection.

This claim has been widely dismissed as frivolous obscurantism, foreclosing the possibility of further research and licensing the wilder forms of religious mysticism. It is nothing of the sort. There is obviously much more to be learned about the relationship between brains and minds, and it will be years, perhaps centuries, before we come up against the “cognitive closure” so courageously identified by Professor McGinn. The fact that such research is destined to describe an asymptotic curve, which approaches but never reaches the limit, does not preclude the necessity of our following it.

JONATHAN MILLER, physician, writer and director, produced the 1983 public television series “States of Mind” and the 1992 series “Madness.”

Introducing the New Grolier Multimedia Encyclopedia™ with exciting features for 1992

*We are filled with awe as we
explore the solar system.
Each discovery comes
alive with spectacular
sights and sounds.
It's getting late, but
we're not ready to
head back
to Earth...*



While exploring the final frontier, we met
Copernicus, saw the Space Shuttle and listened as
Neil Armstrong took a giant leap with one small step.



The Grolier
*Electronic
Encyclopedia* has
long been recog-
nized as the pre-
mier CD-ROM
encyclopedia for

serious, effective research. Now,
Grolier introduces a remarkable
resource that sets a whole new
standard for multimedia encyclo-
pedias—the *New Grolier
Multimedia Encyclopedia*™

**Serious research.
Serious fun.**

Featuring all 21 volumes of the
*Academic American Encyclo-
pedia* on a single CD-ROM,
the *New Grolier Multimedia
Encyclopedia* provides a refer-
ence resource unparalleled
in both quality and
comprehensiveness.

Within seconds,
you can locate articles on
everything from cov-
ered wagons to lunar

landers. And a host of exciting
new features makes the urge to
explore virtually irresistible. You'll
find more color photographs, illus-
trations, sounds and motion
sequences. And timely

coverage of such recent
world events as the
Persian Gulf War and
dissolution of the
Soviet Union.

But that's just the
start. You can actually
listen to famous
speeches, music and
more. And explore
every corner of the
world using over 250
high-quality maps.

The new edition
also features a unique
Timeline with over
5,000 entries that lets
you journey from

prehistory to the present, linking
you directly to articles. And our
new Knowledge Tree takes the
search out of research by letting



you explore broad topics, then
quickly narrow your search to
specific topics.

Still just \$395.

It's been selected Product of the
Year by the Optical Publishing
Association. And it's *still* just
\$395. For more information on
the *New Grolier Multimedia
Encyclopedia*, call toll-free today.


1-800-356-5590



The New
**GROLIER MULTIMEDIA
ENCYCLOPEDIA**

The New Grolier Multimedia Encyclopedia
runs on today's most popular platforms includ-
ing MS-DOS, Macintosh and Windows/MPC.

Grolier Electronic Publishing, Inc., Sherman Tnpl., Danbury CT 06816 □ 203-797-3500



It works for beginners as well as experts.

And no matter how hard you try, you can't exhaust its potential.

In many ways, it's a lot like CompuServe.

For the beginner at interactive computer services, we've got news, travel information, stock quotes, entertainment and games, a complete encyclopedia, free time to sharpen your online skills, and more.

For the experienced user, we feature hardware and software support, special-interest forums, free software, a wealth of online computer expertise, and sophisticated research tools. In fact, no computer service offers the choices that CompuServe does.

Now, for just \$7.95 a month, and a one-time membership fee, you get all the basics as often as you like: news, sports, weather, shopping, a complete encyclopedia, and much more, plus up to 60 E-mail messages a month. And, there are lots of other valuable services available on a nominal pay-as-you-use basis.

Whether you're a beginner or an expert, to choose the right interactive service, just use your brain. Pick CompuServe.

For more information or to order, see your computer dealer or call 1 800 848-8199. Outside the United States, call 614 457-0802.

CompuServe®

The information service you won't outgrow.