SCIENTIFIC AMERICAN
AUGUST 2010
$5.99
WWW.SCIENTIFICAMERICAN.COM
CELEBRATING 165 YEARS
Secrets of Our Success
Once humans almost went extinct.
Here's the untold story of our salvation
"SUPER-EARTHS" Could They Harbor Life?
ROBOT PILLS Doctor in a Capsule
HARDWARE HACKERS The Next Security Threat
PLUS
ORIGINS SPECIAL
MORALITY
ANIMATION
INFLUENZA
FIREWORKS
SWISS CHEESE
BARBED WIRE
HAND WASHING
FAMILIES
AND MORE
© 2010 Scientific American

www.dow.com/hu
®™The DOW Diamond Logo and Human Element and design
are trademarks of The Dow Chemical Company © 2010

I am the Human Element in the relentless pursuit of making things faster and smaller. Electronic materials, chips and semiconductor technology, these are the things I live and breathe every day. Innovation in this world happens at the molecular level. And success is in the power of small.

~Dr. George Barclay
Dow Electronic Materials



SPACE

38 **Planets We Could Call Home**
By Dimitar D. Sasselov and Diana Valencia
A startling number of planets in our galaxy could be similar to Earth—and may even support life.

SPECIAL REPORT

46 **Origins**
How did sex get started? Who invented barbed wire? When did zero become something? These and other beginnings, explained.

54 **When the Sea Saved Humanity**
By Curtis W. Marean
A harsh change in climate might have wiped out *Homo sapiens* long ago if a lucky band of hunter-gatherers hadn't taken shelter in an idyllic corner of Africa.

MEDICINE

62 **Robot Pills**
By Paolo Dario and Arianna Menciassi
Tiny devices may soon voyage through the digestive tract, performing surgery and diagnosing disease.

ENVIRONMENT

66 **Threatening Ocean Life from the Inside Out**
By Marah J. Hardt and Carl Safina
Carbon emissions are making the oceans more acidic, imperiling many species from plankton to squid.

IMAGING

74 **Filming the Invisible in 4-D**
By Ahmed H. Zewail
An amazing form of microscopy can make movies of vanishingly small, nanoscale objects in action.

TECHNOLOGY

82 **The Hacker in Your Hardware**
By John Villasenor
Microchips in many common electronic devices might already be compromised, with dire consequences.

OBJECT LESSON

88 **Plastic Surf**
By Jennifer Ackerman
Photograph by Cary Wolinsky
The unhealthful afterlife of toys and packaging.

ON THE COVER

During a long-ago climate catastrophe a small group of *Homo sapiens* found refuge in a bountiful seashore near Africa's tip. We are all the beneficiaries. IMAGE BY JEAN-FRANCOIS PODEVIN.

PER-ANDERS PETTERSON (*cover background*); FRANK HÜLSBÖMER (*robot pill*)

© 2010 Scientific American

The Norwegian Academy of Science and Letters
announces the

2010 WINNERS OF

THE KAVLI PRIZE℠

Eight scientists whose discoveries have dramatically expanded human understanding in the fields of astrophysics, nanoscience and neuroscience have been recognized with the award of the million-dollar Kavli Prizes.

The laureates were chosen for research that has transformed our knowledge of basic units of matter, laid the foundations for the field of nanotechnology, revealed the molecular basis for the transfer of brain signals and other physiological functions, and made possible the building of telescopes that can see deeper into space and further back in time.

The Kavli Prize is a partnership between The Norwegian Academy of Science and Letters, The Kavli Foundation℠ (US) and The Norwegian Ministry of Education and Research.

HM King Harald presents the Kavli Prize to the 2010 Laureates on September 7, 2010 at a ceremony in Oslo.

THE KAVLI PRIZE IN ASTROPHYSICS

JERRY E. NELSON
Professor,
Lick Observatory,
UC Santa Cruz, US

RAYMOND N. WILSON
Senior Physicist,
European Southern Observatory,
Germany

JAMES ROGER PRIOR ANGEL
Professor,
Steward Observatory,
University of Arizona, US

"for their contributions to the development of giant telescopes"

THE KAVLI PRIZE IN NANOSCIENCE

DONALD M. EIGLER
IBM Fellow,
IBM Almaden Research Center, US

NADRIAN C. SEEMAN
Professor,
New York University, US

"for their development of unprecedented methods to control matter on the nanoscale"

THE KAVLI PRIZE IN NEUROSCIENCE

RICHARD H. SCHELLER
Executive Vice President,
Genentech Research & Early
Development, US

THOMAS C. SÜDHOF
Professor,
Stanford University, US

JAMES E. ROTHMAN
Professor,
Yale University, US

"for discovering the molecular basis of neurotransmitter release"

To learn more about The Kavli Prize℠ please visit
www.kavliprize.no
www.kavlifoundation.org

CONTENTS DEPARTMENTS

6 **From the Editor**

8 **Letters**

12 **50, 100 & 150 Years Ago**

14 **News Scan**

ENERGY & ENVIRONMENT
- Gulf oil: we wanted it—now how do we get rid of it?

RESEARCH & DISCOVERY
- What's next for cells powered by synthetic genomes?
- Lab accidents haunt experimental science.
- Rapid eye movements track targets in dreams.

MEDICINE & HEALTH
- Lamarckian phenomena and heritable diseases.
- Dubious claims still pervade supplements industry.

TECHNOLOGY
- Crash-testing batteries for electric-car safety.
- Instead of telecommuting, try a surrogate robot.

30 **Perspectives**
By the Editors
The way to avoid ruinous oil spills is to fix our national energy policy.

32 **Sustainable Developments**
By Jeffrey D. Sachs
Economic, political and cultural forces have boosted Poland but failed Russia.

34 **Skeptic**
By Michael Shermer
Why Neandertals were not a separate species.



COURTESY OF ANYBOTS

36 **Critical Mass**
By Lawrence M. Krauss
Religious leaders should be held accountable when their irrational ideas turn harmful.

90 **Recommended**
Arctic change. Malaria's grip. Periodic table.

92 **Anti Gravity**
By Steve Mirsky
Physics and medicine are big players on the baseball diamond.

GO TO SCIENTIFICAMERICAN.COM

Meeting of the Minds in Lindau

The 2010 Lindau Nobel Laureate Meetings in Germany brought together dozens of past Nobel Prize winners and hundreds of young scientists to bridge disciplinary, national and—perhaps most important—generational divides.

MEETINGS OF NOBEL LAUREATES LINDAU | EDUCATE. INSPIRE. CONNECT.
www.lindau-nobel.org

More at **www.ScientificAmerican.com/aug2010**

COURTESY OF THE LINDAU NOBEL LAUREATE MEETINGS; COURTESY OF ANYBOTS (*robot*)

Scientific American (ISSN 0036-8733), Volume 303, Number 2, August 2010, published monthly by Scientific American, a trading name of Nature America, Inc., 75 Varick Street, 9th Floor, New York, N.Y. 10013-1917. Periodicals postage paid at New York, N.Y., and at additional mailing offices. Canada Post International Publications Mail (Canadian Distribution) Sales Agreement No. 40012504. Canadian BN No. 127387652RT; QST No. Q1015332537. Publication Mail Agreement #40012504. Return undeliverable mail to Scientific American, P.O. Box 819, Stn Main, Markham, ON L3P 8A2. **Individual Subscription rates:** 1 year $39.97 (USD), Canada $49.97 (USD), International $61 (USD). **Institutional Subscription rates:** Schools and Public Libraries: 1 year $69 (USD), Canada $74 (USD), International $81 (USD). Businesses and Colleges/Universities: 1 year $299 (USD), Canada $304 (USD), International $311 (USD). POSTMASTER: Send address changes to Scientific American, Box 3187, Harlan, Iowa 51537. **Reprints available:** write Reprint Department, Scientific American, 75 Varick Street, 9th Floor, New York, N.Y. 10013-1917; fax: 646-563-7138; reprints@SciAm.com. **Subscription inquiries: U.S. and Canada (800) 333-1199; other (515) 248-7684. Send e-mail to sacust@sciam.com.** Printed in U.S.A.

Copyright © 2010 by Scientific American, a division of Nature America, Inc. All rights reserved.

© 2010 Scientific American

It's your vacation, why fly on anyone's schedule but your own?

Expedia lets you fly there on one airline and back on another. So, you're never at the mercy of one airline's schedule. Which means more flight options, and more time where you want to be.

expedia.com 1-800-EXPEDIA

©2010 Expedia, Inc. All rights reserved. Expedia, Expedia.com and the Airplane logo are either registered trademarks or trademarks of Expedia, Inc. in the U.S., Canada and/or other countries. CST 2029030-50

BOARD OF ADVISERS

LESLIE C. AIELLO
President, Wenner-Gren Foundation for Anthropological Research

ROGER BINGHAM
Professor, Center for Brain and Cognition, University of California, San Diego

G. STEVEN BURRILL
CEO, Burrill & Company

ARTHUR CAPLAN
Emanuel and Robert Hart Professor of Bioethics, University of Pennsylvania

GEORGE M. CHURCH
Director, Center for Computational Genetics, Harvard Medical School

RITA COLWELL
Distinguished Professor, University of Maryland College Park and Johns Hopkins Bloomberg School of Public Health

DREW ENDY
Professor of Bioengineering, Stanford University

ED FELTEN
Director, Center for Information Technology Policy, Princeton University

MICHAEL S. GAZZANIGA
Director, Sage Center for the Study of Mind, University of California, Santa Barbara

DAVID GROSS
Frederick W. Gluck Professor of Theoretical Physics, University of California, Santa Barbara (Nobel Prize in Physics, 2004)

LENE VESTERGAARD HAU
Mallinckrodt Professor of Physics and of Applied Physics, Harvard University

DANNY HILLIS
Co-chairman, Applied Minds

DANIEL M. KAMMEN
Director, Renewable and Appropriate Energy Laboratory, University of California, Berkeley

VINOD KHOSLA
Founder, Khosla Ventures

CHRISTOF KOCH
Lois and Victor Troendle Professor of Cognitive and Behavioral Biology, Caltech

LAWRENCE M. KRAUSS
Director, Origins Initiative, Arizona State University

MORTEN L. KRINGELBACH
Director, Hedonia: TrygFonden Research Group, University of Oxford and University of Aarhus

STEVEN KYLE
Professor of Applied Economics and Management, Cornell University

ROBERT S. LANGER
David H. Koch Institute Professor, M.I.T.

LAWRENCE LESSIG
Professor, Harvard Law School

ERNEST J. MONIZ
Cecil and Ida Green Distinguished Professor. M.I.T.

JOHN P. MOORE
Professor of Microbiology and Immunology, Weill Medical College of Cornell University

M. GRANGER MORGAN
Professor and Head of Engineering and Public Policy, Carnegie Mellon University

MIGUEL NICOLELIS
Co-director, Center for Neuroengineering, Duke University

MARTIN NOWAK
Director, Program for Evolutionary Dynamics, Harvard University

ROBERT PALAZZO
Provost and Professor of Biology, Rensselaer Polytechnic Institute

CAROLYN PORCO
Leader, Cassini Imaging Science Team, and Director, CICLOPS, Space Science Institute

VILAYANUR S. RAMACHANDRAN
Director, Center for Brain and Cognition, University of California, San Diego

LISA RANDALL
Professor of Physics, Harvard University

MARTIN REES
Professor of Cosmology and Astrophysics, University of Cambridge

JOHN REGANOLD
Regents Professor of Soil Science, Washington State University

JEFFREY D. SACHS
Director, The Earth Institute, Columbia University

EUGENIE SCOTT
Executive Director, National Center for Science Education

TERRY SEJNOWSKI
Professor and Laboratory Head of Computational Neurobiology Laboratory, Salk Institute for Biological Studies

MICHAEL SNYDER
Professor of Genetics, Stanford University School of Medicine

MICHAEL E. WEBBER
Associate Director, Center for International Energy & Environmental Policy, University of Texas at Austin

STEVEN WEINBERG
Director, Theory Research Group, Department of Physics, University of Texas at Austin (Nobel Prize in Physics, 1979)

GEORGE M. WHITESIDES
Professor of Chemistry and Chemical Biology, Harvard University

NATHAN WOLFE
Director, Global Viral Forecasting Initiative

R. JAMES WOOLSEY, JR.
Venture Partner, VantagePoint Venture Partners

ANTON ZEILINGER
Professor of Quantum Optics, Quantum Nanophysics, Quantum Information, University of Vienna

JONATHAN ZITTRAIN
Professor, Harvard Law School

Celebrating Science

In 1845 James K. Polk succeeded John Tyler, Jr., as the 11th president. The U.S. annexed Texas as the 28th state, and the young nation's "manifest destiny" to occupy all of North America became a popular ideal. The industrial revolution was burgeoning, easing people's lives with mechanical marvels. By this time, for instance, Cyrus McCormick had created a labor-saving reaper for crops. And with a promise to explain "New Inventions, Scientific Principles, and Curious Works," the painter and inventor Rufus Porter introduced the first issue of a broadsheet called *The Scientific American* on August 28, 1845.

Porter was "a mechanical Johnny Appleseed sowing seed of new and ingenious ideas as he traveled through New England and abroad through his journals," wrote Jean Lipman in *Rufus Porter, Yankee Pioneer* (Clarkson N. Potter, 1968); you can learn more about him at the Rufus Porter Museum in Bridgton, Me. Porter took out more than 100 patents, but his best-known innovation is his revolver mechanism, which he sold in 1844 to Samuel Colt for $100. *Scientific American* reflected his broad interests. The inaugural edition lists patents, describes developments such as Samuel Morse's telegraph and a filter for locomotive smoke, comments on painting portraits and even includes poetry. Find excerpts in a special 50, 100 and 150 Years Ago, starting on page 12; other excerpts from the 1845 issue and a slide show appear at www.ScientificAmerican.com.

True to Porter's restless nature, he sold the publication after only 10 months to Orson Desaix Munn and Alfred Ely Beach, both in their early 20s. Beach was also an inventor—he designed New York City's first subway, the Beach Pneumatic Transit (an 1870 issue featured the plans). *Scientific American* is included in an exhibit at the Newseum in Washington, D.C., on the history of journalism in the 1800s. Munn & Co. had a successful patent agency that shepherded more than 1,000 applications, and it ultimately held *Scientific American* for more than a century.

When Charles Darwin published *On the Origin of Species*, *Scientific American* covered it. Thomas Edison came into the office and demonstrated his work for the staff. With editorials and silver trophies to reward flights of increasing distances, the editors goaded the Wright brothers to reveal details of their flying machines. Albert Einstein once penned an article for our pages. The magazine saw the U.S. through its Sputnik moment and the start of the "space age."

This is *Scientific American*'s 165-year heritage as the country's oldest continuously published magazine. Yet in every issue, the magazine is new again, with its scientist authors, many of them Nobel Prize winners, and top journalists describing the latest in science and technology. In this edition, feature articles cover everything from threats to the computers we now rely on ("The Hacker in Your Hardware," page 82) to devices that will make "fantastic voyages" in medicine a reality ("Robot Pills," page 62) to extrasolar Earth-like planets that might harbor life ("Planets We Could Call Home," page 38). Fittingly, we include a special report on "Origins" (page 46).

So Happy Birthday to *Scientific American* this month. We hope you will join us in celebrating the renewal and positive force of science in our world every day. ■

Mariette DiChristina
editor in chief

ETHAN HILL (*DiChristina*); *SCIENTIFIC AMERICAN*, VOL. 1, NO. 1; AUGUST 28, 1845 (*logo*)

© 2010 Scientific American

BETWEEN A SMARTER GRID AND A SECURE GRID, THERE IS ONE IMPORTANT WORD: HOW.

Everyone agrees that our nation's aging electric grid must be modernized. To handle a future that includes more renewable sources of energy. A future where consumers want more control over their energy usage. But there are challenges in taking our energy grid into the digital world. That's why we're applying our expertise in cyber security and command and control – to grid management. Applying lessons learned securing our nation's most critical systems. Because it's one thing to make the grid smart. But even smarter to make it secure. Securing our future energy grid is all a question of how. And it is the how that we deliver.

lockheedmartin.com/how

LOCKHEED MARTIN

We never forget who we're working for®

SCIENTIFIC AMERICAN®
Established 1845

EDITOR IN CHIEF: Mariette DiChristina
EXECUTIVE EDITOR: Fred Guterl
MANAGING EDITOR: Ricki L. Rusting
SENIOR WRITER: Gary Stix
EDITORS: Davide Castelvecchi, Mark Fischetti, Michael Moyer, George Musser, Christine Soares, Kate Wong
CONTRIBUTING EDITORS: Mark Alpert, Steven Ashley, Stuart F. Brown, Graham P. Collins, W. Wayt Gibbs, Marguerite Holloway, Christie Nicholson, Michelle Press, John Rennie, Michael Shermer, Sarah Simpson

MANAGING EDITOR, ONLINE: Philip M. Yam
NEWS EDITOR, ONLINE: Robin Lloyd
PODCAST EDITOR: Steve Mirsky
ASSOCIATE EDITORS, ONLINE: David Biello, Larry Greenemeier
NEWS REPORTERS, ONLINE: Katherine Harmon, John Matson
ART DIRECTOR, ONLINE: Ryan Reid

DESIGN DIRECTOR: Michael Mrak
ART DIRECTOR, INFORMATION GRAPHICS: Jen Christiansen
PHOTOGRAPHY EDITOR: Monica Bradley
CONTRIBUTING ART DIRECTOR: Edward Bell

COPY DIRECTOR: Maria-Christina Keller

EDITORIAL ADMINISTRATOR: Avonelle Wing
SENIOR SECRETARY: Maya Harty

COPY AND PRODUCTION, NATURE PUBLISHING GROUP:
SENIOR COPY EDITOR, NPG: Daniel C. Schlenoff
COPY EDITOR, NPG: Michael Battaglia
EDITORIAL ASSISTANT, NPG: Ann Chin
MANAGING PRODUCTION EDITOR, NPG: Richard Hunt
SENIOR PRODUCTION EDITOR, NPG: Michelle Wright

PRODUCTION MANAGER: Christina Hippeli
ADVERTISING PRODUCTION MANAGER: Carl Cherebin
PREPRESS AND QUALITY MANAGER: Silvia De Santis
CUSTOM PUBLISHING MANAGER: Madelyn Keyes-Milch
PRODUCTION COORDINATOR, NPG: Lisa Headley

Letters to the Editor
Scientific American
75 Varick Street, 9th Floor,
New York, NY 10013-1917
or editors@SciAm.com

Letters may be edited for length and clarity. We regret that we cannot answer each one. Post a comment on any article at www.ScientificAmerican.com/sciammag

Growth ▪ More Growth ▪ Faulty Circuits

APRIL 2010

"Isn't growth for the sake of growth the philosophy of cancer?"
—John Langerak CHESTER, ME.

Planning for Earth

Peter H. Gleick's "Freshwater Use" [Solutions to Environmental Threats] neglected to mention one obvious water-conservation measure: meter all freshwater and charge for it. Why should some—most notably agribusiness—receive this valuable resource for free? It is already the policy in some countries to assume that all freshwater is the property of the federal government, owned by all citizens equally. If we were to adopt such a policy in the U.S., we could use the money so collected to install free low-flow toilets, showerheads, and so on in all public housing. It could also be used to partially underwrite farmers converting to drip irrigation and soil-moisture sensors, perhaps in exchange for a promise not to raise food prices as a result of increased irrigation costs. Making everyone pay for what they actually use is always good policy: the profit motive encourages conservation.

Robert Russell
Chapel Hill, N.C.

I am both saddened and dismayed that, once again, you devote the better part of an issue to the problems of sustainability while barely mentioning the root cause of most of the threats to our world: population growth. While I concur with the steps you outline to address biodiversity loss, ocean acidification, climate change, and the like, these are but symptoms of the underlying problem. Ultimately, no technological solution can succeed without tackling the issue of demand. As long as the world's population continues to grow, the incessant pressure for more land, more food and more energy will continue, dwarfing any gains to be had through conservation and technology. Modern medicine and modern fertilizers have triggered the population explosion of the past 200 years, but the human drive to procreate hasn't adjusted to the realities of the modern world. We as a species must voluntarily act to reduce our population to sustainable levels, or else Mother Nature will almost certainly do it for us through war, famine and disease. Increased funding for family planning, tax laws that discourage multiple children, and a shift away from a growth-based economy (which depends in part on an increasing population) must be part of any plan for a sustainable future.

Danny Rosenberg
Portland, Ore.

Jonathan Foley's otherwise excellent "Boundaries for a Healthy Planet" falls short, in my view, in one very important respect. In his introduction he mentions the growth of population and of consumption, saying this has expanded to a global assault and observing that we live in a full world. Yet the rest of the article is confined to discussing ways of limiting the consequences of this growth, without any consideration of limiting growth itself. But should not population itself be subject to a threshold?

The world's population continues to

© 2010 Scientific American

LET'S HELP TO KEEP THE SKIES BLUE.
LET'S GO.

We all need clean air. Not just for today's kite-flying trip, but for future generations who want to live and play under clearer blue skies. That's why, for example, at Shell Brazil we've created a fuel oil for factories that can cut soot emissions by up to 75%. It should help Raul and his friends breathe a little easier. Just one of the many things we're doing to help build a better energy future. Let's go. **www.shell.com/letsgo**

grow at more than 360,000 a day. The most often quoted figure of nine billion in 2050 is generally quoted uncritically, but this figure, now updated to 9.2 billion, is only the middle of three projections by the United Nations Population Fund; the actual figure could be as high as 10.5 billion. This will affect for the worse all the other environmental processes identified in the article, especially freshwater, land use and biodiversity.

A recent article in a special issue of *Scientific American* ["Population & Sustainability," by Robert Engelman; SCIENTIFIC AMERICAN EARTH 3.0, June 2009] pointed out that the number of unintended pregnancies is almost the same as the number by which population increases, so it should be possible to address the problem of population increase in a wholly noncoercive way. Moreover, according to the U.N. Population Fund, 200 million women would use family planning but have no access to it. As a result, according to the World Health Organization, 20 million women undergo unsafe abortions with a horrifying death toll as a consequence. Therefore, there is clearly a huge unfulfilled demand for family planning.

Roger Plenty
Stroud, U.K.

DRIP IRRIGATION is one water-saving technique that could help make agriculture sustainable in an increasingly dry world.

BOB ROWAN *Progressive Image/Corbis*

Kicking the Habit

I did not think I would live long enough to read an article dealing with the subject matter in "Breaking the Growth Habit," by Bill McKibben. During my 72 years as a citizen of the U.S., I've lived out West, back East, up North and down South. There was always a local chamber of commerce stating that if a town was not in a perpetual state of growth, it was not healthy. Isn't growth for the sake of growth the philosophy of cancer?

John Langerak
Chester, Me.

Much of what McKibben writes is silly. The idea of completely stopping economic growth is not only crazy but in one fell swoop dooms the couple of billion people living below the poverty level on Earth, as well as their descendants, to bare subsistence living for eternity. That is not only nuts, it is morally indefensible. His ideas on local agriculture are also way off track.

I happen to have lived in a rural community before "industrial farming" existed. I remember the lean diet we had during winter: root vegetables, cabbage as the only salad, canned beans and tomatoes, etcetera, but no fresh green vegetables at all. I do not want to go back to that, nor do I need to. I like having fresh Chilean blueberries available in midwinter, crisp lettuce all year long as well as a cornucopia of other fresh, healthy produce. If McKibben wants to revert back to the lean old days, fine—but do not preach it to me!

Richard L. Huber
New York City

Cartesian Duality

As an investigator, I appreciate how Thomas R. Insel's "Faulty Circuits" ends with a few comments regarding the transition taking place in psychiatry from a subjective, mental-based discipline to one thoroughly grounded in neuroscience. The intrinsic Cartesian dualist bias that has for so long restrained psychiatry may at long last be loosening its grasp, thus allowing psychiatry to join the other medical specialties, thoroughly grounded in sound scientific practice.

Bob Schwartz
Chester, Vt.

ERRATUM In Edward Bell's "8 Wonders of the Solar System," a caption described the Herschel crater on Saturn's moon Mimas as a 13-kilometer-wide depression. The crater's actual width is 139 kilometers.

SCIENTIFIC AMERICAN®
Established 1845

PRESIDENT: Steven Inchcoombe
EXECUTIVE VICE PRESIDENT: Michael Florek
VICE PRESIDENT, OPERATIONS AND ADMINISTRATION: Frances Newburg

MANAGING DIRECTOR, CONSUMER MARKETING: Christian Dorbandt
ASSOCIATE DIRECTOR, CONSUMER MARKETING: Anne Marie O'Keefe
SENIOR MARKETING MANAGER/RETENTION: Catherine Bussey
SENIOR MARKETING MANAGER/ACQUISITION: Patricia Elliott

VICE PRESIDENT AND PUBLISHER: Bruce Brandfon
VICE PRESIDENT, MARKETING AND SALES DEVELOPMENT: Michael Voss
DIRECTOR, GLOBAL MEDIA SOLUTIONS: Jeremy A. Abbate
MANAGER, INTEGRATED MEDIA SALES: Stan Schmidt
SALES DEVELOPMENT MANAGER: David Tirpack
PROMOTION MANAGER: Diane Schube
MARKETING RESEARCH DIRECTOR: Rick Simone
SALES REPRESENTATIVES: Jeffrey Crennan, Chantel Arroyo

DIRECTOR, ANCILLARY PRODUCTS: Diane McGarvey

How to Contact Us

SUBSCRIPTIONS
For new subscriptions, renewals, gifts, payments, and changes of address: U.S. and Canada, 800-333-1199; outside North America, 515-248-7684 or www.ScientificAmerican.com

SUBMISSIONS
To submit article proposals, follow the guidelines at www.ScientificAmerican.com. Click on "Contact Us." We cannot return and are not responsible for materials delivered to our office.

REPRINTS
To order bulk reprints of articles (minimum of 1,000 copies): Reprint Department, Scientific American, 75 Varick Street, 9th Floor, New York, NY 10013-1917; 212-451-8877; reprints@SciAm.com.
For single copies of back issues: 800-333-1199.

PERMISSIONS
For permission to copy or reuse material: Permissions Department, Scientific American, 75 Varick Street, 9th Floor, New York, NY 10013-1917; 212-451-2810; randp@SciAm.com; www.ScientificAmerican.com/permissions.
Please allow three to six weeks for processing.

ADVERTISING
www.ScientificAmerican.com has electronic contact information for sales representatives of Scientific American in all regions of the U.S. and in other countries.

© 2010 Scientific American

Your music never sounded

QuietComfort® 15

Acoustic Noise Cancelling® headphones

Welcome to a better sounding world, where your music comes alive as never before. The new QC®15 headphones are our best, with Bose® technologies that deliver sound more naturally than conventional headphones. And a significant improvement in the noise reduction helps you focus on each nuance of your music, as distractions fade into the background. Mark A. Kellner says in The Washington Times that these headphones "reproduce music with extreme clarity, fidelity and definition." It's a difference you need to hear to believe. We're so sure you'll be delighted, we'll even pay to ship them to your door.

To learn more: 1-800-760-2749, ext. Q8018
Bose.com/headphones

©2010 Bose Corporation. Patent rights issued and/or pending. The distinctive design of the headphone oval ring is a trademark of Bose Corporation. Quote reprinted with permission.

BOSE®
Better sound through research®



even though she had had no contact with males. Virgin births are the norm for the flower-pot snake, a female-only creature that has spread throughout the world, one individual at a time. Mammals, including humans, appear to have been denied the cloning option, however. Our lives seem fated to include plenty of sex, in good times and in bad. —*Brendan Borrell*

that infects human beings—*P. falciparum*—evolved from another version of the parasite, *P. reichenowi,* which currently infects chimpanzees. And it happened a mere 10,000 years ago—a moment in evolutionary terms.

The finding rests on a molecular comparison of the genomes of the two parasites. Stephen Rich, an evolutionary geneticist at the University of Massachusetts Amherst, and his colleagues measured the diversity of the genomes, a rough proxy for age (genomes tend to acquire genetic components over time). *Reichenowi*'s genome can be 20 times more diverse than *falciparum*'s, which means *reichenowi* is much older. "It seems that malaria has been in chimps as long as they've been chimps," Rich says.

Following the trail back to the origin of *reichenowi* is a more complicated problem, not least because malaria is so widespread. "In terrestrial vertebrates, we find it virtually everywhere we look," Rich observes. "We're only getting started." —*Mike May*

DAVID M. PHILLIPS *Photo Researchers, Inc.* (*sperm and egg*); GETTY IMAGES (*firecrackers*)

SNAP, CRACKLE, BANG

The ancient Chinese invented fireworks to scare off 10-foot-tall mountain men

THAT RAUCOUS RITE OF SUMMER—the fireworks display—may have started as a scholarly tradition in ancient China. Before the Chinese got around to inventing paper in the second century A.D., scribes, using a stylus, would etch ideograms on the rounded surface of green bamboo stalks. The medium served as a way for recording transactions and stories. As the stalks dried over the fire, air pockets in the wood would often burst with a loud cracking noise.

The noise, of course, gradually became the whole point of the exercise. The classic *I Ching,* or *Book of Changes,* explains how the cracks and pops succeeded in scaring off the Shan Shan, 10-foot-tall mountain men. Later, the Chinese spiced things up by adding gunpowder to the stalks.

The first fireworks display didn't take place until the 12th century rolled around. In 1267 English philosopher Roger Bacon wrote about "that toy of children" and the "horrible sound" it produces, which "exceeds the roar of sharp thunder." Those sharp bangs evoke nothing more than, yes, another Fourth of July celebration. —*Mike May*

© 2010 Scientific American

THORNY FENCE

The invention of barbed wire was a huge commercial success—and the subject of furious legal battles

AT SOME POINT IN THE HISTORY of civilization, shepherding gave way to farming. That created a need for some way of keeping cows and pigs from wandering freely through the meadows. The fence was born. Wooden fences were among the earliest, but they are expensive and time-consuming to build. By 1870 smooth cable was easy to get hold of and came into wide use on ranches. Cattle would rub their back on the wire, and sometimes one would slip through. Eventually the herds caught on.

That got Michael Kelly, an inventor from New York City, wondering how he might make the wire less comfortable as a bovine back scratcher. He got the idea to twist bits of sharp pointed wire onto ordinary cable, and in 1868 he patented his "thorny fence." It was a big success—and a magnet for lawsuits. "Almost overnight it developed into a source of wealth and furious litigation colored by impassioned charges and countercharges of patent infringement and greed," says historian Robert T. Clifton.

Joseph Glidden of DeKalb, Ill., also hit legal snags over an improved wire that used two strands to lock the barbs in place. In 1892 his case went before the U.S. Supreme Court, which ruled in his favor, making him the undisputed father of an invention that more than any other marked the closing of the West's open range. *—Mike May*

SCRUBS

A rise in maternity ward deaths led one physician to discover the importance of hand washing

IN THE MID-1840S Hungarian physician Ignaz Semmelweis saw with alarm that 15 percent of new mothers in his Vienna General Hospital were dying of an illness called puerperal fever. Semmelweis was desperate to prevent the illnesses, but he didn't know how. As he pondered the problem, he learned that his friend, forensic pathologist Jakob Kolletschka, had died from what sounded like the same illness. It happened only a few days after a student accidentally pricked Kolletschka with a scalpel that had been used to dissect a cadaver.

The news gave Semmelweis pause. Medical students at his hospital would routinely go right from the morgue to the maternity ward without ever washing their hands. Were they carrying an infection to the mothers? Was that why they were dying? Could hand washing help?

To test his dirty-hands hypothesis, Semmelweis made his students wash their hands in a mixture of water and chlorine (soap and water did not eliminate the cadaver smell). Fevers in the maternity ward quickly dropped by 10 percent. Hand washing became standard procedure at Semmelweis's hospital.

It took 40 years for the policy to take hold widely. Even today hospital workers don't follow it as consistently as they should. According to an ongoing study from the Maryland Health Quality and Cost Council, 90 percent of staff wash their hands when someone is looking, but only 40 percent do when alone. *—Mike May*

STEVEN PUETZER *Getty Images* (*hand washing*); GETTY IMAGES (*barbed wire*)

© 2010 Scientific American

MORAL ANIMAL

A sense of right and wrong starts with innate brain circuitry

THE ROOTS OF MODERN MORALITY have long been a point of contention among psychologists, philosophers and neuroscientists. Do our ethical foundations arise from our relatively recent ability to reason or from our ancient emotions? Studies have recently lent support to the notion that we owe much of our sense of right and wrong to our animal ancestors.

Evidence that morality comes before reason is supported by primate studies. A chimpanzee, for instance, will sometimes drown to save its peers and refuse food if doing so prevents others from injury. That's not to say they are morally sophisticated beings, but "it's not as if morality and our moral rules are just a pure invention of the religious or philosophical mind," explains Frans de Waal, a primatologist and psychologist at Emory University. De Waal's work suggests that our morality is an outgrowth of our ancestors' social tendencies, an indication that it is at least in part an evolved trait (an idea Charles Darwin shared). Dogs, too, seem to have a keen sense of "wild justice," says Marc Bekoff, a professor emeritus at the University of Colorado at Boulder. He has observed a sense of morality among dogs at play. "Animals know right from wrong," he notes.

If morality is innate rather than learned, then it should have left biological traces. Studies suggest that moral decisions involve certain parts of the brain associated with prosocial tendencies and emotional regulation, such as the ventromedial prefrontal cortex. In brain scans, this region lights up when subjects choose to donate money to charity, and those with damage to this region make unexpected moral judgments. Some ethical dilemmas also activate brain regions involved in rational decision making, such as one called the anterior cingulate cortex—a finding that implies that higher-order brain functions may also contribute to our morality, even if it's rooted in emotions.

Ultimately, de Waal says, we need to thank our evolutionary ancestors for far more than just bestial urges. "When humans kill each other or commit genocide, we say we're acting like animals," he says. But "you can see the same sort of thing with regard to our positive behavior."

—Melinda Wenner Moyer



URBAN BUG

Packed living conditions made the influenza virus a leading public health threat

HIPPOCRATES DESCRIBED THE SYMPTOMS of the flu some 2,400 years ago. But the influenza virus didn't become a true menace until the rise of stable, densely populated settlements and the growth of animal husbandry. This crowding of people and their animals furnished the virus with ample opportunities to jump from one species to another, acquiring deadly attributes along the way.

The first influenza pandemics were recorded during the 1500s. The one that occurred in 1580 traced a path that epidemiologists today would recognize: it began in Asia during the summer and then spread to Africa, Europe and America over the next six months. Another big epidemic hit in 1789, the year that George Washington took office, "before modern means of rapid travel were available and when a man could go no faster than his horse could gallop," wrote virologist and epidemiologist Richard E. Shope in 1958. Even so, he said, it "spread like wildfire."

Shope knew influenza well: in 1931 he became the first scientist to transmit the virus between two animals, by transferring mucus from one pig's nose to another's. Because Shope had filtered bacteria from the mucus beforehand, his experiment suggested, for the first time, that the flu was caused by a virus. Two years later a group of U.K. scientists became the first to isolate a human form of the virus, from a sick ferret. *—Melinda Wenner Moyer*

FORMER LIFE OF THE ELECTRIC CAR

A century ago taxicabs had batteries, not gas guzzlers, under their hoods

IN WHAT MAY HAVE BEEN THE FIRST attempt at an electric car, Scottish inventor Robert Anderson built a "crude electric carriage" in the mid- to late 1830s. It didn't get far. For one thing, its battery wasn't good enough. (Today's green car engineers can sympathize.) It also faced stiff competition from steam-powered cars.

When rechargable batteries started to appear in the mid-1800s, electric vehicles got a fillip. In 1897 the Electric Carriage and Wagon Company in Philadelphia assembled a fleet of electric-powered taxis for New York City. By 1902 the Pope Manufacturing Company in Hartford, Conn., had built around 900 electric vehicles, most of which were used as cabs. That same year Studebaker, which had gotten its start in horse-drawn wagons, entered the car market in Indiana with an electric model. Through the early 1900s electric vehicles ran smoother and quieter than their gas-guzzling, internal-combustion-engine-powered rivals.

Where the electric car stumbled was range—it couldn't go far between rechargings. By 1920 gas-powered rivals emerged as the clear winner, a twist of history that engineers are now working furiously to undo. *—Mike May*

ALFRED PASIEKA *Photo Researchers, Inc.* (*flu virus*); CORBIS BETTMANN (*carriage*)

© 2010 Scientific American

THE FIRST HUMVEE

Wheeled vehicles may have first arisen as a tool of war

SIR C. LEONARD WOOLLEY'S 1922 excavation of the Royal Cemetery of Ur—a Sumerian site located in modern-day Iraq—was, by early 20th-century standards, a major media event. Thomas Edward Lawrence, a.k.a. Lawrence of Arabia, who had achieved fame for his dashing exploits during the Arab Revolt several years earlier, helped to organize the expedition. British mystery writer Agatha Christie paid a visit to the site and penned *Murder in Mesopotamia* as a tribute (she would later marry Woolley's assistant). All this fuss over a box with a picture of a wheel on it.

It wasn't just any box, of course. It was the Standard of Ur, a 4,600-year-old container, the size of a shoebox (*above*), encrusted in lapis lazuli. Most important, it featured an illustration of ancient warfare that included the oldest uncontested image of the wheel in transportation. A series of images depicted tanklike carriages, each with four solid wheels braced to their axles and a team of horses propelling them forward. The wheeled carriages clearly provided soldiers with better protection against ambush than the poor foot soldiers had, who are shown squirming to avoid horses' hooves.

This ancient Humvee wasn't the only way fifth-millennium engineers deployed wheels. The Sumerians, Egyptians and Chinese all used wheels for spinning pots, and Egyptians moved massive stones with log rollers. Wheels never caught on for ordinary transport because they weren't useful on the sandy soils of the world's trade routes, says Richard Olson, a historian and author. Camels remained the all-terrain vehicle of choice for another 2,000 years or so.

Wheeled vehicles didn't take off until the advent of roads. The Egyptians built extensive dirt roads and paved some of them with sandstone, limestone and even a surfacing of petrified wood. By as far back as 3,500 years ago, they fashioned a metal wheel with six spokes, and from the Middle East to Russia, agile, two-wheeled chariots became all the rage. *—Brendan Borrell*

GRAVITY'S TUG

The first black holes are almost as old as the universe itself

THE IDEA THAT A BLACK HOLE could possibly exist came from an English rector, John Michell. In 1783 he calculated that the force of gravity exerted by a massive star could prevent light from escaping its surface. Michell's work was largely forgotten for 200 years. In 1971 astrophysicists noticed flickering x-rays coming from the constellation Cygnus, 6,000 light-years away: the radiation indicated that a black hole was apparently circling a star. As with any black hole, it formed as a star ran out of fuel and collapsed in on itself. If the sun were to somehow become a black hole, it would be less than three miles across, trapping light in the warped space that enfolds it. For Earth to become a black hole, it would be the size of a marble.

The first black holes in the universe arose nearly 14 billion years ago, contends Abraham Loeb, an astrophysicist at Harvard University. At that time, gas began to condense into clouds that fragmented into massive stars 100 times the size of the sun, which in turn collapsed into black holes. Fortunately, the spinning of early galaxies limited the growth of the black holes at their cores, allowing stars to form.

Physicists have now begun to make something akin to black holes on Earth. Chinese researchers built concentric cylinders that mimic a black hole, bending microwave radiation in on itself as it passes from outer to inner surfaces. And a real black hole could still improbably pop out of the Large Hadron Collider near Geneva. *—Brendan Borrell*

© 2010 Scientific American

RAINBOW CELLS

Biodiversity was the first step toward complex life

Witnesses were absent for the comings and goings of the first life some four billion years ago, but scientists are pretty sure the typical Earth creature in those days consisted of no more than a single cell. That doesn't mean the planet was a dull sea of sameness. Single-celled creatures may have acquired genetic diversity early on.

Here's why. When cells divide, mistakes have a way of creeping into genetic material. Variants that enhance a cell's ability to survive and reproduce become more common over successive generations. This basic fact of evolution applied to the early Earth. "Variation is necessary for there to be evolution by natural selection in the first place," explains Andrew Hamilton, a philosopher of science at Arizona State University. "Biodiversity originated at the point that there was variation on which selection could operate."

Today we think of biodiversity in terms of multicellular life, but flowering plants and animals didn't arrive until relatively recently (540 million years ago). Although some evidence suggests that having a wide variety of species makes an ecosystem more stable, the jury is still out. It is of no comfort to know that the worst catastrophe would still preserve some biodiversity—even if only for the lowly cell. —*Melinda Wenner Moyer*



© TRUSTEES OF BRITISH MUSEUM (*container*); GARCIA/CORBIS (*pasta*)

ZERO

How nothing became something

NOBODY KNEW HOW MUCH we needed nothing until we had a number for it. Without zero, negative and imaginary numbers would have no meaning, and it would be impossible to solve quadratic equations, a mainstay of applied math. Without zero to act as a placeholder to distinguish, say, 10 from 100, all but the simplest arithmetic requires an abacus or counting board. "If we didn't have zero, our system of numbers would be incomplete," says Charles Seife, author of *Zero: The Biography of a Dangerous Idea.* "It would really break down without zero."

Zero arrived on the scene in two installments. Around 300 B.C., Babylonians developed a proto-zero—two slanted wedges pressed into clay tablets—that served as a placeholder in their funky sexagesimal, or base 60, number system. By the fifth century, the concept of zero migrated to India and made its symbolic entrance as a dot carved on a wall at the Chaturbhuja Temple in Gwalior. Then, like a pebble dropped into a puddle, the symbol for zero expanded to an "0" and became a number with properties all its own: an even number that is the average of –1 and 1. In 628 mathematician Brahmagupta pontificated on the frightening properties of zero: multiply anything by zero, and it, too, turns to nothing. Independently, Mayans in the Americas developed their own zero to assist in the study of astronomy.

Over time the expansion of the Islamic empire spread the Indian zero back to the Middle East and, eventually, to the Moors in Spain, where it became one of 10 Arabic numerals, as we refer to them today. European scholars clung to their Roman numerals. Zero's official endorsement by the Western world came by way of Italian mathematician Fibonacci (Leonardo of Pisa), who included it in a textbook in 1202.

—*Brendan Borrell*

NOODLING THE NOODLES

It took thousands of years to go from mush to spaghetti

FOOD HISTORIAN FRANCINE SEGAN asserts that pasta emerged more than 5,000 years ago when an enterprising chef happened upon the now seemingly obvious idea of mushing flour and water together to create something that looked surprisingly like lasagna. "It breaks my heart to tell you this," says Segan, invoking her own Italian heritage, "but the first to make those noodles might have been the ancient Greeks. Lots of references in ancient Greek writing—even in 3,000 B.C.—talk about layers that sound a lot like lasagna."

Spaghetti took longer but appears to have taken shape in Italy. A popular misconception has Marco Polo introducing Italians to pasta when he returned from China in 1295, but Italy already had pasta by then. "In Sicily there is a town called Trabia," wrote Arab geographer Muhammad al-Idrisi in 1154. "In this town they made a food of flour in the form of strings." He describes a true pasta industry: foodstuffs were first dried in the sun and then shipped by boat to other regions of Italy and even other countries.

A few hundred years later Leonardo da Vinci invented a machine that turned dough into edible strings. Technical glitches kept his pasta maker from achieving the mechanization of the industry he had hoped for. Still, the Italians succeeded in refining the art of pasta making, crafting more elaborate forms of mushed dough than anyone else.

—*Mike May*

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

© 2010 Scientific American

Cave today known simply as PP13B, near Mossel Bay, South Africa, sheltered humans between 164,000 and 35,000 years ago, at a time when *Homo sapiens* was in danger of dying out. These people may have been the ancestors of us all.

© 2010 Scientific American

When the Sea Saved Humanity

Shortly after *Homo sapiens* arose, harsh climate conditions nearly extinguished our species. Recent finds suggest that the small population that gave rise to all humans alive today survived by exploiting a unique combination of resources along the southern coast of Africa

BY CURTIS W. MAREAN

WITH THE GLOBAL POPULATION of humans currently approaching seven billion, it is difficult to imagine that *Homo sapiens* was once an endangered species. Yet studies of the DNA of modern-day people indicate that, once upon a time, our ancestors did in fact undergo a dramatic population decline. Although scientists lack a precise timeline for the origin and near extinction of our species, we can surmise from the fossil record that our forebears arose throughout Africa shortly before 195,000 years ago. Back then the climate was mild and food was plentiful; life was good. But around 195,000 years ago, conditions began to deteriorate. The planet entered a long glacial stage known as Marine Isotope Stage 6 (MIS6) that lasted until roughly 123,000 years ago.

A detailed record of Africa's environmental conditions during glacial stage 6 does not exist, but based on more recent, better-known glacial stages, climatologists surmise that it was almost certainly cool and arid and that its deserts were probably significantly expanded relative to their modern extents. Much of the landmass would have been uninhabitable. While the planet was in the grip of this icy regime, the number of people plummeted perilously—from more than 10,000 breeding individuals to just hundreds. Estimates of exactly when this bottleneck occurred and how small the population became vary among genetic studies, but all of them indicate that everyone alive today is descended from a small population that lived in one region of Africa sometime during this global cooling phase.

I began my career as an archaeologist working in East Africa and studying the origin of modern humans. But my interests began to shift when I learned of the population bottleneck that geneticists had started talking about in the early 1990s. Humans today exhibit very low genetic diversity relative to many other species with much smaller population sizes and geographic ranges—a phenomenon best explained by the occurrence of a population crash in early *H. sapiens*. Where, I wondered, did our ancestors manage to survive during the climate catastrophe? Only a handful of regions could have had the natural resources to support hunter-gatherers. Paleoanthropologists argue vociferously over which of these areas was the ideal spot. The southern coast of Africa, rich in shellfish and edible plants year-round, seemed to me as if it would have been a particularly good refuge in tough times. So, in 1991, I decided I would go there and look for sites with remains dating to glacial stage 6.

KEY CONCEPTS

- At some point between 195,000 and 123,000 years ago, the population size of *Homo sapiens* plummeted, thanks to cold, dry climate conditions that left much of our ancestors' African homeland uninhabitable. Everyone alive today is descended from a group of people from a single region who survived this catastrophe.
- The southern coast of Africa would have been one of the few spots where humans could survive during this climate crisis, because it harbors an abundance of shellfish and edible plants.
- Excavations of a series of sites in this region have recovered items left behind by what may have been that progenitor population.
- The discoveries confirm the idea that advanced cognitive abilities evolved earlier than previously thought—and may have played a key role in the survival of the species during tough times.

—The Editors

PER-ANDERS PETTERSON

© 2010 Scientific American

SEASIDE SANCTUARY

Between 195,000 and 123,000 years ago, the planet was locked in an ice age known as Marine Isotope Stage 6, rendering much of the African continent cool and arid—unsuitable for the plants and animals that *Homo sapiens* ate. Only a few regions could have supported our species, namely, those with grassland or Mediterranean scrub vegetation. The southern coast would have been a particularly plentiful oasis, thanks to the edible fynbos plants that grow only here and the dense shellfish beds nurtured by the Agulhas current and the Benguela upwelling of nutrient-rich cold water from the sea bottom.

[THE AUTHOR]

CURTIS W. MAREAN is a professor at the School of Human Evolution and Social Change at Arizona State University and a member of the Institute of Human Origins. He studies the origins of modern humans, the prehistory of Africa, paleoclimates and paleoenvironments, and animal bones from archaeological sites. Marean is particularly interested in human occupation of coastal ecosystems. He is the principal investigator for the South African Coast Paleoclimate, Paleoenvironment, Paleoecology, Paleoanthropology (SACP4) project, funded by the National Science Foundation.

My search within that coastal area was not random. I had to find a shelter close enough to the ancient coastline to provide easy access to shellfish and elevated enough that its archaeological deposits would not have been washed away 123,000 years ago when the climate warmed and sea levels surged. In 1999 my South African colleague Peter Nilssen and I decided to investigate some caves he had spotted at a place called Pinnacle Point, a promontory near the town of Mossel Bay that juts into the Indian Ocean. Scrambling down the sheer cliff face, we came across a cave that looked particularly promising—one known simply as PP13B. Erosion of the sedimentary deposits located near the mouth of the cave had exposed clear layers of archaeological remains, including hearths and stone tools. Even better, a sand dune and a layer of stalagmite capped these remnants of human activity, suggesting that they were quite old. By all appearances, we had hit the jackpot. The following year, after a local ostrich farmer built us a 180-step wooden staircase to allow safer access to the site, we began to dig.

Since then, my team's excavations at PP13B and other nearby sites have recovered a remarkable record of the activities undertaken by the people who inhabited this area between approximately 164,000 and 35,000 years ago, hence during the bottleneck and after the population began to recover. The deposits in these caves, combined with analyses of the ancient environment there, have enabled us to piece together a plausible account of how the prehistoric residents of Pinnacle Point eked out a living during a grim climate crisis. The remains also debunk the abiding notion that cognitive modernity evolved long after anatomical modernity: evidence of behavioral sophistication abounds in even the oldest archaeological levels at PP13B. This advanced intellect no

PER-ANDERS PETTERSON (*Marean*); DON FOLEY (*map*); LUCY READING-IKKANDA (*globe*)

© 2010 Scientific American

HIGH AND DRY

Finding archaeological sites dating to glacial stage 6 required searching for shelters that were close enough to the sea to allow relatively easy access to shellfish yet elevated enough that their ancient remains would not have washed away when the sea level rose 123,000 years ago. PP13B and other caves carved into the sheer cliff face of a promontory called Pinnacle Point meet those requirements and have yielded a plethora of remains dating to this critical juncture in human prehistory.

PER-ANDERS PETTERSON

doubt contributed significantly to the survival of the species, enabling our forebears to take advantage of the resources available on the coast.

While elsewhere on the continent populations of *H. sapiens* died out as cold and drought claimed the animals and plants they hunted and gathered, the lucky denizens of Pinnacle Point were feasting on the seafood and carbohydrate-rich plants that proliferated there despite the hostile climate. As glacial stage 6 cycled through its relatively warmer and colder phases, the seas rose and fell, and the ancient coastline advanced and retreated. But so long as people tracked the shore, they had access to an enviable bounty.

A COASTAL CORNUCOPIA

FROM A SURVIVAL STANDPOINT, what makes the southern edge of Africa attractive is its unique combination of plants and animals. There a thin strip of land containing the highest diversity of flora for its size in the world hugs the shoreline. Known as the Cape Floral Region, this 90,000-square-kilometer strip contains an astonishing 9,000 plant species, some 64 percent of which live only there. Indeed, the famous Table Mountain that rises above Cape Town in the heart of the Cape Floral Region has more species of plants than does the entire U.K. Of the vegetation groups that occur in this realm, the two most extensive are the fynbos and the renosterveld, which consist largely of shrubs. To a human forager equipped with a digging stick, they offer a valuable commodity: the plants in these groups produce the world's greatest diversity of geophytes—underground energy-storage organs such as tubers, bulbs and corms.

Geophytes are an important food source for modern-day hunter-gatherers for several reasons. They contain high amounts of carbohydrate; they attain their peak carbohydrate content reliably at certain times of year; and, unlike aboveground fruits, nuts and seeds, they have few predators. The bulbs and corms that dominate the Cape Floral Region are additionally appealing because in contrast to the many geophytes that are highly fibrous, they are low in fiber relative to the amount of energy-rich carbohydrate they contain, making them more easily digested by children. (Cooking further enhances their digestibility.) And because geophytes are adaptations to dry conditions, they would have been readily available during arid glacial phases.

The southern coast also has an excellent source of protein to offer, despite not being a prime hunting ground for large mammals. Just offshore, the collision of nutrient-rich cold waters from the Benguela upwelling and the warm Agulhas current creates a mix of cold and warm eddies along the southern coast. This varied ocean environment nurtures diverse and dense beds of shellfish in the rocky intertidal zones and sandy beaches. Shellfish are a very high quality source of protein and omega-3 fatty acids. And as with geophytes, glacial cooling does not depress their numbers. Rather, lower ocean temperatures result in a greater abundance of shellfish.

SURVIVAL SKILLS

WITH ITS COMBINATION of calorically dense, nutrient-rich protein from the shellfish and low-fiber, energy-laden carbs from the geophytes, the southern coast would have provided an ideal diet for early modern humans during glacial stage 6. Furthermore, women could obtain both these

© 2010 Scientific American

GONE SHELLFISHING

Shellfish, which are rich in protein, are thought to have aided survival of the Pinnacle Point population because they abound year-round in the rocky intertidal zone along the southern coast of Africa (*upper left*). Brown mussels (*lower left*) have turned up in even the earliest levels of PP13B, dating to 164,000 years ago, revealing that humans began exploiting marine resources earlier than previously thought. In addition to mussels, the occupants of the Pinnacle Point sites collected various kinds of limpets as well as sea snails called alikreukel for food and gathered empty shells of helmet snails for their aesthetic appeal (*right*).

PER-ANDERS PETTERSON

resources on their own, freeing them from relying on men to provision them and their children with high-quality food. We have yet to unearth proof that the occupants of PP13B were eating geophytes—sites this old rarely preserve organic remains—although younger sites in the area contain extensive evidence of geophyte consumption. But we have found clear evidence that they were dining on shellfish. Studies of the shells found at the site conducted by Antonieta Jerardino of the University of Barcelona show that people were gathering brown mussels and local sea snails called alikreukel from the seashore. They also ate marine mammals such as seals and whales on occasion.

Previously the oldest known examples of humans systematically using marine resources dated to less than 120,000 years ago. But dating analyses performed by Miryam Bar-Matthews of the Geological Survey of Israel and Zenobia Jacobs of University of Wollongong in Australia have revealed that the PP13B people lived off the sea far earlier than that: as we reported in 2007 in the journal *Nature,* marine foraging there dates back to a stunning 164,000 years ago. By 110,000 years ago the menu had expanded to include species such as limpets and sand mussels.

This kind of foraging is harder than it might seem. The mussels, limpets and sea snails live on the rocks in the treacherous intertidal zone, where an incoming swell could easily knock over a hapless collector. Along the southern coast, safe harvesting with sufficiently high returns is only possible during low spring tides, when the sun and moon align, exerting their maximum gravitational force on the ebb and flow of the water. Because the tides are linked to the phases of the moon, advancing by 50 minutes a day, I surmise that the people who lived at PP13B—which 164,000 years ago was located much farther inland, two to five kilometers from the water, because of lower sea levels—scheduled their trips to the shore using a lunar calendar of sorts, just as modern coastal people have done for ages.

Harvesting shellfish is not the only advanced behavior in evidence at Pinnacle Point as early as 164,000 years ago. Among the stone tools are significant numbers of "bladelets"—tiny flakes twice as long as they are wide—that are too small to wield by hand. Instead they must have been attached to shafts of wood and used as projectile weapons. Composite toolmaking is indicative of

© 2010 Scientific American

considerable technological know-how, and the bladelets at PP13B are among the oldest examples of it. But we soon learned that these tiny implements were even more complex than we thought.

Most of the stone tools found at coastal South African archaeological sites are made from a type of stone called quartzite. This coarse-grained rock is great for making large flakes, but it is difficult to shape into small, refined tools. To manufacture the bladelets, people used fine-grained rock called silcrete. There was something odd about the archaeological silcrete, though, as observed by Kyle S. Brown of the Institute of Human Origins at Arizona State University, an expert stone tool flaker on my team. After years of collecting silcrete from all over the coast, Brown determined that in its raw form the rock never has the lustrous red and gray coloring seen in the silcrete implements at Pinnacle Point and elsewhere. Furthermore, the raw silcrete is virtually impossible to shape into bladelets. Where, we wondered, did the toolmakers find their superior silcrete?

A possible answer to this question came from Pinnacle Point Cave 5-6, where one day in 2008 we found a large piece of silcrete embedded in ash. It had the same color and luster seen in the silcrete found at other archaeological deposits in the region. Given the association of the stone with the ash, we asked ourselves whether the ancient toolmakers might have exposed the silcrete to fire to make it easier to work with—a strategy that has been documented in ethnographic accounts of native North Americans and Australians. To find out, Brown carefully "cooked" some raw silcrete and then attempted to knap it. It flaked wonderfully, and the flaked surfaces shone with the same luster seen in the artifacts from our sites. We thus concluded that the Stone Age silcrete was also heat-treated.

We faced an uphill battle to convince our colleagues of this remarkable claim, however. It was archaeology gospel that the Solutrean people in France invented heat treatment about 20,000 years ago, using it to make their beautiful tools. To bolster our case, we used three independent techniques. Chantal Tribolo of the University of Bordeaux performed what is called thermoluminescence analysis to determine whether the silcrete tools from Pinnacle Point were intentionally heated. Then Andy Herries of the University of New South Wales in Australia employed magnetic susceptibility, which looks for changes in the ability of rock to be magnetized—another indicator of heat exposure among iron-rich rocks. Finally, Brown used a gloss meter to measure the luster that develops after heating and flaking and compare it with the luster on the tools he made. Our results, detailed last year in the journal *Science,* showed that intentional heat treatment was a dominant technology at Pinnacle Point by 72,000 years ago and that people there employed it intermittently as far back as 164,000 years ago.

The process of treating by heat testifies to two uniquely modern human cognitive abilities. First, people recognized that they could substantially alter a raw material to make it useful—in this case, engineering the properties of stone by heating it, thereby turning a poor-quality rock into high-quality raw material. Second, they could invent and execute a long chain of processes. The making of silcrete blades requires a complex series of carefully designed steps: building a sand pit to insulate the silcrete, bringing the heat slowly up to 350 degrees Celsius, holding the temperature steady and then dropping it down slowly. Creating and carrying out the sequence and passing technologies down from generation to generation probably required language. Once established, these abilities no doubt helped our ancestors outcompete the archaic human species they encountered once they dispersed from Africa. In particular, the complex pyrotechnology detected at Pinnacle Point would have given early modern humans a distinct advantage as they en-

DIGGING FOR DINNER

Geophytes, the underground energy-storage organs of certain kinds of plants (*left*), swell with edible carbohydrates at certain times of the year. The distinctive vegetation that hugs the southern coast of Africa, particularly the fynbos plants (*above*), produces especially nutritious and easily digested geophytes, which presumably served as a staple for the early modern humans who lived in this region during glacial stage 6.

COURTESY OF J. C. DE VYNCK

© 2010 Scientific American

CUTTING-EDGE TECHNOLOGY

Stone tools found in PP13B include sophisticated implements such as microblades (*bottom two rows*), which would have been attached to a wooden shaft to form projectile weapons. The toolmakers also appear to have heat-treated the stone to make it easier to shape—a technique that was believed to have originated much later and in France.

PER-ANDERS PETTERSON

tered the cold lands of the Neandertals, who seem to have lacked this technique.

SMART FROM THE START

IN ADDITION to being technologically savvy, the prehistoric denizens of Pinnacle Point had an artistic side. In the oldest layers of the PP13B sequence, my team has unearthed dozens of pieces of red ochre (iron oxide) that were variously carved and ground to create a fine powder that was probably mixed with a binder such as animal fat to make paint that could be applied to the body or other surfaces. Such decorations typically encode information about social identity or other important aspects of culture—that is, they are symbolic. Many of my colleagues and I think that this ochre constitutes the earliest unequivocal example of symbolic behavior on record and pushes the origin of such practices back by tens of thousands of years. Evidence of symbolic activities also appears later in the sequence. Deposits dating to around 110,00 years ago include both red ochre and seashells that were clearly collected for their aesthetic appeal, because by the time they washed ashore from their deepwater home, any flesh would have been long gone. I think these decorative seashells, along with the evidence for marine foraging, signal that people had, for the first time, begun to embed in their worldview and rituals a clear commitment to the sea.

The precocious expressions of both symbolism and sophisticated technology at Pinnacle Point have major implications for understanding the origin of our species. Fossils from Ethiopia show that anatomically modern humans had evolved by at least 195,000 years ago. The emergence of the modern mind, however, is more difficult to establish. Paleoanthropologists use various proxies in the archaeological record to try to identify the presence and scope of cognitive modernity. Artifacts made using technologies that require outside-the-box connections of seemingly unrelated phenomena and long chains of production—like heat treatment of rock for tool manufacture—are one proxy. Evidence of art or other symbolic activities is another, as is the tracking of time through proxies such as lunar phases. For years the earliest examples of these behaviors were all found in Europe and dated to after 40,000 years ago. Based on that record, researchers concluded that there was a long lag between the origin of our species and the emergence of our peerless creativity.

But over the past 10 years archaeologists working at a number of sites in South Africa have found examples of sophisticated behaviors that predate by a long shot their counterparts in Europe. For instance, archaeologist Ian Watts, who works in South Africa, has described hundreds to thousands of pieces of worked and unworked ochre at sites dating as far back as 120,000 years ago. Interestingly, this ochre, as well as the pieces at Pinnacle Point, tends to be red despite the fact that local sources of the mineral exhibit a range of hues, suggesting that humans were preferentially curating the red pieces—perhaps associating the color with menstruation and fertility. Jocelyn A. Bernatchez, a Ph.D. student at Arizona State, thinks that many of these ochre pieces may have been yellow originally and then heat-treated to turn them red. And at Blombos Cave, located about 100 kilometers west of Pinnacle Point, Christopher S. Henshilwood of the University of Bergen in Norway has discovered pieces of ochre with systematic engravings, beads made of snail shells and refined bone tools, all of which date to around 71,000 years ago [see "The Morning of the Modern Mind," by Kate Wong; SCIENTIFIC AMERICAN, June 2005]. These sites, along with those at Pin-

© 2010 Scientific American

nacle Point, belie the claim that modern cognition evolved late in our lineage and suggest instead that our species had this faculty at its inception.

I suspect that a driving force in the evolution of this complex cognition was strong long-term selection acting to enhance our ancestors' ability to mentally map the location and seasonal variation of many species of plants in arid environments and to convey this accumulated knowledge to offspring and other group members. This capacity laid the foundation for many other advances, such as the ability to grasp the link between the phases of the moon and the tides and to learn to schedule their shellfish-hunting trips to the shore accordingly. Together the readily available shellfish and geophytes provided a high-quality diet that allowed people to become less nomadic, increased their birth rates and reduced their child mortality. The larger group sizes that resulted from these changes would have promoted symbolic behavior and technological complexity as people endeavored to express their social identity and build on one another's technologies, explaining why we see such sophisticated practices at PP13B.

FOLLOW THE SEA

PP13B PRESERVES a long record of changing occupations that, in combination with the detailed records of local climate and environmental change my team has obtained, is revealing how our ancestors used the cave and the coast over millennia. Modeling the paleocoastline over time, Erich C. Fisher of the University of Florida has shown that the conditions changed quickly and dramatically, thanks to a long, wide, gently sloping continental shelf off the coast of South Africa called the Agulhas bank. During glacial periods, when sea levels fell, significant amounts of this shelf would have been exposed, putting considerable distance—up to 95 kilometers—between Pinnacle Point and the ocean. When the climate warmed and sea levels rose, the water advanced over the Agulhas bank again, and the caves were seaside once more.

Judging from rainfall and vegetation patterns evident in records from stalagmites spanning the time between 350,000 and 50,000 years ago, we see that the fynbos probably followed the retreating coast out onto the now submerged continental shelf and back again, keeping the geophytes and shellfish in close proximity. As for the people, during these periods of low population density they were free to target the best part of the landscape, and that was the intersection of the geophytes and shellfish—so I suspect they followed the sea. The tracking of resources would explain why PP13B appears to have been occupied intermittently.

Our excavations at PP13B have intercepted the people who may very well be the ancestors of everyone on the planet as they shadowed the shifting shoreline. Yet if I am correct about these people and their connection to the coast, the richest record of the progenitor population lies underwater on the Agulhas bank. There it will remain for the near future, guarded by great white sharks and dangerous currents. We can still test the hypothesis that humans followed the sea by examining sites on the current coast such as PP13B and another site we are excavating called PP5-6. But we can also study locations where the continental shelf drops steeply and the coast was always near—investigations that my colleagues and I are currently initiating.

The genetic, fossil and archaeological records are reasonably concordant in suggesting that the first substantial and prolonged wave of modern human migration out of Africa occurred around 50,000 years ago. But questions about the events leading up to that exodus remain. We still do not know, for example, whether at the end of glacial stage 6 there was just one population of *H. sapiens* left in Africa or whether there were several, with just one ultimately giving rise to everyone alive today. Such unknowns are providing my team and others with a very clear and exciting research direction for the foreseeable future: our fieldwork needs to target the other potential progenitor zones in Africa during that glacial period and expand our knowledge of the climate conditions just before that stage. We need to flesh out the story of these people who eventually pushed out of their refuge, filled up the African continent and went on to conquer the world. ■

PER-ANDERS PETTERSON

PROBING THE PAST

Continued excavation of PP13B (*above*) and other caves along the southern coast of Africa should reveal more about the progenitor population of humans who survived the bottleneck and colonized the globe.

MORE TO EXPLORE

The Origin of Modern Human Behavior: Critique of the Models and Their Test Implications. Christopher S. Henshilwood and Curtis W. Marean in *Current Anthropology,* Vol. 44, No. 5, pages 627–651; December 2003.

Early Human Use of Marine Resources and Pigment in South Africa during the Middle Pleistocene. Curtis W. Marean et al. in *Nature,* Vol. 449, pages 905–908; October 18, 2007.

Fire as an Engineering Tool of Early Modern Humans. Kyle S. Brown et al. in *Science,* Vol. 325, pages 859–862; August 14, 2009.

Coastal South Africa and the Coevolution of the Modern Human Lineage and the Coastal Adaptation. Curtis W. Marean in *Trekking the Shore: Changing Coastlines and the Antiquity of Coastal Settlement.* Edited by Nuno Bicho, Jonathan A. Haws and Loren G. Davis. Springer (in press).

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

© 2010 Scientific American



Robot Pills

A voyage through the human body is no longer mere fantasy. Tiny devices may soon perform surgery, administer drugs and help diagnose disease

BY PAOLO DARIO AND ARIANNA MENCIASSI

© 2010 Scientific American

THE MOVIE *FANTASTIC VOYAGE*, the story of a miniaturized team of doctors traveling through blood vessels to make lifesaving repairs in a patient's brain, was pure science fiction when it came out in 1966. By the time Hollywood remade the film in 1987 as *Innerspace,* a comedy, real-world engineers had already begun building prototypes of pill-size robots that could voyage through a patient's gastrointestinal tract on a doctor's behalf. Patients began swallowing the first commercially built pill cameras in 2000, and since then doctors have used the capsules to get unprecedented views of places, such as the inner folds of the small intestine, that are otherwise difficult to reach without surgery.

One important aspect of *Fantastic Voyage* that has remained fantasy is the notion that such tiny pill cameras could maneuver under their own power, swimming toward a tumor to get a biopsy, checking out inflammation in the small intestine, or even administering drug treatments to an ulcer. In recent years, however, researchers have made great strides in converting the basic elements of a passive camera pill into an active miniature robot. Advanced prototypes, now being tested in animals, have legs, propellers, sophisticated imaging lenses and wireless guidance systems. Soon these tiny robots may be ready for clinical trials. Right now they are testing the limits of miniaturized robotics.

TRANSFORMING PASSIVE PILLS

THE DIGESTIVE TRACT is the first frontier. The first wireless camera pill, M2A, introduced in 1999 by the Israeli company Given Imaging, and subsequent models established the usefulness of examining the gastrointestinal tract with a wireless device. The practice, known as capsule endoscopy, is now routinely used in medicine. Unfortunately, the lack of human control in a passive camera pill leads to a high rate of false negative results—the cameras miss problem areas, which is unacceptable for a diagnostic tool. If the purpose of peering inside the body is to screen for disease or to get a closer look at a suspected problem, a doctor wants most of all to be able to stop the camera and maneuver it to inspect a region of interest.

Transforming a passive capsule into a more reliable device for gastrointestinal screening requires adding moving appendages, or actuators, to propel the pill through the body or act as tools for manipulating tissues. Operating those moving parts demands two-way high-speed wireless data transmission of images and instructions. The pills must, in effect, become tiny robots able to respond quickly to a technician's orders. All these components need sufficient power to complete their tasks during a journey that could last up to 12 hours. And all this must fit into a two-cubic-centimeter container—about the size of a Gummi Bear—that a patient can comfortably swallow.

The same year M2A debuted, the Intelligent Microsystem Center (IMC) in Seoul, Korea, initiated a 10-year project to develop a new generation of capsular endoscopes with advanced features. Such a robotic pill would have onboard sensors and a light source for imaging. It would have mechanisms for delivering drug therapies and taking biopsies. And it would have the ability to locomote, under an endoscopist's wireless remote control. Since 2000, additional companies and research groups have entered the field. For instance, 18 European teams formed a consortium with the IMC to develop capsular robots for cancer detection and treatment. Our group at the Scuola Superiore Sant'Anna in Pisa, Italy, with the medical supervision and guidance of Marc O. Schurr of novineon in Tubingen, Germany, handles the scientific and technical coordination of that project, called VECTOR, for *v*ersatile *e*ndoscopic *c*apsule for gastrointestinal *t*umor *r*ecognition and therapy.

These academic and industry groups have come up with many innovative ideas. In particular, they have posed a variety of solutions to the central challenge: how to control the movement of capsular devices inside the body. Most of them take one of two fundamental approaches.

The first entails directing the movement of the capsule with onboard actuators—moving parts such as paddles, legs, propellers or similar appendages integrated into the shell of the device and capable of deploying once the pill is inside the digestive tract. The actuators, powered by miniature motors, are most often used to direct the capsule's movements, but in some designs legs can also push aside tissue around the capsule, to get a better look at something or to help the capsule pass through a collapsed section of intestine. Motors and actuator mechanisms such as gears are pretty large compared with the total volume of a swallowable capsule, which makes the integration of other essential parts—the imaging sensor or a therapeutic module such as a biopsy tool— challenging. In addition, to distend tissue, a capsule must exert a significant force—equivalent to 10 or 20 times its weight. The ef-

KEY CONCEPTS

- Pill cameras made possible unprecedented internal views of the entire digestive tract, but the uses and accuracy of those passive capsules are limited.
- Active pill-size robotic capsules are being developed for use in screening, diagnosis and therapeutic procedures.
- Miniaturizing robotic components to perform tasks inside the body poses novel engineering challenges. Those challenges are giving rise to creative solutions that will influence robotics and other medical technologies in general.

—The Editors

FRANK HÜLSBÖMER

© 2010 Scientific American

MINI BOTS FOR A WIDE RANGE OF JOBS

To make miniature robots that can operate in the digestive tract, engineers must find ways of wirelessly controlling their locomotion and fine movements. And they must fit the required tools, imaging sensors and power supply into a capsule small enough for a patient to swallow. Here are some examples of the diverse tasks engineers want tiny robots to do and the ways they are trying to overcome the technical challenges.



[THE AUTHORS]

PAOLO DARIO and ARIANNA MENCIASSI are professors of biomedical robotics at the Scuola Superiore Sant'Anna in Pisa, Italy. Dario, inventor of the first self-propelled colonoscopic robot in the 1990s, has also pioneered wireless robotic capsular endoscopes through work with South Korea's Intelligent Microsystems Center and partnerships with European robotics investigators. Menciassi, who has collaborated with Dario for 10 years, specializes in microengineering for minimally invasive therapies and medical nanotechnology.

fort requires high torques from the motors, which consumes considerable power (approximately half a watt). This power drain puts a strain on battery technology, which limits how long these devices can operate.

To conserve battery power, the best trade-off might be to use actuators only for propulsion and find other ways to distend tissue. Having a patient drink half a liter of clear liquid just before swallowing a propeller-driven capsule, for instance, would distend the stomach for up to 20 minutes before the fluid drains away into the small intestine. In that time, the pill could swim around under wireless control and examine the stomach lining.

The bulk and power requirements of onboard actuators have led some researchers to focus on another approach: applying magnetic fields generated outside the body to remotely control the movement of the capsules. In 2005 Olympus and Siemens introduced a magnetic guidance system for its passive camera pill that causes the pill to rotate. The corkscrew motion generates a light friction that helps the capsule burrow through narrow stretches of the digestive tract, such as the small intestine, according to literature from Siemens.

Although using magnets to guide an endoscopic capsule through the intestines is straightforward, achieving reliable control with magnets alone is extremely difficult. Magnetic fields lose power with distance, and with the irregular geometry of the intestine, sudden changes in field strength can cause the capsule to jump or can entirely sever magnetic control over the pill. In practice, this instability may cause an operator to lose contact with the pill and to be unable to find it again. It is possible to compensate by adding more magnets, which would give greater control and stability, but that might require cumbersome configurations of the electromagnetic coils.

TAILORED HYBRIDS

IN LIGHT OF the limitations of both internal and external approaches to controlling capsule movements, we believe that we need to combine these

FRANK HÜLSBÖMER (*three capsules*); COURTESY OF SIEMENS (*magnetic propulsion capsule*); JOSH MCKIBLE (*illustrations*); COURTESY OF FEDERICA RADICI (*Dario*); COURTESY OF ARIANNA MENCIASSI (*Menciassi*)

© 2010 Scientific American

REMOTE SURGERY

One way to expand the range of tasks that robot pills can perform is to design them for self-assembly. The patient would swallow a dozen or more pills; once inside the stomach, the pills would combine with one another to form one big, powerful robot. Surgeons would operate the device wirelessly. When the surgery was completed, the robot would break apart into capsules, which would pass harmlessly through the digestive tract.

Possible surgical configuration



two methods to find a solution that will be comfortable for the patient and offer reliable diagnostics. External magnetic locomotion is adequate for a rough positioning of a capsule inside the intestine; leglike actuators are useful to shift position or maneuver for a better view.

Our research group has designed such a hybrid capsule with four motor-driven legs and tested it in a pig, whose intestines are the same dimensions as those of humans. The capsule's legs remain closed while the device is being swallowed and during most of its trip through the digestive tract. An external magnetic field generator held close to the abdomen guides the capsule forward. When the capsule reaches a segment of intestine whose walls have collapsed, it lifts the surrounding tissue by extending its legs, which move the capsule slightly forward through the opening.

In most areas of the small and large intestines, a hybrid locomotion system would provide the control doctors need for thorough visual inspection. Different situations will call for different solutions. The VECTOR project, for instance, has developed three capsule concepts for the small intestine alone: one is a passive camera pill for normal screening; a second is a diagnostic capsule with active locomotion and spectroscopic imaging that can detect abnormalities underneath the tissue surface. The same spectroscopic sensor is incorporated in the third planned VECTOR capsule, which would also carry a biopsy tool that could take a tissue sample and store it inside the capsule, to be retrieved when the capsule exits the patient.

The ability to perform biopsies and other more complex therapeutic actions, such as surgical procedures, would make capsular endoscopic robots an even more powerful medical tool. But critical problems such as power supply, space constraints and limited torque make many more ambitious therapeutic tasks requiring complicated motions and multiple actuators impossible to achieve with a single two-cubic-centimeter pill.

For these reasons, we are now working on an advanced concept: surgical robots that configure themselves inside the body. Here is how it might work. The patient would drink a stomach-distending fluid, then swallow as many as 10 to 15 capsules. Each capsule would be a miniaturized component with magnets at either end. Once inside the stomach, the capsules, under remote guidance, would quickly assemble themselves into the desired configuration. A surgeon would then use the assembled robot as a wireless tool that can operate without the need to make a single incision from outside the body. When the surgery is done, the magnetic bonds between the capsules could be reconfigured or broken, allowing the parts to make their exit harmlessly through the digestive tract.

We have an early prototype based on two-centimeter capsules with customizable internal parts and actuators. One or more capsules could have a camera, others could have onboard tools, and all would be controlled wirelessly.

Miniaturized robotic components may eventually find wider use throughout the body for a variety of purposes. Guidance systems and camera sensors developed for capsule endoscopy are already influencing related biomedical technologies, such as the newest versions of traditional endoscopes and laparoscopic surgery tools. Beyond health care, these technologies are part of a broad trend toward miniaturization and wirelessly controlled multifunctional robotics. Capsule robots will undoubtedly have an influence on robotic machines in the larger world. ■

HYBRID CAPSULE guided by external magnets navigates the colon of a pig, using extendable legs to adjust its position and push aside tissue.

MORE TO EXPLORE

Wireless Capsule Endoscopy: From Diagnostic Devices to Multipurpose Robotic Systems. Andrea Moglia, Arianna Menciassi, Marc Oliver Schurr and Paolo Dario in *Biomedical Microdevices,* Vol. 9, No. 2, pages 235–243; December 12, 2006.

Wireless Therapeutic Endoscopic Capsule: In Vivo Experiment. P. Valdastri et al. in *Endoscopy 2008,* Vol. 40, No. 12, pages 979–982; December 2008.

Wireless Reconfigurable Modules for Robotic Endoluminal Surgery. Kanako Harada, Ekawahyu Susilo, Arianna Menciassi and Paolo Dario in *Proceedings of the 2009 IEEE International Conference on Robotics and Automation.* Kobe, Japan, May 2009.

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

FRANK HÜLSBÖMER (*left*); COURTESY OF ARIANNA MENCIASSI (*right*)

© 2010 Scientific American

Carbon dioxide emissions are making the oceans more acidic, imperiling the growth and reproduction of species from plankton to squid

BY MARAH J. HARDT AND CARL SAFINA

Threatening Ocean Life

PHOTOGRAPH BY JAMIE CHUNG

© 2010 Scientific American

from the Inside Out

KEY CONCEPTS

- The pH of seawater worldwide is dropping (acidifying) as oceans absorb ever more carbon dioxide from the atmosphere.
- Experiments show that the struggle by copepods, snails, sea urchins and brittlestars to balance the changing pH inside their bodies impairs their ability to reproduce and grow. Many species are unlikely to genetically adapt to ocean acidification, because the change is occurring too quickly.
- As species wither, the marine food chain could be disrupted; human action is needed to curtail further acidification.

—The Editors

© 2010 Scientific American

"SLOW SPERM ... now that's a problem," said Jonathan Havenhand, his British accent compounding the gravity of the message. "That means fewer fertilized eggs, fewer babies and smaller populations." We were sharing a hilly cab ride along the glistening northern coast of Spain to attend an international symposium about the effects of climate change and excess atmospheric carbon dioxide on the world's oceans. As researchers, we were concerned about the underappreciated effects of changing ocean chemistry on the cells, tissues and organs of marine species. In laboratory experiments at the University of Gothenburg in Sweden, Havenhand had demonstrated that such changes could seriously impede the most fundamental strategy of survival: sex.

Ocean acidification—a result of too much carbon dioxide reacting with seawater to form carbonic acid—has been dubbed "the other CO_2 problem." As the water becomes more acidic, corals and animals such as clams and mussels have trouble building their skeletons and shells. But even more sinister, the acidity can interfere with basic bodily functions for all marine animals, shelled or not. By disrupting processes as fundamental as growth and reproduction, ocean acidification threatens the animals' health and even the survival of species. Time is running out to limit acidification before it irreparably harms the food chain on which the world's oceans—and people—depend.

RAPID SEA CHANGE

THE OCEAN'S INTERACTION with CO_2 mitigates some climate effects of the gas. The atmospheric CO_2 concentration is almost 390 parts per million (ppm), but it would be even higher if the oceans didn't soak up 30 million tons of the gas every day. The world's seas have absorbed roughly one third of all CO_2 released by human activities. This "sink" reduces global warming—but at the expense of acidifying the sea. Robert H. Byrne of the University of South Florida has shown that in just the past 15 years, acidity has increased 6 percent in the upper 100 meters of the Pacific Ocean from Hawaii to Alaska. Across the planet, the average pH of the ocean's surface layer has declined 0.12 unit, to approximately 8.1, since the beginning of the industrial revolution.

That change may not sound like much, but because the pH scale is logarithmic, it equates to a 30 percent increase in acidity. Values of pH

pH PRIMER

The pH of a solution is determined by the concentration of hydrogen ions (H^+) in it. A value of 7.0 is neutral; lower is acidic; higher is basic. "Acidification" means a drop in value, anywhere along the scale.



GEORGE RETSECK (*main illustration*)



[BASICS]

ACID IN THE OCEAN

The ocean consists of relatively stable layers created by particular combinations of salinity and temperature (*left*). Sunlight at the top energizes phytoplankton—the base of the food chain—which ultimately nourish all marine life, including copepods that migrate between layers, feeding animals in the deeper water (*center, left*). As CO_2 in the atmosphere rises, though, it increases the acidity of the water (*center, right*)—a phenomenon already being observed (*far right*). Experiments indicate such acidity will harm sea life at various depths.

LAYERS

Atmospheric CO_2 lowers pH in the surface (sunlight) layer directly, and mixing between layers gradually reduces pH farther down, to lesser degrees. Animals in deeper layers are more vulnerable to smaller changes, however.

FOOD CHAIN

Every night a squadron of tiny crustaceans called copepods migrate up and within the surface layer to feed on even tinier zooplankton. They descend by daybreak, serving as food for deeper species. Changing pH could affect their survival, disrupting the food chain.

© 2010 Scientific American

measure hydrogen ions (H^+) in solution. A value of 7.0 is neutral; lower values are increasingly acidic, and higher values are basic. Although 8.1 is mildly basic, the declining trend constitutes acidification. Marine life has not experienced such a rapid shift in millions of years. And paleontology studies show that comparable changes in the past were linked to widespread loss of sea life. It appears that massive volcanic eruptions and methane releases around 250 million years ago may have as much as doubled atmospheric CO_2, leading to the largest mass extinction ever. More than 90 percent of all marine species vanished. A completely different ocean persisted for four million to five million years, which contained relatively few species.

If we continue to emit greenhouse gases at current rates, scientists estimate that atmospheric CO_2 will reach 500 ppm by 2050 and 800 ppm by 2100. The pH of the upper ocean could drop to 7.8 or 7.7—as much as a 150 percent increase in acidity compared with preindustrial times.

Most people envision the ocean as a giant pool of water. But the ocean is more like a layer cake, with each layer created by unique combinations of salinity and temperature. The warmest and freshest (least salty) floats from the surface down 50 to 200 meters, sometimes deeper. Plentiful oxygen and sunlight support the blooming base of the food chain: single-celled phytoplankton that, like plants, use sunlight to create sugar. The phytoplankton nourish zooplankton—small animals ranging from minuscule shrimplike crustaceans to the larvae of giant fish. Zooplankton are eaten by small fish, which feed bigger animals, and so on.

Winds help to mix the surface and deeper layers, sending oxygen down and bringing nutrients up. But the flux of nutrients between surface and seafloor also occurs through the movement of animals, alive and dead. An extensive class of tiny crustaceans called copepods migrate every night, under the cover of darkness, from middle

Geologic records show that comparable pH changes in the past wiped out almost all sea life.

JEN CHRISTIANSEN (*right panel*); SOURCE: "DIRECT OBSERVATIONS OF BASIN-WIDE ACIDIFICATION OF THE NORTH PACIFIC OCEAN," BY ROBERT H. BYRNE ET AL., IN *GEOPHYSICAL RESEARCH LETTERS*, VOL. 37, L02601; JANUARY 20, 2010



ACID CREATION

Carbon dioxide molecules in the air are absorbed by seawater at the surface. The CO_2 then reacts with the water (H_2O) to form carbonic acid (H_2CO_3). Most of this weak acid dissociates into hydrogen ions (H^+) and bicarbonate ions (HCO_3^-). Some bicarbonate also dissociates, producing more H^+. The increase in H^+ ions reduces pH.

GREATER ACIDITY OVER TIME

In just 15 years, pH fell by about 6 percent in the Pacific Ocean surface layer (*above*) along a line from Hawaii to Alaska, as measured during expeditions by the University of South Florida. Oddly, pH is normally highest at the surface (*left*) because various ions mix there. It drops with depth until about 3,000 meters; beyond that, pH remains nearly constant.

© 2010 Scientific American

Emissions targets should limit pH declines to no more than 0.1 over the next century, given the consequences.

and even deep layers to the surface to dine on the banquet created by the day's rays. Many fish and squid follow their movements, while deep dwellers wait for that bountiful food to rain down, in the form of sinking remains. As organisms rise and fall, they pass through waters with different pH values. But as acidification changes this pH profile, it could harm the organisms.

THE INSIDE ANGLE

AT THE SCALE of individual marine animals, acidification can force creatures to spend more energy on restoring and maintaining their internal pH balance, diverting energy away from important processes such as growth and reproduction.

Even small increases in seawater CO_2 concentration can cause rapid diffusion into the bodies of water-breathing animals. Once inside, CO_2 reacts with internal fluids, creating hydrogen ions, making the bodily fluids or tissue more acidic. Species employ various mechanisms to balance their internal pH. These actions include producing negative ions such as bicarbonate that soak up, or buffer, the extra hydrogen ions; pumping ions in and out of cells and intercellular spaces; and reducing metabolism to absorb fewer ions and "wait out" the period of high H^+ concentration. But none of these mechanisms is meant to handle a sustained drop in pH. As an organism struggles to regain an acid-base balance, it sacrifices energy. Basic life functions such as synthesizing protein and maintaining a strong immune system can also become compromised.

Most species possess at least some buffer molecules. Fish and other active species stockpile them to reduce temporary pH declines that result from extended swimming bursts. Just like in a runner, muscles shift to anaerobic (nonoxygen based) metabolism during sprints, which uses up ATP (the main fuel molecule) more quickly, causing extra H^+ ions to accumulate. But few species can stockpile enough buffering to last across extended timescales. If small pH changes occurred gradually over tens of thousands of years, a species might evolve adaptations, for example, by retaining chance genetic mutations that result in greater production of buffer molecules. But species generally cannot adapt to changes occurring over mere hundreds of years or less. Similar changes produced in the lab over days to weeks are lethal.

In past eras when CO_2 concentrations rose, species with less well-buffered systems fared poorly. Declines in pH may especially harm deep-sea species, whose stable environment leaves them ill equipped to adapt to change. (For this reason, proposed strategies to combat climate change by pumping large quantities of CO_2 into the deep sea are worrisome; they could destabilize the habitats of a wide array of creatures.)

POOR GROWTH AND REPRODUCTION

THE INTERNAL EFFECTS of ocean acidification vary across different developmental stages of life. A small but growing body of research points to a variety of potential trouble.

Indeed, the very first spark of life—fertilization—can be affected. In the lab, scientists sim-

GEORGE RETSECK (*illustration*); "NEAR-FUTURE LEVEL OF CO_2-DRIVEN OCEAN ACIDIFICATION RADICALLY AFFECTS LARVAL SURVIVAL AND DEVELOPMENT IN THE BRITTLESTAR *OPHIOTHRIX FRAGILIS*," BY SAMUEL DUPONT ET AL., IN *MARINE ECOLOGY PROGRESS SERIES*, VOL. 373; 2008 (*brittlestar*); ROBERT SUTTON *University of Plymouth* (*eggs*)

[DEFENSE MECHANISMS]

THE STRUGGLE TO NEUTRALIZE

More CO_2 in seawater sends more hydrogen ions inside animals' bodies, making tissue more acidic. Species can attempt different countermeasures, but all sap energy. Cells, for example, can try to pump out the extra H^+ ions: energy from an ATP molecule pushes the ions through a one-way channel in the cell membrane (*a*). Or the body can produce more buffer molecules such as carbonate (HCO_3^-) that bind to the H^+ ions and carry them away (*b*).



▶ CRIPPLED CRITTERS

Laboratory experiments show that the development and fertility of ocean animals suffer in more acidic waters. After being raised for eight days in 8.1 pH water, typical of today's oceans, the larvae of brittlestars were mature and symmetric (*left, top*), but larvae raised at 7.7 pH were deformed (*left, bottom*), and none reached full development. Snail eggs fertilized and raised at 8.05 pH all had embryos, which had grown shells after 18 days (*right, top*), but in 7.6 pH water (*right, bottom*) some eggs were empty, and embryos had still not developed shells.

© 2010 Scientific American

ulate acidification by pumping extra CO_2 bubbles through seawater tanks. As Havenhand had explained during our cab ride, sperm of the Australian sea urchin *Heliocidaris erythrogramma* moved 16 percent less and swam 12 percent slower when experimenters lowered seawater pH by 0.4 (within the range predicted by 2100). Fertilization success dropped by 25 percent. In the wild, a 25 percent reduction could lead to significantly diminished adult populations over time. Although individual sea urchins release millions of sperm and eggs, the sperm do not remain viable for very long; they have to find and fertilize an egg within a few minutes. In a big, turbulent ocean, sluggish sperm may never reach their destination at all.

Acidification also thwarts early larval stages of several species. Samuel Dupont, down the hall from Havenhand at Gothenburg, exposed larvae of a temperate brittlestar—a relative of the common sea star—to pH reduced by 0.2 to 0.4 unit. Many showed abnormal development, and fewer than 0.1 percent survived more than eight days. In another study, fewer embryos of the snail *Littorina obtusata* hatched when exposed to lower pH waters, and those that did hatch moved less frequently and more slowly than normal.

A change of 0.2 to 0.4 pH all at once is more dramatic than species in the wild are experiencing, and some species might be able to adapt to gradual change. But for others, the effects of even slight acidification come on strong and fast. Scientists suspect ocean acidification explains recent mortality in larval oysters along the coast of Oregon, for example, sending some oyster growers scrambling to find enough babies to stay in business.

Adult animals suffer as well, especially when it comes to growth. Sea urchins and snails move slowly, but *growing* slowly is problematic. In 2005 researchers at Kyoto University in Japan determined that a CO_2 concentration 200 ppm higher than today's value, pumped into seawater for six months, reduced growth rates for the sea urchin species *Hemicentrotus pulcherrimus* and *Echinometra mathaei* and for the strawberry conch *Strombus luhuanu*. The 200-ppm increase is equal to that predicted over the next four to five decades. Slowed growth leaves individuals smaller for longer, making them more susceptible to predators and potentially reducing their reproductive output.

Acidification also makes it harder for some phytoplankton species to absorb iron, a micronutrient critical for growth. Researchers at Princeton University indicate that a 0.3 pH decline could reduce phytoplankton iron uptake by 10 to 20 percent. In addition to being an important link in the food chain, phytoplankton produce vast amounts of oxygen that we breathe.

In other experiments, the sediment-dwelling brittlestar *Amphiura filiformis* grew arms at greater rates under lower pH but lost significant muscle mass. Strong muscles are required for

MODELS VS. MEASUREMENTS IN THE PACIFIC



Ocean pH predictions by climate models (*top*) parallel measurements taken from 1989 to 2009 at a University of Hawaii recording station in the north-central Pacific Ocean (*bottom*). The correlation indicates that if atmospheric CO_2 continues to rise, pH will drop significantly.

JEN CHRISTIANSEN (*graphs*); SOURCES: "DIRECT EFFECTS OF CO_2 CONCENTRATION ON GROWTH AND ISOTOPIC COMPOSITION OF MARINE PLANKTON," BY DIETER A. WOLF-GLADROW ET AL., IN *TELLUS SERIES B-CHEMICAL AND PHYSICAL METEOROLOGY*, VOL. 51, NO. 2; 2002; AND "PHYSICAL AND BIOGEOCHEMICAL MODULATION OF OCEAN ACIDIFICATION IN THE CENTRAL NORTH PACIFIC," BY JOHN E. DORE ET AL., IN *PNAS*, VOL. 106, NO. 30; JULY 28, 2009



© 2010 Scientific American

[THE AUTHORS]

MARAH J. HARDT, founder of OceanInk, is a research scientist, writer and consultant on the Big Island of Hawaii. A coral reef ecologist by training and a former fellow at the Blue Ocean Institute in Cold Spring Harbor, N.Y., she works to solve diverse issues ranging from global overfishing to freshwater conservation. **CARL SAFINA** is founding president of the Blue Ocean Institute and an adjunct professor at Stony Brook University. A MacArthur Fellow, he is author of *Song for the Blue Ocean: Encounters along the World's Coasts and beneath the Seas* and *Eye of the Albatross: Visions of Hope and Survival*. Safina's upcoming book *The View from Lazy Point: A Natural Year in an Unnatural World* will appear this year.



FUTURE DATA

Carbon dioxide and pH sensors were deployed on buoys in the Pacific Ocean in December 2009 by the Scripps Institution of Oceanography, which provides data to California Current Ecosystem research projects. Such information will improve forecasts of ocean acidification trends.

feeding, building burrows and escaping predators. A pH decline of 0.3 to 0.5 suppressed the immune system response of the common blue mussel within one month. Reduced strength, growth, immune function or reproduction can cause long-term population declines—bad news for the victims, as well as for the many other species (including humans) that rely on them for food and even habitat. Grazing by sea urchins, for example, helps to keep coral reefs and kelp forests healthy, and the mixing of sediments by the brittlestars' movements is critical to making the sediments livable for many other species.

For some creatures, ocean acidification can simply mean the end. When a sample of copepod species common off the California coast (*Paraeuchaeta elongata*) was exposed to water that was 0.2 pH below normal, half of the organisms died within a week. The fish we prefer to eat, from tuna to salmon or striped bass, depend on an abundance of specific copepods to support the prey that supports them.

Several species of fish, such as the spotted wolffish (*Anarhichas minor*), have shown remarkable tolerance in the lab, because they maintain a relatively large stockpile of buffers and store extra oxygen in their tissue, which is handy because H^+ ions interfere with the blood's ability to absorb oxygen from the water. Even very adaptable fish, however, may struggle if their food supply dwindles. Other species are not so well prepared. Highly active squid, for example, have no oxygen stores—they use all they have all the time. Less oxygen in their blood would limit their ability to hunt, avoid predators and find mates. For the commercially important squid *Illex illecebrosus*, a pH drop of just 0.15 could cause significant harm.

The message of lab studies as well as the geologic record is that ocean acidification forces animals to struggle harder, which today they are already doing because of other human-induced stressors such as warming waters, pollution and overfishing.

ACID ADAPTATION?

LAB EXPERIMENTS PERSIST for weeks to months. Climate change occurs over decades and centuries. Some species could adapt, especially if they have a short reproductive cycle. Each time an animal reproduces, genetic mutations can arise in the offspring that might help the next generation adjust to new circumstances. Ninety years—the predicted time frame for pH to decline by 0.3 to 0.5 unit—is extremely short, however, for genetic adaptation by species that reproduce at relatively slow rates and that may already be stressed by the 30 percent pH decrease. Species extinctions often result from slow declines over centuries or more; a decline of just 1 percent of individuals per generation could cause extinction in less than a century.

Alarmingly, the pH drop observed so far and the predicted trajectory under current emissions trends are 100 times faster than any changes in prior millennia. Left unchecked, CO_2 levels will create a very different ocean, one never experienced by modern species.

Adaptation is even more unlikely because the effects of acidification, and the other struggles creatures face, interact. For example, increased CO_2 levels can narrow the temperature range in which an individual can survive. We already see such constraints on corals and some algae, which become heat-stressed at lower temperatures than normal if exposed to higher CO_2.

OPTIONS FOR THE FUTURE

SCIENTISTS HAVE CONSISTENTLY underestimated rates of climate change, from Arctic ice melt to sea-level rise. Increasingly, experts recommend limiting atmospheric CO_2 to prevent dangerous levels of global warming. But the targets should be set with ocean acidification in mind as well. Unabated acidification could completely restructure marine ecosystems, with cascading effects across the food chain. Some species might thrive on a new combination of plankton while others suffer, but there is no telling if the species that we depend on most (or like the best) will be the winners. The changes could also hurt tourism and erase potential pharmaceutical and biomedical resources.

Ocean acidification also changes the rules for the planet's entire carbon cycle. Although the oceans now absorb a vast quantity of human emissions, the absorption rate slows as the seawater CO_2 concentration increases, and CO_2 "backs up" at the sea surface. As a result, atmospheric CO_2 concentration will rise even faster, accelerating global weather changes.

Such consequences warrant emissions targets that limit pH declines to no more than 0.1 over the next century. More and more, reducing the atmospheric CO_2 level to 350 ppm seems like the rational target. Stabilizing at 450 ppm by 2100, as some have suggested, could perhaps keep an additional pH decline to 0.1. But even that number could doom coral reefs and make it impossible for some animals to build shells, especially

COURTESY OF MARCY HARDT (*Hardt*); COURTESY OF PATRICIA PALADINES (*Safina*); COURTESY OF UWE SEND'S LAB, HTTP://MOORING.UCSD.EDU (*buoy*)

© 2010 Scientific American

[SENSORY CONFUSION]

THE NEW LSD

Increasingly acidic ocean water interferes with the internal body chemistry of many species. Researchers are also finding that acidification may be altering survival in other unusual ways.

For example, many marine species use subtle olfactory cues to seek prey, mates or suitable habitat. Certain clownfish differentiate between attractive and repulsive smells to decide which reef and anemone to settle on for the rest of their days. In laboratory studies, clownfish larvae raised in seawater with a pH 0.2 to 0.4 lower than today's ocean average swim toward negative cues and do not respond to positive ones. Their nasal organs look normal, but scientists think the acidification may disrupt the way chemical signals travel through the neurological system. It's a new kind of LSD: lost smell disorder. More studies are needed to determine if and how LSD may affect fish communities worldwide.

Complex, pH-regulated interactions between molecules in seawater also increase or decrease sound reception. Ocean acidification pumps up the volume; if pH declines by another 0.3 unit (within the scope predicted by 2100), oceans could be 40 percent louder. Although no studies have directly linked increased noise caused by acidification with changes in animal survival, the findings raise a warning flag, because marine wildlife, especially marine mammals, depend on sound for navigation, communication, hunting and courtship. *—M.J.H. and C.S.*

CLOWNFISH

GEORGETTE DOUWMA *Nature Picture Library*

in the Southern Ocean, which encircles Antarctica. Because of its cold temperatures and unique circulation patterns, the Southern Ocean will start dissolving shell and skeletal structures sooner than other oceans. It is far easier to prevent further acidification than to reverse changes once they occur; natural buffering systems would need hundreds to thousands of years to restore pH to preindustrial levels.

What can be done? For a start, the Obama administration should enact a National Ocean Policy—the first ever for the U.S.—because it could effectively coordinate action to combat these multiple threats. The U.S. Environmental Protection Agency should move forward with including CO_2 as a pollutant under the Clean Water Act, giving states authority to enforce CO_2 emissions limits. Establishing marine protected areas would allow species to recover from overexploitation; higher numbers would give their populations and gene pools more resilience in responding to climate changes. Adjusting fishery catch limits so they meet scientific recommendations rather than political desires would help. And signing the United Nations Convention on the Law of the Sea, which the U.S. has put off for decades, would make the nation a leader in marine stewardship.

More science is needed, too. Funding to support research initiatives by the European Project on Ocean Acidification and to implement the Federal Ocean Acidification Research and Monitoring Act will deepen understanding of acidification's effects. But a dramatically scaled-up monitoring network to detect acidification is also required. An international team, led by Richard Feely of the Pacific Marine Environmental Laboratory in Seattle and Victoria J. Fabry of California State University, San Marcos, has created a blueprint for integrating acidification monitoring into existing ocean tracking programs, such as OceanSITES, and the recommendations should be followed as soon as possible. In addition, expanding efforts to combine field data with laboratory experiments, such as the California Current Ecosystem Interdisciplinary Biogeochemical Moorings project, will ensure that scientists' experiments simulate realistic conditions.

Ultimately, the solution to ocean acidification lies in a new energy economy. In light of recent lethal coal mine and offshore drilling explosions and the catastrophic Gulf of Mexico oil spill, the U.S. has more reason than ever to forge a safer energy strategy for the planet. Only a dramatic reduction in fossil fuel use can prevent further CO_2 emissions from contaminating the seas. An explicit plan to shift from finite, dangerous energy sources to renewable, clean energy sources offers nations a more secure path forward. And it offers the planet, especially the oceans, a chance for a healthy future. ■

MORE TO EXPLORE

The Dangers of Ocean Acidification. Scott C. Doney in *Scientific American,* Vol. 294, No. 3, pages 58–65; March 2006.

Impacts of Ocean Acidification on Marine Fauna and Ecosystem Processes. Victoria J. Fabry et al. in *ICES Journal of Marine Science,* Vol. 65, No. 3, pages 414–432; April 2008.

European Project on OCean Acidification: **www.epoca-project.eu**

For more research and policy considerations, see the Ocean Acidification Network (**www.oceanacidification.net**), notably the Monaco Declaration, and the Research Priorities Report of the Ocean in a High CO_2 World II Symposium.

U.S. Ocean Carbon and Biogeochemistry program FAQ: **www.whoi.edu/OCB-OA/FAQs**

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

© 2010 Scientific American



© 2010 Scientific American

BRYAN CHRISTIE DESIGN

FILMING THE INVISIBLE IN 4-D

Picture this: a movie revealing the inner workings of a cell or showing a nanomachine in action. A new microscopy is making such imaging possible

BY AHMED H. ZEWAIL

THE HUMAN EYE is limited in its vision. We cannot see objects much thinner than a human hair (a fraction of a millimeter) or resolve motions quicker than a blink (a tenth of a second). Advances in optics and microscopy over the past millennium have, of course, let us peer far beyond the limits of the naked eye, to view exquisite images such as a micrograph of a virus or a stroboscopic photograph of a bullet at the millisecond it punched through a lightbulb. But if we were shown a movie depicting atoms jiggling around, until recently we could be reasonably sure we were looking at a cartoon, an artist's impression or a simulation of some sort.

In the past 10 years my research group at the California Institute of Technology has developed a new form of imaging, unveiling motions that occur at the size scale of atoms and over time intervals as short as a femtosecond (a million billionth of a second). Because the technique enables imaging in both space and time and is based on the venerable electron microscope, I dubbed it four-dimensional (4-D) electron microscopy. We have used it to visualize phenomena such as the vibration of cantilevers a few billionths of a meter wide, the motion of sheets of carbon atoms in graphite vibrating like a drum after being "struck" by a laser pulse, and the transformation of matter from one state to another. We have also imaged individual proteins and cells.

Four-dimensional electron microscopy promises to answer questions in fields ranging from materials science to biology: how to understand the behavior of materials from the bottom up, from the atomic to macroscopic scale; how nanoscale or microscale machines (NEMS and MEMS) function; and how

KEY CONCEPTS

- Four-dimensional electron microscopy produces "movies" of nanoscale processes occurring over time intervals as short as femtoseconds (10^{-15} second).
- The technique builds up each frame of the movie from thousands of individual shots taken at precisely defined times.
- It has applications in a wide range of fields, including materials science, nanotechnology and medicine.

—The Editors

© 2010 Scientific American

proteins or assemblies of biological molecules fold and become organized into larger structures, a vital process in the functioning of all living cells. Four-dimensional microscopy can also reveal the atomic arrangements of nanoscale structures (which determine the properties of new nanomaterials), and, potentially, track electrons moving around in atoms and molecules on the timescale of attoseconds (a billion billionth of a second). Along with the advances in basic science, the potential applications are wide-ranging, including the design of nanomachines and new kinds of medicines.

CATS AND ATOMS IN MOTION

ALTHOUGH 4-D MICROSCOPY is a cutting-edge technique that relies on advanced lasers and concepts from quantum physics, many of its principles can be understood by considering how scientists developed stop-motion photography more than a century ago. In particular, in the 1890s, Étienne-Jules Marey, a professor at the Collège de France, studied fast motions by placing a rotating disk with slits in it between the moving object and a photographic plate or strip, producing a series of exposures similar to modern motion picture filming.

Among other studies, Marey investigated how a falling cat rights itself so that it lands on its feet. With nothing but air to push on, how did cats instinctively perform this acrobatic feat without violating Newton's laws of motion? The fall and the flurry of legs took less than a second—too fast for the unaided eye to see precisely what happened. Marey's stop-motion snapshots provided the answer, which involves twisting the hindquarters and forequarters in opposite directions with legs extended and retracted. High divers, dancers and astronauts learn similar motions to turn themselves.

Another approach, stroboscopic photography, relied on short light flashes to capture events occurring on much shorter timescales than is possible with mechanical shutters. The flashes make an object moving in the dark momentarily visible to a detector such as an observer's eye or a photographic plate. In the mid-20th century Harold Edgerton of the Massachusetts Institute of Technology greatly advanced stroboscopic photography by developing electronics that could produce reliable, repetitive, microsecond flashes of light.

The falling-cat experiment requires shutter times or stroboscopic flashes short enough for the photographs to show the animal clearly despite its motion. Suppose the cat has righted itself

[HOW IT WORKS]

THE FOUR-DIMENSIONAL ELECTRON MICROSCOPE

A standard electron microscope records still images of a nanoscopic sample by sending a beam of electrons through the sample and focusing it onto a detector. By employing single-electron pulses, a four-dimensional electron microscope produces movie frames representing time steps as short as femtoseconds (10^{-15} second).



❶ A clocking pulse—such as a femtosecond laser pulse—excites the specimen to set the process of interest in motion at a precisely defined time zero.

❷ An electron-generating laser pulse is created in synchrony with the clocking pulse, but it is then delayed by a controlled amount.

❸ A pulse containing a single electron passes through the sample at a precise time T after time zero.

❹ Magnetic lenses "focus" the electron onto a charge-coupled device, which registers it as a single pixel to include in the T frame of the nanomovie.



▲ Each frame of the nanomovie is built up by repeating this process thousands of times with the same delay and combining all the pixels from the individual shots. Researchers may also use the microscope in other modes, such as with one many-electron pulse per frame, depending on the kind of movie to be obtained. The single-electron mode produces the finest spatial resolution and captures the shortest time spans in each frame.

© 2010 Scientific American

half a second after being released. At that instant the cat will be falling at five meters per second, so by using one-millisecond flashes we will ensure that the cat falls no more than five millimeters during each exposure so that the image of the cat will be only slightly blurred by its motion. To slice the acrobatics into 10 snapshots, the photographs must be taken every 50 milliseconds.

If we wish to observe the behavior of a molecule instead of a feline, how fast must our stroboscopic flashes be? Many changes in molecular or material structure involve atoms moving a few angstroms (one angstrom equals 10^{-10} meter). To map out such motion requires a spatial resolution of less than one angstrom. Atoms often move at speeds of about one kilometer per second in these transformations, requiring stroboscopic flashes no longer than 10 femtoseconds to observe them with better than 0.1-angstrom definition. As long ago as the 1980s researchers used femtosecond laser pulses to time chemical processes involving moving atoms, but without imaging the positions of the atoms in space—the wavelength of the light is hundreds of times longer than the spacing between atoms in molecules or materials [see "The Birth of Molecules," by Ahmed H. Zewail; Scientific American, December 1990].

Accelerated electrons have long produced images at atomic scales—as in electron microscopes—but only with targets fixed in place and imaged over time intervals of milliseconds or longer, being limited by the speed of the camera. The atom-scale movies we sought thus required the spatial resolution of an electron microscope but with femtosecond electron pulses to "illuminate" the targets. The illuminating packets of electrons are called probe pulses.

Another issue is clocking of the motion—having a well-defined instant in time when the motion begins. We will not get useful images if all the probe pulses take snapshots before the motion starts or after it finishes. In photographing the cat, the recording begins when the cat is released. For ultrafast recording, a femtosecond initiation pulse called the clocking pulse launches the material or the process to be studied.

Even with probing and clocking under control, the issue of synchronization remains. Here the typical ultrafast experiment drastically departs from the cat analogy. Marey could complete his experiment by dropping one cat once, if everything went according to plan. And it did not matter much if the series of exposures began, say, five, 10 or 17 milliseconds after the cat's release. Ultrafast microscopy, however, may probe millions of atoms or molecules for each clocking pulse or may build up images by repeating an experiment thousands of times. Imagine if Marey had been restricted to capturing only a narrow vertical strip of the field of view with each cat drop. To build up the series of full snapshots of the falling cat, he would have had to repeat the experiment many times, recording along a slightly different vertical strip each time. For the various strips to combine sensibly and form a meaningful whole image, he would need to prepare the cat in the same starting configuration for each drop and carefully synchronize the release with the shutter openings in the same way each time. (The technique would also rely on the cat moving in the same fashion every time. I suspect molecules are more reliable than cats in that respect.)

The starting configurations must be accurate to a small fraction of the cat's size, and the time synchronization must be accurate to less than the shutter durations. Similarly, in ultrafast imaging of atoms or molecules, the launch configuration must be defined to subangstrom resolution, and the relative timing of clocking and probe pulses must be of femtosecond precision. The timing of probe pulses relative to the clocking is accomplished by sending either of these pulses along a path with an adjustable length. For a pulse traveling at the speed of light, setting the path length to an accuracy of one micron corresponds to setting the relative timing with 3.3-femtosecond accuracy.

A further major and fundamental problem remained to be overcome before we could make movies with electrons. Unlike photons, electrons are charged and repel one another. Crowding a lot of them into a pulse spoils both the temporal and spatial resolutions because the electrons' mutual repulsion blows the pulse apart. In the 1980s Oleg Bostanjoglo of the Technical University of Berlin did achieve imaging using pulses having as few as 100 million electrons, but the resolutions were no better than nanoseconds and microns (later significantly improved to the submicron level by researchers at Lawrence Livermore National Laboratory).

My group attacked this challenge by developing single-electron imaging, which built on our earlier work with ultrafast electron diffraction. Each probe pulse contains a single electron and thus provides only a single "speck of light" in the final movie. Yet thanks to each pulse's careful timing and another property known as the coherence of the pulse, the many specks add up to form a useful image of the object. A similar feat



▲ A 50-nanometer-wide cantilever made of a nickel-titanium alloy oscillates after a laser pulse excites it. Blue boxes highlight the movement. The full movie (online at **ScientificAmerican.com/aug2010/nanomovies**) has one frame every 10 nanoseconds. Material properties determined from these oscillations would influence the design of nanomechanical devices.

GEORGE RETSECK (*illustrations*); COURTESY OF CALIFORNIA INSTITUTE OF TECHNOLOGY (*movie frames*)

© 2010 Scientific American

[CASE STUDY]

A NANOSCOPIC ROSETTA STONE

Four-dimensional microscopy of graphite nanocrystals, some only a few atomic layers thick, demonstrated three different imaging modes, producing data about the material in a variety of "languages." The studies examined how the nanocrystals responded when a laser pulse slammed into them from above.



❶ Diffraction patterns revealed the motion of each crystal's atomic layers as they were pushed together and then rebounded in the picoseconds after the laser struck at time zero and as they subsequently oscillated up and down for hundreds of picoseconds.



❷ Images of the nanocrystal measured these oscillations as they proceeded at different locations. Over tens of microseconds, the crystal's initially chaotic motion Ⓐ (*suggested here by arrows*) evolved into a coordinated drumming motion of the entire crystal Ⓑ.



❸ Measurements of the energy lost by imaging electrons in collisions (*depicted here*) with the graphite's electrons indicated how the carbon bonds in the material became more like the bonds in diamond during compression of the layers and more like the bonds in graphene (an isolated layer of carbon atoms) during expansion.





▲ Movie frames show the graphite nanocrystal oscillating like a drumhead after being struck by a laser pulse. The frames show a 24-micron-wide area at 250-nanosecond intervals (every fifth frame of the movie). Subtle permanent buckling of the graphite surface produces the dark bands, which move when the surface ripples. Red boxes have been added to guide the eye. The movie is online at **ScientificAmerican.com/aug2010/nanomovies**

GEORGE RETSECK (*illustrations*); COURTESY OF CALIFORNIA INSTITUTE OF TECHNOLOGY (*movie frames*)

© 2010 Scientific American

is sometimes exhibited as one of the characteristic oddities of quantum mechanics: electrons pass through two slits one at a time, each one contributing a single speck at some random location on a detection screen. Yet all the specks add up to form predictable patterns of light and darkness characteristic of interfering waves.

Single-electron imaging was the key to 4-D ultrafast electron microscopy (UEM). We could now make movies of molecules and materials as they responded to various situations, like so many startled cats twisting in the air.

DECIPHERING NANOMATTER

ONE OF OUR FIRST TARGETS was graphite, the "lead" material in pencils. We chose graphite in part because it is an unusual material, with applications in environments as extreme as those in nuclear reactor cores, and because it has close relatives that are just as remarkable. Graphite consists of carbon atoms arranged in a hexagonal pattern to form sheets reminiscent of chicken wire. Relatively weak bonds hold the sheets together in a stack. Writing with an ordinary pencil relies on pieces of the graphite sloughing off and adhering to the paper. The pencil marks include tiny quantities of the strongest material known to science—graphene, which consists of isolated single sheets of carbon atoms. Researchers are studying graphene vigorously for a variety of electronics applications. Furthermore, when soft graphite is subjected to extreme pressure, its atoms rearrange to form diamond, one of the hardest known substances.

To study graphite's response to mechanical shocks, we took nanoscale crystals of the substance—some only nanometers thick, or a few sheets of atoms—and struck them with intense femtosecond laser pulses, which served as the clocking pulses for our microscope. Each laser pulse pushed the graphite's layers of atoms momentarily closer together, setting them oscillating up and down [*see box on opposite page*]. Our electron microscope sent its electrons through these oscillating graphite layers to produce two kinds of picture: a real-space image (much like a photograph of the graphite surface) or a diffraction pattern, which is a regular array of spots whose precise configuration provides information about the arrangement and separations of atoms in the graphite lattice. In particular, we could track the layers oscillating up and down by the movements of the spots in the diffraction pattern. The oscillations had frequencies of about 10 to 100 gigahertz (10^{10} to 10^{11} cycles per second). No imaging experiment had previously observed such high-frequency resonances unfolding over time.

From our measurements we determined the elasticity of graphite perpendicular to the planes of atoms—how the material responds to compressing or stretching forces acting in that direction. Imagine that the graphite crystal is a stack of rigid metal plates connected by springs and that the laser pulse is a large sledgehammer striking the top plate. We measured the properties of the springs.

The metal-plate analogy is reasonable as long as our "camera" is zoomed in very close. If the camera figuratively "pulls back," however, more of the tiny graphite crystal comes into view. Now the hammer is striking one region of the top metal sheet, and it becomes apparent that the sheets are flexing, with the compression and expansion propagating out from the impact point in waves.

When we pull back the camera even farther and take images more slowly, yet another kind of dynamics comes into view. Now we see how the laser pulse sets the entire nanoscopically thin crystal oscillating, like a drumhead hit by a drumstick. We saw that in the first few microseconds after the laser pulse hit, the crystal's motion appeared chaotic, but as time went on the entire crystal settled down into a well-defined resonant oscillation—it drummed!

For these oscillations, the material property that sets the resonance frequency is the elasticity of the graphite planes—their response to being stretched or compressed in the plane. We found that the graphite is much more resistant to being deformed in the planes of carbon atoms than it is to having those planes pulled apart or pushed together. The results can be explained by considering that the chemical bonds joining the carbon atoms in each hexagonal layer are much stronger than the bonds linking adjacent planes to one another.

Although studies of bulk samples of graphite produce similar data about graphite's elasticity, the information we obtained tells us much more. It addresses questions of two types that are fundamental to our understanding of how materials behave at the nanoscale: first, at what length scale does the description of a substance in terms of a continuum material with properties such as elasticity break down? Second, can we extrapolate from the behavior at atomic scales of length and time to reproduce the known macroscopic properties of a material? With graphite, we found that even quite nanoscopic samples (only

By integrating the fourth dimension, we are turning still pictures into the movies needed to watch matter's behavior—from atoms to cells—unfolding in time.

[THE AUTHOR]

AHMED H. ZEWAIL received the 1999 Nobel Prize in Chemistry for his studies of the transition states of chemical reactions using femtosecond spectroscopy. He is Linus Pauling Chair professor of chemistry, director of the Physical Biology Center for Ultrafast Science and Technology, and professor of physics, all at the California Institute of Technology. In 2009 he was appointed to the President's Council of Advisors on Science and Technology and was named the first U.S. Science Envoy to the Middle East.

COURTESY OF AZ FOUNDATION

© 2010 Scientific American

This technique has produced images of bacterial cell membranes and protein vesicles with femtosecond- and nanometer-scale resolutions.



▲ An *Escherichia coli* bacterium was imaged with photon-induced near-field electron microscopy. A femtosecond laser pulse generated an evanescent electromagnetic field in the cell's membrane at time zero. By collecting only the imaging electrons that gained energy from this field, the technique produces high-contrast, high-spatial resolution of the membrane (*top*). The false-color contour plot depicts the intensity recorded. The method can capture events occurring on very short timescales, as is evinced by the field's significant decay after 200 femtoseconds (*middle*). The field vanishes by 2,000 femtoseconds (*bottom*).

a few dozen atomic layers thick) behave surprisingly like the bulk material. Would this description still be valid near the graphene limit?

The movies of graphite I have described thus far all relied on collisions of our probe electrons with the sample in which they lose no energy—like rubber balls bouncing off something hard. Sometimes, however, a probe electron may lose energy, by exciting an electron in a carbon atom. The amount of energy lost depends on the kind of bond in which the atom's electron was involved. A very old technique called electron energy loss spectroscopy can measure such losses; the energy spectra obtained provide information about the bonding in a material and the chemical elements that compose it. Using this method with our ultrafast electron microscope, we showed that during the compression phase, the bonding inside the graphite shifted toward the kind of bond that is characteristic of diamond. In the expansion phase, the bonding of the surface atoms shifted toward that of graphene. Conventional electron energy loss spectroscopy is far too slow to observe these changes.

FROM CANTILEVERS TO CELLS

MY GROUP HAS NOW carried out four-dimensional microscopy on a number of materials in addition to graphite. In iron, we made diffraction images to follow the crystal structure changing from what is called body-centered cubic to face-centered cubic, a process that occurs in many industrial applications at high temperatures, including production of steel. We saw two dynamic processes unfold when we heated the iron from room temperature to nearly 1,500 kelvins in about a nanosecond. First, specks of the face-centered phase developed, or nucleated, at locations in the crystal relatively slowly—on a nanosecond timescale—out of the incoherent motions of iron atoms. Second, these regions of the new phase grew at the speed of sound, meaning that the process took only picoseconds (10^{-12} second) to encompass the hot iron. This rapidly spreading transformation involves numerous atoms being displaced in a coordinated fashion, a curious kind of "emergence" of a large-scale change in the crystal from the innumerable underlying nanoscopic motions. Understanding of this phenomenon might lead to better ways to handle iron and steel (and many other materials) in industrial processes.

One of the most powerful applications of 4-D ultrafast electron microscopy is seeing nanosystems and microsystems as they function in real time. For instance, we imaged the resonant oscillations of nanoscopic cantilevers, which had not been accomplished before for such high-frequency motions. From our results we determined a range of quantities that describe the cantilevers' material properties and their motion, and we saw that they functioned coherently for nearly 10^{11} oscillations. Researchers can use such data to test the theoretical models that guide design of microelectromechanical and nanoelectromechanical systems, which in turn may lead to new kinds of such devices or new uses for them.

Four-dimensional imaging with ultrafast electron microscopy also has potential biological applications. To fully understand how the body functions, investigators need to know not only the structures of the various proteins and other molecular and cellular structures involved but also their dynamics—how a protein folds, how it selectively recognizes other molecules, what role the water around it plays, and so on. Some biological functions involve ultrafast steps. For instance, our vision and photosynthesis in plants both rely on photons of light triggering femtosecond-scale processes. Although many proteins function, and malfunction, on timescales much longer than femtoseconds, the atomic and molecular motions in the initial femtoseconds can determine whether these macromolecules ultimately fold properly into a useful structure or into one that, say, causes Alzheimer's disease.

One study on protein folding illustrates the kind of techniques needed and the results that are possible. My colleagues and I investigated how quickly a short length of protein would fold into one turn of a helix by heating the water in which the protein was immersed—a so-called ultrafast temperature jump. (Helices occur in innumerable proteins.) We found that short helices formed more than 1,000 times faster than researchers have thought—arising in hundreds of picoseconds to a few nanoseconds rather than the microseconds commonly believed. Knowing that such rapid folding occurs may lead to new understanding of biochemical processes, including those involved in diseases.

Biological imaging with our 4-D ultrafast technology often relies on a well-established technique called cryoelectron microscopy, in which a sample in water is plunged quickly into liquid ethane (which boils at –89 degrees Celsius). The water freezes into a glassy solid that does not diffract electrons and spoil imaging (and the sample itself!) as ordinary ice crystals do. We have obtained images of bacterial cells and

COURTESY OF CALIFORNIA INSTITUTE OF TECHNOLOGY

© 2010 Scientific American

[FUTURE WORK]

WATCHING BIOLOGY'S CLOCKWORK

By adapting a technique called cryoimaging, researchers plan to do four-dimensional microscopy of biological processes such as proteins folding. A glassy (noncrystalline) ice will hold the protein. For each shot of the movie, a laser pulse will melt the ice around the sample, causing the protein to unfold in the warm water. The movie will record the protein refolding before the water cools and refreezes. The protein could be anchored to the substrate to keep it in the same position for each shot.



GEORGE RETSECK

protein crystals in this way. In the future we hope to watch proteins embedded in such vitreous water fold and unfold: a clocking pulse will boost the temperature enough to melt a tiny droplet of the water around the protein, which will unfold and then promptly refold. When the water cools and refreezes, it renders the molecule ready for another clocking pulse. The same approach could allow us to visualize the dynamics of bacterial flagella and of the fatty acid bilayers that make up cell membranes. As with our graphite studies, ultrafast electron energy loss spectroscopy should let us map changes in bonding. Capturing the image before the biosystem moves or disintegrates should provide sharper images than currently possible in cryomicroscopy.

Variants of ultrafast electron microscopy might well push below the nanoscale in structural dynamics studies and below a femtosecond in the imaging of the electron distribution in matter. Very recently, my Caltech group demonstrated two new techniques. In one, convergent-beam UEM, the electron pulse is focused and probes only a single nanoscopic site in a specimen. The other, near-field UEM, enables imaging of the evanescent electromagnetic waves ("plasmons") created in nanoscopic structures by an intense laser pulse—a phenomenon that underlies an exciting new technology known as plasmonics [see "The Promise of Plasmonics," by Harry A. Atwater; Scientific American, April 2007]. This technique has produced images of bacterial cell membranes and protein vesicles with femtosecond- and nanometer-scale resolution.

In recent years Ferenc Krausz of Ludwig Maximilian University of Munich, Paul Corkum of the University of Ottawa and others have opened up the attosecond regime to optical (light-based) studies using extremely short laser pulses. At Caltech, we have proposed several ultrafast electron microscopy schemes for attosecond-scale electron-based imaging, and we are now pursuing the experimental realization in collaboration with Herman Batelaan of the University of Nebraska–Lincoln.

The electron microscope is extraordinarily powerful and versatile. It can operate in three distinct domains: real-space images, diffraction patterns and energy spectra. It is used in applications ranging from materials and mineralogy to nanotechnology and biology, elucidating static structures in tremendous detail. By integrating the fourth dimension, we are turning still pictures into the movies needed to watch matter's behavior—from atoms to cells—unfolding in time. ■

MORE TO EXPLORE

A Revolution in Electron Microscopy. John M. Thomas in *Angewandte Chemie International Edition,* Vol. 44, No. 35, pages 5563–5566; September 5, 2005.

Microscopy: Photons and Electrons Team Up. F. Javier García de Abajo in *Nature,* Vol. 462, page 861; December 17, 2009.

Four-Dimensional Electron Microscopy. Ahmed H. Zewail in *Science,* Vol. 328, pages 187–193; April 9, 2010.

Biological Imaging with 4D Ultrafast Electron Microscopy. David J. Flannigan, Brett Barwick and Ahmed H. Zewail in *Proceedings of the National Academy of Sciences USA*, Vol. 107, No. 22, pages 9933–9937; June 1, 2010.

4D Electron Microscopy: Imaging in Space and Time. Ahmed H. Zewail and John M. Thomas. Imperial College Press, 2010.

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

© 2010 Scientific American

The Hacker in Your Hardware

As if software viruses weren't bad enough, the microchips that power every aspect of our digital world are vulnerable to tampering in the factory. The consequences could be dire

BY JOHN VILLASENOR

KEY CONCEPTS

- Integrated circuits are increasingly complex and capable—but also increasingly vulnerable to attack.
- The circuits typically include designs from many sources. A "Trojan" attack hidden in one of these designs could surface long after the circuit has left the factory.
- A few relatively simple measures could go a long way toward protecting hardware from malicious hackers.

—The Editors

YOUR ONCE RELIABLE MOBILE phone suddenly freezes. The keypad no longer functions, and it cannot make or receive calls or text messages. You try to power off, but nothing happens. You remove the battery and reinsert it; the phone simply returns to its frozen state. Clearly, this is no ordinary glitch. Hours later you learn that yours is not an isolated problem: millions of other people also saw their phones suddenly, inexplicably, freeze.

This is one possible way that we might experience a large-scale hardware attack—one that is rooted in the increasingly sophisticated integrated circuits that serve as the brains of many of the devices we rely on every day. These circuits have become so complex that no single set of engineers can understand every piece of their design; instead teams of engineers on far-flung continents design parts of the chip, and it all comes together for the first time when the chip is printed onto silicon. The circuitry is so complex that exhaustive testing is impossible. Any bug placed in the chip's code will go unnoticed until it is activated by some sort of trigger, such as a specific date and time—like the Trojan horse, it initiates its attack after it is safely inside the guts of the hardware.

The physical nature of hardware attacks makes them potentially more problematic than worms, viruses and other malicious software. A virus can jump from machine to machine, but it can also in principle be wiped clean from any system it infects. In contrast, there is no fix for a hardware attack short of replacing the infected units. At least, not yet.

The difficulty of fixing a systemic, malicious hardware problem keeps cybersecurity experts up at night. Anything that uses a microprocessor—which is to say, just about everything electronic—is vulnerable. Integrated circuits lie at the heart of our communications systems and the world's electricity supply. They position the

KENN BROWN *Mondolithic Studios*

© 2010 Scientific American



© 2010 Scientific American

flaps on modern airliners and modulate the power in your car's antilock braking system. They are used to access bank vaults and ATMs and to run the stock market. They form the core of almost every critical system in use by our armed forces. A well-designed attack could conceivably bring commerce to a halt or immobilize critical parts of our military or government.

Because Trojan hardware can hide for years before it is activated, it is possible—perhaps likely—that hardware bugs have already been planted. And although no large-scale hardware attacks have yet been confirmed, they are inevitable.

As we know all too well from combating software-based cyberattacks, a relatively small proportion of people who use their technical skills for malicious purposes can have an big impact. Thus, rather than asking whether or not hardware attacks will occur, the better questions are: What forms will these attacks take? What consequences will they have? And, perhaps most important of all, what can we do to detect and stop them or at least minimize their effects?

BLOCK BY BLOCK

AN INTEGRATED CIRCUIT, or chip, is simply an electronic circuit etched onto a single piece of a semiconductor material, most often silicon. Modern integrated circuits are physically quite small—no more than a few square centimeters and often much smaller—but can contain several billion transistors. The very complexity of modern chips creates the vulnerabilities that make Trojan attacks possible.

Modern chips are divided into subunits called blocks that perform different functions. In a mobile phone's processor, for example, one block might be memory that can be used to store frames of video captured by the camera. A second block might compress that video into an MPEG file, and a third block might convert those files into a form that can be transmitted over the antenna. Data move among these blocks across a system bus, which acts like a highway connecting the different parts of the chip.

When a company embarks on the design of a new integrated circuit, it first maps out what functional blocks the circuit will need. Some of these blocks will be designed in-house, either from scratch or as a modification of a block design used in the company's earlier chips. Others will be licensed from third parties that might specialize in a certain type of functionality—receiving data from an antenna, for example.

The block from the third party does not come as a physical piece of silicon, because the goal in building the integrated circuit is to have all the functional blocks printed onto the same surface. Instead the block comes as a data file that fully describes how the block should be etched onto the silicon. The file can be thousands of lines long, making it a practical impossibility for a human to read the file and understand everything that is going on. The block provider will also typically supply some software that the block purchaser uses to model how the block will respond to a variety of situations. Before any circuits are printed, the lead company will join all the model blocks into a computer simulation to ensure the chip will function as expected. Only when the model passes a battery of tests will the company begin the time-consuming and expensive process of fabricating the physical integrated circuits.

Here is where the vulnerability lies: because the rogue hardware requires a specific trigger to become active, chipmakers will have to test

[CIRCUIT VULNERABILITIES]

GLOBAL SECURITY PROBLEM

A single chip can incorporate circuits designed in locations across the globe by hundreds of people working at many different firms. This globalization of chip design makes new product development faster and less expensive. It also creates risks, because it is difficult to detect rogue circuitry hidden among the hundreds of millions of transistors before the chip ships.

▼ **INSIDE AN INTEGRATED CIRCUIT:** A chip contains a set of functional blocks, each dedicated to performing a specific task. Data move from block to block across a system bus, and traffic flow in the bus is controlled by yet another block, the bus arbiter. In a cell phone, for instance, data might travel from memory (*1*) to a block that performs computations (*2*) to a block that encodes and decodes information (*3*) to blocks that exchange data with off-chip locations (*4*) and (*5*).



HARDWARE DESIGN BY THE NUMBERS

1,550
Estimated number of companies worldwide involved in integrated-circuit design, including 700 in North America, 600 in Asia, and 250 in Europe, the Middle East and Africa

2,500
The approximate number of new chips designed every year

$235 billion
Global semiconductor sales revenue in 2009

QUICKHONEY

© 2010 Scientific American

▼ **WHEN THINGS GO WRONG:** A chip containing a Trojan will function normally until the rogue hardware is awakened by a trigger and instructed to attack. Triggers could take many possible forms, including the arrival of a certain date and time or a "wake-up call" from the outside world arriving as a specially encoded packet of data. The Trojan can then execute one of two types of attacks—overt and covert.

In an overt attack (*below*), the rogue hardware stops the chip from functioning properly. In this example, a corrupted block refuses to release its access to the system bus, thereby preventing other blocks from communicating with one another. In this case, the chip would cease to function altogether.

In a covert attack (*right*), the rogue circuitry does not give any indication of its presence. Covert attacks are of particular concern because there is no obvious evidence that anything is wrong. Underneath, however, the rogue circuitry can be sending confidential data to an off-chip location or working in concert with other compromised systems to launch other attacks.





their models against every possible trigger to ensure that the hardware is clean. This is simply not possible—the universe of possible triggers is far too large. In addition to internal triggers such as a date-based trigger described in the mobile phone example, hackers could employ external triggers such as the reception of a text or e-mail message containing a specific set of characters. Companies test as best as they can, even though this necessarily means testing only a very small percentage of possible inputs. If a block behaves as expected, it is assumed to be functioning correctly.

AN ISSUE OF TRUST

IN THE EARLY DAYS of integrated-circuit design, no one had to worry about hackers. The first designs were created completely in-house, executed by small teams who were working toward a common purpose. Because of this organizational security, the designers established open protocols that assumed different parts of the chip would behave as expected. (The history echoes the choices that were made in the early days of the Internet, when a small academic community built an open platform that assumed everyone would behave nicely. That assumption has not withstood the growth of the Internet.)

In today's world, however, the design process for a single, large integrated circuit can involve contributions from hundreds or even thousands of people at locations on multiple continents. As this design goes through various stages of development, portions of the design are stored on many different physical platforms and repeatedly exchanged among many parties. For example, an American manufacturer might combine designs from separate branches of the company with designs from third-party vendors in the U.S., Europe and India, then fabricate the chip in a Chinese factory. These global networks have become a fact of life in recent years, and they have provided large savings in cost and efficiency. But they make security far more complicated than back in the days when things were done in one facility. Given the sheer number of people and complexity involved in a large integrated-circuit design, there is always a risk that an unauthorized outsider might gain access and corrupt the design without detection.

A very small—but not zero—risk also exists that a design could be corrupted by someone with internal access. While the overwhelming majority of people involved in any aspect of circuit design will endeavor to deliver designs of the highest quality, as with any security issue, malicious actions taken by even a very small minority of those with inside access acting maliciously can create significant problems.

MODERN MIND: The AMD Opteron 6100 processor, shown here at actual size, is a package of two integrated circuts containing a total of more than 1.8 billion transistors. The various colors correspond to different functional blocks.



COURTESY OF AMD

© 2010 Scientific American

HOW TO STOP HARDWARE HACKERS

A secure integrated circuit contains a modest amount of extra hardware that polices the chip from the inside. When hostile behavior is detected, a set of security measures can spring into place within microseconds to identify the source of the attack and react with countermeasures to surmount it. Here are a few strategies being developed.

SECURITY MEASURE	INTENDED EFFECT	HOW IT WORKS
Memory gatekeeper	Preempts any attempt by a rogue block to access off-limits areas of memory; the measure helps to prevent spying or data corruption	A gatekeeper ensures that blocks can access only authorized portions of memory, and flags any attempts at unauthorized access
Secure system bus	Prevents a malicious takeover of a system bus, which could result in a complete circuit shutdown or dramatically slowed operation	The bus analyzes statistical patterns of bus access by different functional blocks and flags strange behavior
Input/output monitor	Impedes covert spying—when the chip attempts to copy data to off-chip locations	The circuit analyzes the movement of data on and off the chip and compares this flow with the expected pattern, flagging any aberrations
On-site block testing	Defends against a Trojan attack that attempts to damage a formerly "healthy" block	An on-chip integrity checker occasionally tests the blocks to make sure they continue to operate as expected
Extra programmable hardware logic	Allows the circuit to quarantine a compromised block and replicate its function	The extra logic is configured to replace the quarantined functional block—though likely at a slower speed
Attack alert system	Allows other circuits to preemptively protect themselves against impending attacks	A circuit under attack institutes countermeasures and sends a warning about the attack to other devices containing the same circuit

Ideally, would-be attackers would never get the opportunity to gain access to an integrated circuit during the design and manufacturing process, thereby ensuring that hardware attacks never occur. This is the strategy that the Defense Advanced Research Projects Agency (DARPA), the research agency run by the Pentagon, has pursued with its Trust in Integrated Circuits program. DARPA is designing processes to ensure that all the steps in the design and manufacturing chain are carried out by companies and people known to be trustworthy and working in secure environments. (In addition, the agency is funding research into new ways to test chips before they are placed into U.S. weapons systems.) Yet in the real world, actions taken to secure the design process are never perfect.

Hardware designers should also build circuits that identify and respond to attacks even as they are taking place, like an onboard police force. Although a community should certainly engage in all reasonable measures to discourage potential criminals from committing crimes, any responsible community also recognizes that such efforts, no matter how well intentioned and thorough, will never be 100 percent effective. It is critical to have a police force that can respond quickly and appropriately when crimes do occur.

SECURING THE CIRCUIT

A CIRCUIT THAT can effectively detect and respond to attacks is called a secure circuit. These chips have a modest amount of extra circuitry specifically designed to look for behavior that may reveal a problem. If an attack is suspected, the secure circuit will identify the type of attack and attempt to minimize the resulting damage.

In the example of the frozen cell phone, the failure may have been caused by a single block that was acting out of order. That block interacts with all the other blocks over the system bus. This bus, in turn, has a bus arbiter—a traffic cop that decides what information can travel over the bus at what time. Yet the traffic cop analogy is not perfect. While a traffic cop can instruct traffic when to start and when to stop, a bus arbiter has less authority. It can grant permission for a block to start sending information through the bus, but the block can retain that access for as long as it wants—a vestige of the long-ago assumption that blocks would always behave properly. Herein lies the problem.

In a typical system, a block will retain access to the system bus for only as long as necessary before it relinquishes it for use by other blocks.

QUICKHONEY

© 2010 Scientific American

The bus arbiter sees that the system bus is available and then assigns it to another block. But if a block keeps control of the bus indefinitely, no further data will be able move within the integrated circuit, and the system will freeze.

In contrast, a secure integrated circuit performs constant checks to ensure that the communications among different blocks have not been disrupted. When it detects one block monopolizing access to the bus, the secure integrated circuit can respond by quarantining the malicious block. It can then use its store of programmable logic hardware to replace the lost functionality. This process will likely slow the overall operation, but it will at least keep the device working.

An overt attack is probably not the most pernicious threat, however. A covert attack could be much worse. In a covert attack, the device appears to operate normally, but in reality it is acting with malicious intent. A mobile phone, for instance, might secretly begin to transmit a copy of all incoming and outgoing text messages to a third party. An unsuspecting observer would not notice anything wrong, and the attack could continue indefinitely.

A secure integrated circuit would provide a critically important defense against this type of attack. The chip would constantly monitor the amount and type of data moving on and off the integrated circuit and statistically compare this movement with the expected data flows. Any anomaly would be flagged as a potential data leak, and the chip would either alert the user or begin to staunch the flow on its own.

In addition to taking steps to counter the effects of a Trojan attack on its own operations, an integrated circuit can notify other devices of the type of assault, potentially allowing them to take preemptive actions to avoid it (or at least to minimize its effects). Such notification is not as far-fetched as it might seem given the level of network connectivity that almost all systems now have. For example, if a circuit experiencing an attack can identify the initiating trigger, it can alert other circuits to screen for that particular message.

The measures described here will be effective only if the parts of the circuit responsible for managing security are themselves secure and trustworthy. This might seem like a circular argument—another way of saying that the only way to secure a circuit is to secure a circuit—but the elements of the circuit devoted to security constitute only a small fraction of the overall design. They can be designed in-house to ensure that only highly trusted parties have access.

WHAT TO DO NEXT

THANKS TO the efforts of governments, academic researchers and the commercial sector, enormous progress has been made in Internet security. The same cannot be said of the state of integrated-circuit security, which lies roughly where Internet security was 15 years ago: there is growing awareness that the issue is worthy of attention, but defensive strategies have not yet been fully developed, much less put into practice.

A comprehensive approach to preventing hardware attacks requires action on several levels. Strategies that aim to ensure compromised hardware never gets out the door, such as DARPA's program, are a good start. But most important, we must begin to implement secure design measures such as the ones discussed here that can defend against attacks as they occur. These defenses will not come free. As with security in other domains, integrated-circuit security will require the expense of time, money and effort. A wide spectrum of options represents trade-offs between the effectiveness of the security and the cost of implementing it. Fortunately, it is possible to deliver effective security at modest costs.

A secure integrated circuit contains a small amount of extra logic. In research my group has conducted at the University of California, Los Angeles, we have found that the increase in integrated-circuit size is typically several percent. There is also generally a cost in operating speed, given that the steps taken to ensure that functional blocks are behaving appropriately can consume some clock cycles that might otherwise be used for core operational tasks. Again, however, we have found the speed reduction to be small in relative terms, and in some cases no speed reduction happens at all if the security measures are performed using logic and functional blocks that are temporarily dormant.

Keeping hardware secure will inevitably become an arms race requiring continual innovation to stay ahead of the latest attacks, as has been the case in the software world. Whereas new circuits cannot be downloaded over the Internet in the manner used to fix security holes identified in software, modern integrated circuits have a number of reconfigurable aspects that, with appropriate steps taken during the integrated-circuit design process, could be used to automatically replace parts of hardware that become incapacitated in the event of an attack. Engineered flexibility is our best defense.

Even if hardware attacks are inevitable, that does not mean that they have to be successful. ■

[THE AUTHOR]

JOHN VILLASENOR joined the electrical engineering department at the University of California, Los Angeles, in 1992. Prior to that he worked on developing methods to image the earth from space at the NASA Jet Propulsion Laboratory and completed an M.S. and Ph.D. at Stanford University. At U.C.L.A. his research focuses on the methods, technologies and systems used to capture information in the world around us, convert it into digital form, and move it efficiently and securely from one place to another.

MORE TO EXPLORE

Trojan Detection Using IC Fingerprinting. D. Agrawal et al. in *Proceedings of the 2007 IEEE Symposium on Security and Privacy.* Berkeley, Calif., May 2007.

Old Trick Threatens the Newest Weapons. John Markoff in *New York Times;* October 26, 2009.

A Trojan-Resistant System-on-Chip Bus Architecture. L. Kim, J. Villasenor and C. Koc in *Proceedings of the 2009 IEEE Military Communications Conference.* Boston, October 2009.

Securing the Information Highway. Wesley K. Clark and Peter L. Levin in *Foreign Affairs,* November/December 2009.

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

COURTESY OF RORY EARNSHAW

© 2010 Scientific American

Plastic Surf

Small remnants of toys, bottles and packaging have an unhealthful afterlife in the ocean

BY JENNIFER ACKERMAN ■ PHOTOGRAPH BY CARY WOLINSKY

SIREN CALL FOR BIRDS
Seabirds are known to pick up plastic fragments, especially red ones, mistaking them for food at the sea surface. Even in remote areas, scientists are finding birds, such as Antarctic prions and sub-Antarctic skuas, that have ingested plastic marine debris.

LIKE ROCKS INTO SAND
The same mechanical action of waves against shoreline that turns rocks into sand can smooth and round the edges of plastics and help to grind them into tiny fragments and even powder, but they do not disappear.

BY NOW EVEN SCHOOLCHILDREN know that the plastics we discard every year in the millions of tons persist in the environment for hundreds of years. And we have all heard of the horrors caused by such debris in the sea: fur seals entangled by nylon nets, sea otters choking on polyethylene six-pack rings, and plastic bags or toys stuck in the guts of sea turtles. This photograph, showing plastic fragments collected in just an hour at a cove near Gloucester, Mass., hints at a lesser-known but equally disturbing story: much smaller bits of plastic that are accumulating in oceans all over the world can potentially harm marine life and possibly even human health.

Although plastic does not get digested by microbes, as food and paper are, it does slowly "photodegrade": ultraviolet light and heat from the sun increase its brittleness, causing it to weaken, crack and break up into smaller and smaller fragments. Indeed, a handful of sand or cup of seawater from nearly anywhere in the world will probably be peppered with microplastics—pieces that are tinier than a small pea and often invisible. Scientists fear the possible effects of this plastic confetti on zooplankton and other creatures at the base of the marine food web, which get consumed by larger organisms—turtles, fish, birds—and, ultimately, by us.

The bits evoke worry for several reasons. They may block the feeding appendages or digestive tracts of small invertebrates. Also, chemicals incorporated into plastic products during their manufacture can threaten health when ingested—among them the endocrine disruptor bisphenol A. Moreover, plastic debris floating in the ocean acts as a magnet and sponge for toxic substances, such as DDT, dioxins and PCBs, absorbing concentrations 100 to a million times that of surrounding seawater. Life-forms eating pollutant-loaded microplastics may therefore be exposing themselves—and those who feed on them—to concentrated poisons. A solution, many scientists agree, is designing plastic products that can be recycled throughout much of their material life. ■

Jennifer Ackerman is a science writer based in Virginia.

Comment on this article at **www.ScientificAmerican.com/sciammag/aug2010**

© 2010 Scientific American

READY TO CRACK
Scientists have identified at least nine types of plastic waste in the oceans, from acrylic and nylon to polyester, polypropylene (found in ropes and containers), polycarbonate (the hard plastic in eyeglass lenses) and polystyrene (used in making Styrofoam). Polystyrene begins to break into smaller pieces within a year, and even hard plastics such as polycarbonate can come apart in the sea, potentially leaking chemicals as they do.

© 2010 Scientific American

Arctic Change ▪ Malaria's Grip ▪ Periodic Table

BY KATE WONG

THE CHANGING ARCTIC LANDSCAPE
by Ken D. Tape. University of Alaska Press, 2010 ($35)

Photographer Ken D. Tape pairs old and new images of sites in northern Alaska to document the impact of climate change on the Arctic.

BROOKS RANGE, 1973

BROOKS RANGE, 2007

EXCERPT

THE FEVER: HOW MALARIA HAS RULED HUMANKIND FOR 500,000 YEARS
by Sonia Shah. Farrar, Straus and Giroux, 2010 ($26)

Every year malaria infects more than 500 million people; every 30 seconds someone dies from it. Journalist Sonia Shah examines the various factors that have allowed the disease to flourish despite scientists' best efforts to combat it. Where malaria is endemic, many people, Shah says, do not accept that the disease is transmitted solely by mosquitoes, or else they consider it largely benign. Here she describes how Malawi's dominant ethnic group, the Chewa, view it.

"For us, malaria is a disease caused by a protozoan parasite transmitted by mosquito. For the Lake Malombe Chewa, malaria—which the locals call *malungo,* and lump together with other malaria-like illnesses—is a disease caused by mosquitoes … and spirits and jealousy and hexes and bad weather and hard work and dirty water and rotten food, among other things.…

"Like intelligent design and other forms of magical thinking, these beliefs are not unrelated to actual shortcomings in the scientific explanations with which they compete. Every time mosquitoes bit Lake Malombe Chewa and they did not fall ill with *malungo*, their disbelief in the mosquito theory of malaria transmission strengthened. Ditto for every time they took an antimalarial drug and it failed to work. If the drug didn't work, this meant that the *malungo* was not caused by mosquitoes.

"What these beliefs mean is that while our malaria is an eminently preventable disease, for the Chewa, as for other rural peoples living traditional lives, it is anything but. Malaria is everywhere, caused by everything.…

"It isn't that the Chewa villagers don't understand that destroying mosquitoes' larval habitats, or sleeping under bed nets, or taking prophylactic drugs, or sealing up their houses, helps prevent malaria. And it isn't that they aren't interested in preventing malaria.… It is that, as with people everywhere, there's little interest in fixes that are time-consuming or temporary, or that promise only—in their minds—marginal efficacy. Even if some *malungo* can be alleviated by people avoiding mosquito bites, they can't possibly avoid exposure to the weather, or to hard work, or to the envy of their neighbors."

ALSO NOTABLE

- **The Disappearing Spoon: And Other True Tales of Madness, Love, and the History of the World from the Periodic Table of the Elements**
  by Sam Kean. Little, Brown, 2010 ($24.99)
- **The Book of Shells: A Life-Size Guide to Identifying and Classifying Six Hundred Seashells**
  by M. G. Harasewych and Fabio Moretzsohn. University of Chicago Press, 2010 ($55)
- **Voyager: Seeking Newer Worlds in the Third Great Age of Discovery**
  by Stephen J. Pyne. Viking, 2010 ($29.95)
- **The Weather of the Future: Heat Waves, Extreme Storms, and Other Scenes from a Climate-Changed Planet**
  by Heidi Cullen. HarperCollins, 2010 ($19.99)
- **Here's Looking at Euclid: A Surprising Excursion through the Astonishing World of Math**
  by Alex Bellos. Free Press, 2010 ($25)
- **The Artificial Ape: How Technology Changed the Course of Human Evolution**
  by Timothy Taylor. Palgrave Macmillan, 2010 ($27)
- **The Youth Pill: Scientists at the Brink of an Anti-Aging Revolution**
  by David Stipp. Current, 2010 ($26.95)
- **Ancient Bodies, Modern Lives: How Evolution Has Shaped Women's Health**
  by Wenda Trevathan. Oxford University Press, 2010 ($34.95)
- **The Zodiac of Paris: How an Improbable Controversy over an Ancient Egyptian Artifact Provoked a Modern Debate between Religion and Science**
  by Jed Z. Buchwald and Diane Greco Josefowicz. Princeton University Press, 2010 ($35)

DENNIS TRABANT (*left*); KEN D. TAPE (*right*)

© 2010 Scientific American

"As the master craftsman who opened the famous Lincoln Pocket Watch in Washington, D.C., I recently reviewed the Stauer Meisterzeit timepiece. The assembly and the precision of the mechanical movement are the best in its class."

—George Thomas
Towson Watch Company

Authentic Stauer Historical Reproductions

We Can Only Find One

A rare chance to claim a unique piece of watchmaking history for under $100!

Eighty-six years ago, a watchmaker in Paris famous for building the magnificent clocks at Versailles created a legendary timepiece. He invented the first watch with an automatic mechanical drive. These innovative movements required no batteries and never needed to be manually wound. Only seven of these ultra-rare watches were ever made and we've studied the one surviving masterpiece in a watch history museum. Inspired by history, classic design and technology, our Stauer ***Meisterzeit*** has been painstakingly handcrafted to meet the demanding standards of vintage watch collectors.

Why the new "antique" is better than the original. The original timepiece was truly innovative, but, as we studied it closely, we realized that we could engineer ours with a much higher level of precision. The 27-ruby-jewel movement utilizes an automatic self-winding mechanism inspired by a patent from 1923, but built on $31 million in state-of-the-art Swiss-made machinery. With an exhibition back, you can see into the heart of the engineering and view the rotor spin—it's powered by the movement of your body.

This limited edition Stauer ***Meisterzeit*** allows you to wear a watch far more exclusive than most new "upscale" models. Here is your chance to claim a piece of watchmaking history in a rare design that is priced to wear everyday.

Elegant and accurate. This refined beauty has a fastidious side. Each movement and engine-turned rotor is tested for 15 days and then certified before it leaves the factory.

The best part is that with our special price, you can wear a superb classic historical reproduction watch and laugh all the way to the bank. Stauer specializes in classic timeless watches and jewelry that are made for the millionaires who want to keep their millions. This watch will quickly move to heirloom status in your household.

View the precision movement of the *Meisterzeit* through the rear exhibition port.

Try it for 30 days and if you are not thrilled with the beauty and construction of the ***Meisterzeit***, simply return it for a refund of the purchase price.

Extremely limited availability. Since it takes about 6 months to build each ***Meisterzeit***, the release is a limited edition, so please be sure to order yours soon.

WATCH SPECS:
- 18K Gold-clad case and bezel
- Precision 27-jeweled movement
- Interior dials display day and month
- Croc-embossed leather strap
- Fits 6 3/4"–8 1/4" wrist

76% OFF

Exclusively Through Stauer

Stauer Meisterzeit Watch—~~$395~~
Now **only** ***$95*** +S&P ***Save $300!***

Call now to take advantage of this limited offer.

1-888-324-4351

Promotional Code MZW297-02
Please mention this code when you call.

Stauer
14101 Southcross Drive W.,
Dept. MZW297-02
Burnsville, Minnesota 55337
www.stauer.com

Smart Luxuries—Surprising Prices

Take Me Out of the Ball Game

Physics and medicine are the biggest players on the diamond

BY STEVE MIRSKY

Baseball is a game of trajectories. And as Yogi Berra supposedly said, you can observe a lot just by watching. For example, at Yankee Stadium on May 29, I observed New York slugger Alex Rodriguez hit a pitch by Cleveland Indians David Huff back up the middle and off the pitcher's head. In fact, the ball hit Huff's head so hard that it rolled nearly all the way to the right-field wall. The ball, that is, not Huff's head. He collapsed in a heap and remained face down on the mound for several minutes. Huff eventually left on a stretcher. Home team fans who then watched the Yankees blow a six-run lead left in a huff.

Anyway, many in the crowd feared that Huff was seriously injured. Having observed physics teachers years earlier, however, I was guardedly optimistic—precisely because the ball had ricocheted so far and so fast. Had the ball rebounded from Huff's dome only a few feet straight back toward home plate, I would have been concerned that the poor pitcher had become the second player in major league baseball history to be killed on the job. In that scenario, much of the ball's energy of motion would have been imparted to the pitcher. But said energy appeared to have been expended on sending the ball skittering into the right-field corner, with only a small amount having been transferred to Huff's head.

Indeed, after being checked out at nearby New York–Presbyterian Hospital, Huff returned to the scene of the bean, seemingly little the worse for wear, before the game was even over. He was helped in this effort by the fact that the 13–11 Indians win took an excruciating four hours and 22 minutes, which felt much longer for those of us near the event horizon.

(The only man killed playing major league baseball is Cleveland Indians shortstop Ray Chapman, who, after being struck in the temple by a pitch from Carl Mays of the Yankees in 1920, never regained consciousness. A scant 51 years later the powers-that-were made protective headgear mandatory.)

On the same afternoon as A-Rod's double off Broca's area, a far more serious injury was sustained by Kendry Morales, first baseman for a team somehow seriously referred to as the Los Angeles Angels of Anaheim. Morales hit a game-winning grand-slam home run in the bottom of the ninth, trotted around the bases, leapt onto home plate and shattered his left ankle. He'll need surgery and could miss the rest of the season. Morales is still lucky compared with other world-class athletes that after similar injuries used to be taken out back and shot and more recently are euthanized by large-animal veterinarians.

That same evening Philadelphia Phillies pitcher Roy Halladay pitched the 20th perfect game in major league history, retiring all 27 Florida Marlins who futilely waved their tribute sticks, I mean bats, at him. The *Philadelphia Inquirer*'s Bill Lyon wrote that Halladay "made the ball dive, and he made it rise."

Lyon is hitting .500. No pitcher throwing overhand can really make the ball rise, says University of New Hampshire psychology professor Kenneth Fuld, who has taught a course called Visual Perception. (New Hampshire was where I came closest to playing professional baseball, when I was given room and board for a week so I could be a ringer on the Dartmouth College math department softball team. Given sufficient initial conditions, those guys could tell you exactly where an opponent's batted ball would land when, in keeping with math department tradition, they didn't catch it.)

According to Fuld, whose son, Sam, is an outfielder on the Chicago Cubs' 40-man roster, straight fastballs delivered overhand always drop slightly on their way to home plate. But the batter perceives the trajectory as level. An unusually fast pitch, which drops merely less than expected, will then appear to rise. It's an illusion, like when you think your stationary train is moving because a train on the next track moves or when paying six bucks for a hot dog at Yankee Stadium doesn't seem at all crazy, because you already shelled out $125 for your seat. ■

PHOTOGRAPH BY FLYNN LARSEN; ILLUSTRATION BY MATT COLLINS

© 2010 Scientific American

If 41 city mpg doesn't charge you up, its battery will.

MOTOR TREND'S 2010 CAR OF THE YEAR.*

THE FORD **FUSION HYBRID**

Our next-generation battery pack helps Fusion Hybrid deliver over 700 miles on a single tank, for up to 41 city mpg. All of which make Fusion Hybrid the most fuel-efficient midsize sedan in America.*

fordvehicles.com

Ford

Drive one.

*2010 EPA-estimated 41 city/36 hwy/39 combined mpg. 17.5-gallon tank. Actual mileage will vary. Midsize class per R. L. Polk & Co.

Advanced adaptive optics help create a sharper, truer view of the inside of your eye...

Reduced distortion, higher resolution

For years, researchers have examined the *fundus* or inner lining of the eye by looking through the pupil with an ophthalmoscope. However, the eye's cornea and lens have always *distorted* their view. And early techniques for correcting the distortion were too costly and low in resolution...



Adaptive Optics Scanning Laser Ophthalmoscope using LCOS-SLM

So in 2005, Hamamatsu started working on a better solution. The answer was *Liquid Crystal on Silicon Spatial Light Modulator* technology (LCOS-SLM). It applies controlled low voltages across a special liquid crystal mirror to dynamically alter the refractive index at each pixel and precisely correct wavefront distortions.

Hamamatsu is opening the new frontiers of Light

In the future, this technology may enable new generations of lower-cost, high-resolution fundus imaging. Such as scanning laser ophthalmoscopes that can clearly focus on individual visual receptor cells and microscopic blood vessels.



Get focusable views of visual receptor cells, blood vessels and nerve fibers (left to right).

Or the technology may help to make earlier diagnosis of eye and circulatory system diseases. It's just one more way Hamamatsu is opening the new frontiers of light — to brighten our world.

http://jp.hamamatsu.com/en/rd/publication/

HAMAMATSU

Photon is Our Business

Hamamatsu's work on this technology was conducted under a research project of the New Energy and Industrial Technology Development Organization of Japan (NEDO). The adaptive optics scanning laser ophthalmoscope is being developed and reviewed with cooperation from NIDEK Co., Ltd. and the Faculty of Medicine at Kyoto University under this project.